

Table of Contents



Basic Trig Functions



$$\sin(\theta) = \frac{\text{opp}}{\text{hyp}}$$

$$\cos(\theta) = \frac{\text{adj}}{\text{hyp}}$$

$$\tan(\theta) = \frac{\text{opp}}{\text{adj}}$$

$$\csc(\theta) = \frac{1}{\sin(\theta)}$$

$$\sec(\theta) = \frac{1}{\cos(\theta)}$$


$$\cot(\theta) = \frac{1}{\tan(\theta)}$$

COMMONLY USED FACTS


- Graphs of sine, cosine, tangent
- Sine, cosine, and tangent of reference angles: 0, $\frac{\pi}{6}$, $\frac{\pi}{4}$, $\frac{\pi}{3}$, $\frac{\pi}{2}$
- How to use reference angles to find sine, cosine and tangent of other angles
- Identities: $\sin^2 x + \cos^2 x = 1$; $\tan^2 x + 1 = \sec^2 x$; $\sin^2 x = \frac{1 - \cos(2x)}{2}$; $\cos^2 x = \frac{1 + \cos 2x}{2}$
- Conversion between radians and degrees

CLP-1 has an appendix on high school trigonometry that you should be familiar with.

Reference Angles



Derivative of Sine



Consider the derivative of $f(x) = \sin(x)$.

$$\frac{\mathrm{d}}{\mathrm{d}x}\{\sin x\} = \lim_{h\to 0}\frac{\sin(x+h)-\sin(x)}{h}$$


$$= \lim_{h\to 0}\frac{\sin(x)\cos(h)+\cos(x)\sin(h)-\sin(x)}{h}$$

$$= \lim_{h\to 0}\frac{\sin(x)(\cos(h)-1)}{h} + \lim_{h\to 0}\frac{\cos(x)\sin(h)}{h}$$

$$= \sin(x)\lim_{h\to 0}\frac{\cos(0+h)-\cos(0)}{h} + \cos(x)\lim_{h\to 0}\frac{\sin(h)}{h}$$

$$= \sin(x)\,\frac{\mathrm{d}}{\mathrm{d}x}\{\cos(x)\}\Big|_{x=0} + \cos(x)\lim_{h\to 0}\frac{\sin(h)}{h} \qquad = \qquad \boxed{\cos(x)\lim_{h\to 0}\frac{\sin(h)}{h}}$$

since $\cos(x)$ has a horizontal tangent, and hence has derivative zero, at $x = 0$.


1
h
sin(h)
h
tan(h)


Derivatives of Sine and Cosine

From before,

$$\frac{\mathrm{d}}{\mathrm{d}x}\{\sin(x)\} = \cos(x) \cdot \lim_{h \to 0} \frac{\sin(h)}{h} = \cos(x)$$

Derivative of Cosine

Now for the derivative of cos. We already know the derivative of sin, and it is easy to convert between sin and cos using trig identities.



$$\sin x = \frac{b}{c} = \cos\left(\frac{\pi}{2} - x\right)$$
$$\cos x = \frac{a}{c} = \sin\left(\frac{\pi}{2} - x\right)$$

When we use radians:

Derivatives of Trig Functions

$\frac{d}{dx}\{\sin(x)\} = \cos(x)$

$\frac{d}{dx}\{\cos(x)\} = -\sin(x)$

$\frac{d}{dx}\{\tan(x)\} =$


$\frac{d}{dx}\{\sec(x)\} =$

$\frac{d}{dx}\{\csc(x)\} =$


$\frac{d}{dx}\{\cot(x)\} =$

Honorable Mention

$$\lim_{x \to 0} \frac{\sin x}{x} = 1$$



$y = \sin x$, radians



$y = \sin x$, degrees

Other Trig Functions

$$\tan(x) = \frac{\sin(x)}{\cos(x)}$$

Other Trig Functions

$$\sec(x) = \frac{1}{\cos(x)}$$

$$\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}x}[\sec(x)] &= \frac{\mathrm{d}}{\mathrm{d}x}\left[\frac{1}{\cos(x)}\right] \\
&= \frac{\cos(x)(0) - (1)(-\sin(x))}{\cos^2(x)} \\
&= \frac{\sin(x)}{\cos^2(x)} \\
&= \frac{1}{\cos(x)}\frac{\sin(x)}{\cos(x)} \\
&= \sec(x)\tan(x)
\end{aligned}$$

Other Trig Functions

$$\csc(x) = \frac{1}{\sin(x)}$$

$$\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}x}[\csc(x)] &= \frac{\mathrm{d}}{\mathrm{d}x}\left[\frac{1}{\sin(x)}\right] \\
&= \frac{\sin(x)(0) - (1)\cos(x)}{\sin^2(x)} \\
&= \frac{-\cos(x)}{\sin^2(x)} \\
&= \frac{-1}{\sin(x)}\frac{\cos(x)}{\sin(x)} \\
&= -\csc(x)\cot(x)
\end{aligned}$$

Other Trig Functions

$$\cot(x) = \frac{\cos(x)}{\sin(x)}$$

$$\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}x}[\cot(x)] &= \frac{\mathrm{d}}{\mathrm{d}x}\left[\frac{\cos(x)}{\sin(x)}\right] \\
&= \frac{\sin(x)(-\sin(x)) - \cos(x)\cos(x)}{\sin^2(x)} \\
&= \frac{-1}{\sin^2(x)} \\
&= -\csc^2(x)
\end{aligned}$$

Memorize

$$\frac{\mathrm{d}}{\mathrm{d}x}\{\sin(x)\} = \cos(x) \qquad \frac{\mathrm{d}}{\mathrm{d}x}\{\sec(x)\} = \sec(x)\tan(x)$$
$$\frac{\mathrm{d}}{\mathrm{d}x}\{\cos(x)\} = -\sin(x) \qquad \frac{\mathrm{d}}{\mathrm{d}x}\{\csc(x)\} = -\csc(x)\cot(x)$$
$$\frac{\mathrm{d}}{\mathrm{d}x}\{\tan(x)\} = \sec^2(x) \qquad \frac{\mathrm{d}}{\mathrm{d}x}\{\cot(x)\} = -\csc^2(x)$$

$$\lim_{x\to 0}\frac{\sin x}{x} = 1$$

Let $f(x) = \dfrac{x\tan(x^2+7)}{15e^x}$. Use the definition of the derivative to find $f'(0)$.

Differentiate $(e^x + \cot x)\left(5x^6 - \csc x\right)$.

Let $h(x) = \begin{cases} \frac{\sin x}{x} & , \quad x < 0 \\ \frac{ax+b}{\cos x} & , \quad x \geq 0 \end{cases}$

Which values of a and b make $h(x)$ continuous at $x = 0$?

Practice and Review


$$f(x) = \begin{cases} x^2 \cos\left(\frac{1}{x}\right) & , \quad x \neq 0 \\ 0 & , \quad x = 0 \end{cases}$$

Is $f(x)$ differentiable at $x = 0$?

$$g(x) = \begin{cases} e^{\frac{\sin x}{x}} & , \quad x < 0 \\ (x-a)^2 & , \quad x \geq 0 \end{cases}$$

What value(s) of a makes $g(x)$ continuous at $x = 0$?

A ladder 3 metres long rests against a vertical wall. Let θ be the angle between the top of the ladder and the wall, measured in radians, and let y be the height of the top of the ladder. If the ladder slides away from the wall, how fast does y change with respect to θ?
When is the top of the ladder sinking the fastest? The slowest?



Suppose a point in the plane that is r centimetres from the origin, at an angle of θ ($0 \leq \theta \leq \frac{\pi}{2}$), is rotated by $\pi/2$ radians. What is its new coordinate (x, y)? If the point rotates at a constant rate of a radians per second, when is the x coordinate changing fastest and slowest with respect to θ?

