
# Table of Contents



# Back to Limits!

$$\lim_{x\to\infty} \frac{x^2}{5} \qquad \lim_{x\to\infty} \frac{5}{x^2} \qquad \lim_{x\to 0} \frac{x^2}{5} \qquad \lim_{x\to 0} \frac{5}{x^2}$$



ans

# Back to Limits!

$$\lim_{x\to\infty} \frac{x^2}{5} = \infty \qquad \lim_{x\to\infty} \frac{5}{x^2} \qquad \lim_{x\to 0} \frac{x^2}{5} \qquad \lim_{x\to 0} \frac{5}{x^2}$$



ans

# Back to Limits!

$$\lim_{x\to\infty} \frac{x^2}{5} = \infty \qquad \lim_{x\to\infty} \frac{5}{x^2} = 0 \qquad \lim_{x\to 0} \frac{x^2}{5} \qquad \lim_{x\to 0} \frac{5}{x^2}$$



ans

# Back to Limits!

$$\lim_{x\to\infty} \frac{x^2}{5} = \infty \qquad \lim_{x\to\infty} \frac{5}{x^2} = 0 \qquad \lim_{x\to 0} \frac{x^2}{5} = 0 \qquad \lim_{x\to 0} \frac{5}{x^2}$$



ans

# Back to Limits!

$$\lim_{x\to\infty} \frac{x^2}{5} = \infty \qquad \lim_{x\to\infty} \frac{5}{x^2} = 0 \qquad \lim_{x\to 0} \frac{x^2}{5} = 0 \qquad \lim_{x\to 0} \frac{5}{x^2} = \infty$$

# Back to Limits!

$$\lim_{x\to\infty} \frac{x^2}{5} = \infty \qquad \lim_{x\to\infty} \frac{5}{x^2} = 0 \qquad \lim_{x\to 0} \frac{x^2}{5} = 0 \qquad \lim_{x\to 0} \frac{5}{x^2} = \infty$$

## Indeterminate Forms – Definition 3.7.1

Suppose $\lim_{x\to a} f(x) = \lim_{x\to a} g(x) = 0$. Then the limit

$$\lim_{x\to a} \frac{f(x)}{g(x)}$$

is an indeterminate form of the type $\frac{0}{0}$.
Suppose $\lim_{x\to a} F(x) = \lim_{x\to a} G(x) = \infty$ (or $-\infty$). Then the limit

$$\lim_{x\to a} \frac{F(x)}{G(x)}$$

is an indeterminate form of the type $\frac{\infty}{\infty}$.

# Back to Limits!

$$\lim_{x\to\infty} \frac{x^2}{5} = \infty \qquad \lim_{x\to\infty} \frac{5}{x^2} = 0 \qquad \lim_{x\to 0} \frac{x^2}{5} = 0 \qquad \lim_{x\to 0} \frac{5}{x^2} = \infty$$

## Indeterminate Forms – Definition 3.7.1

Suppose $\lim_{x\to a} f(x) = \lim_{x\to a} g(x) = 0$. Then the limit

$$\lim_{x\to a} \frac{f(x)}{g(x)}$$

is an indeterminate form of the type $\frac{0}{0}$.
Suppose $\lim_{x\to a} F(x) = \lim_{x\to a} G(x) = \infty$ (or $-\infty$). Then the limit

$$\lim_{x\to a} \frac{F(x)}{G(x)}$$

is an indeterminate form of the type $\frac{\infty}{\infty}$.

**When you see an indeterminate form, you need to do more work.**

# Indeterminate Forms

$$\lim_{x \to 5} \frac{x^2 - 3x - 10}{x - 5}$$

indeterminate form of the type $\frac{0}{0}$



ans

# Indeterminate Forms

$$\lim_{x\to 5} \frac{x^2-3x-10}{x-5}$$

indeterminate form of the type $\frac{0}{0}$

To evaluate, factor the top:

$$\lim_{x\to 5} \frac{x^2-3x-10}{x-5} = \lim_{x\to 5} \frac{(x-5)(x+2)}{x-5} = \lim_{x\to 5} x+2 = \boxed{7}$$

# Indeterminate Forms

$$\lim_{x\to 5} \frac{x^2-3x-10}{x-5}$$

indeterminate form of the type $\frac{0}{0}$

To evaluate, factor the top:

$$\lim_{x\to 5} \frac{x^2-3x-10}{x-5} = \lim_{x\to 5} \frac{(x-5)(x+2)}{x-5} = \lim_{x\to 5} x+2 = \boxed{7}$$

$$\lim_{x\to \infty} \frac{3x^2-4x+2}{8x^2-5}$$

indeterminate form of the type $\frac{\infty}{\infty}$

Q



ans

# Indeterminate Forms

$$\lim_{x\to 5} \frac{x^2-3x-10}{x-5}$$

indeterminate form of the type $\frac{0}{0}$

To evaluate, factor the top:

$$\lim_{x\to 5} \frac{x^2-3x-10}{x-5} = \lim_{x\to 5} \frac{(x-5)(x+2)}{x-5} = \lim_{x\to 5} x+2 = \boxed{7}$$

$$\lim_{x\to \infty} \frac{3x^2-4x+2}{8x^2-5}$$

indeterminate form of the type $\frac{\infty}{\infty}$

To evaluate, pull out $x^2$:

$$\lim_{x\to \infty} \frac{3x^2-4x+2}{8x^2-5} = \lim_{x\to \infty} \frac{x^2(3-\frac{4}{x}+\frac{2}{x^2})}{x^2(8-\frac{5}{x^2})} = \lim_{x\to \infty} \frac{3-\frac{4}{x}+\frac{2}{x^2}}{8-\frac{5}{x^2}} =$$

$$\lim_{x\to \infty} \frac{3-0+0}{8-0} = \boxed{\frac{3}{8}}$$

# Indeterminate Forms and the Derivative

$$\lim_{x \to 0} \frac{3 \sin x - x^4}{x^2 + \cos x - e^x}$$

indeterminate form of the type $\frac{0}{0}$

Q





ans

# Indeterminate Forms and the Derivative

$$\lim_{x\to 0} \frac{3\sin x - x^4}{x^2 + \cos x - e^x}$$

indeterminate form of the type $\frac{0}{0}$

Suppose $\lim_{x\to a} f(x) = \lim_{x\to a} g(x) = 0$. Suppose also that $f$ and $g$ are continuous and differentiable at $a$, and $g'(a) \neq 0$. Then:

$$\begin{aligned}
\lim_{x\to a} \frac{f(x)}{g(x)} &= \lim_{x\to a} \frac{f(x)-0}{g(x)-0} = \lim_{x\to a} \frac{f(x)-f(a)}{g(x)-g(a)} \\
&= \lim_{x\to a} \frac{f(x)-f(a)}{g(x)-g(a)} \frac{(x-a)^{-1}}{(x-a)^{-1}} \\
&= \lim_{x\to a} \left( \frac{\frac{f(x)-f(a)}{x-a}}{\frac{g(x)-g(a)}{x-a}} \right) = \frac{f'(a)}{g'(a)}
\end{aligned}$$

# Indeterminate Forms and the Derivative

$$\lim_{x\to 0} \frac{3\sin x - x^4}{x^2 + \cos x - e^x}$$

indeterminate form of the type $\frac{0}{0}$

$$\begin{aligned}
\lim_{x\to 0} \frac{3\sin x - x^4}{x^2 + \cos x - e^x} &= \frac{\frac{\mathrm{d}}{\mathrm{d}x}[3\sin x - x^4]|_{x=0}}{\frac{\mathrm{d}}{\mathrm{d}x}[x^2 + \cos x - e^x]|_{x=0}} \\
&= \frac{[3\cos x - 4x^3]|_{x=0}}{[2x - \sin x - e^x]|_{x=0}} \\
&= \frac{3-0}{0-0-1} = \boxed{-3}
\end{aligned}$$

A

## L'Hôpital's Rule: First Part – Theorem 3.7.2

Let $f$ and $g$ be functions such that $\lim_{x\to a} f(x) = 0 = \lim_{x\to a} g(x)$.

If $f'(a)$ and $g'(a)$ exist and $g'(a) \neq 0$, then $\lim_{x\to a} \dfrac{f(x)}{g(x)} = \dfrac{f'(a)}{g'(a)}$.

If $f$ and $g$ are differentiable on an open interval containing $a$, and if $\lim_{x\to a} \dfrac{f'(x)}{g'(x)}$ exists, then $\lim_{x\to a} \dfrac{f(x)}{g(x)} = \lim_{x\to a} \dfrac{f'(x)}{g'(x)}$.

This works even for $a = \pm\infty$.

Extremely Important Note:
L'Hôpital's Rule only works on indeterminate forms.

## L'Hôpital's Rule: Second Part – Theorem 3.7.2

Let $f$ and $g$ be functions such that $\lim_{x\to a} f(x) = \infty = \lim_{x\to a} g(x)$.

If $f'(a)$ and $g'(a)$ exist and $g'(a) \neq 0$, then $\lim_{x\to a} \dfrac{f(x)}{g(x)} = \dfrac{f'(a)}{g'(a)}$.

If $f$ and $g$ are differentiable on an open interval containing $a$, and if $\lim_{x\to a} \dfrac{f'(x)}{g'(x)}$ exists, then $\lim_{x\to a} \dfrac{f(x)}{g(x)} = \lim_{x\to a} \dfrac{f'(x)}{g'(x)}$.

This works even for $a = \pm\infty$.

Extremely Important Note:
L'Hôpital's Rule only works on indeterminate forms.

Evaluate:

$$\lim_{x \to 2} \frac{3x \tan(x-2)}{x-2}$$

Evaluate:

$$\lim_{x\to 2} \frac{3x\tan(x-2)}{x-2}$$

$$\lim_{x\to 2} \frac{3x\tan(x-2)}{x-2} \qquad \text{form } \frac{0}{0}$$


$$\underset{=}{\text{l'H}} \frac{3\left[x\sec^2(x-2)+\tan(x-2)\right]_{x=2}}{1}$$

$$=3\left[2\sec^2 0+\tan 0\right] = \boxed{6}$$

# Little Harder

$$\lim_{x \to 0} \frac{x^4}{e^x - \cos x - x}$$

indeterminate form of the type $\frac{0}{0}$



# Little Harder

$$\lim_{x\to 0} \frac{x^4}{e^x - \cos x - x}$$

indeterminate form of the type $\frac{0}{0}$

$$\lim_{x\to 0} \frac{x^4}{e^x - \cos x - x} \overset{?}{\underset{?}{=}} \left.\frac{4x^3}{e^x + \sin x - 1}\right|_{x=0} = \frac{0}{0}$$

oops

# Little Harder

$$\lim_{x\to 0} \frac{x^4}{e^x - \cos x - x}$$

indeterminate form of the type $\frac{0}{0}$

$$\lim_{x\to 0} \frac{x^4}{e^x - \cos x - x} \overset{?}{\underset{?}{=}} \left.\frac{4x^3}{e^x + \sin x - 1}\right|_{x=0} = \frac{0}{0}$$

oops

Iterate!

$$\lim_{x\to 0} \frac{x^4}{e^x - \cos x - x} \overset{\text{l'H}}{=} \lim_{x\to 0} \frac{4x^3}{e^x + \sin x - 1}$$
$$\overset{\text{l'H}}{=} \lim_{x\to 0} \frac{12x^2}{e^x + \cos x} = \frac{0}{2} = \boxed{0}$$

Evaluate:

$$\lim_{x \to \infty} \frac{\log x}{\sqrt{x}}$$

Evaluate:

$$\lim_{x\to\infty} \frac{\log x}{\sqrt{x}}$$

$$\lim_{x\to\infty} \frac{\log x}{\sqrt{x}} \overset{\text{l'H}}{=} \lim_{x\to\infty} \frac{\frac{1}{x}}{\frac{1}{2\sqrt{x}}}$$

$$= \lim_{x\to\infty} \frac{2\sqrt{x}}{x}$$

$$= \lim_{x\to\infty} \frac{2}{\sqrt{x}} = \boxed{0}$$

# Other Indeterminate Forms

$$\lim_{x \to \infty} e^{-x} \log x$$

form $0 \cdot \infty$

Q





ans

# Other Indeterminate Forms

$$\lim_{x\to\infty} e^{-x}\log x$$

form $0\cdot\infty$

$$\lim_{x\to\infty} e^{-x}\log x = \lim_{x\to\infty} \frac{\log x}{e^x} \qquad \text{form } \frac{\infty}{\infty}$$

$$\overset{\text{l'H}}{=} \lim_{x\to\infty} \frac{1/x}{e^x}$$

$$= \lim_{x\to\infty} \frac{1}{xe^x} = \boxed{0}$$

# VOTE VOTE VOTE

Which of the following can you immediately apply L'Hôpital's rule to?

A. $\dfrac{e^x}{2e^x+1}$

B. $\lim\limits_{x\to 0} \dfrac{e^x}{2e^x+1}$

C. $\lim\limits_{x\to \infty} \dfrac{e^x}{2e^x+1}$

D. $\lim\limits_{x\to \infty} e^{-x}(2e^x+1)$

E. $\lim\limits_{x\to 0} \dfrac{e^x}{x^2}$

# VOTE VOTE VOTE

Which of the following can you immediately apply L'Hôpital's rule to?

A. $\dfrac{e^x}{2e^x+1}$

B. $\lim\limits_{x\to 0} \dfrac{e^x}{2e^x+1}$

C. $\lim\limits_{x\to \infty} \dfrac{e^x}{2e^x+1}$

D. $\lim\limits_{x\to \infty} e^{-x}(2e^x+1)$

E. $\lim\limits_{x\to 0} \dfrac{e^x}{x^2}$

A

# Votey McVoteface

Suppose you want to use L'Hôpital's rule to evaluate $\lim_{x \to a} \frac{f(x)}{g(x)}$, which has the form $\frac{0}{0}$. How does the quotient rule fit into this problem?


A. You should use the quotient rule because the function you are differentiating is a quotient.

B. You will not use the quotient rule because you differentiate the numerator and the denominator separately

C. You may use the quotient rule because perhaps $f(x)$ or $g(x)$ is itself in the form of a quotient

D. You will not use L'Hôpital's rule because $\frac{0}{0}$ is not an appropriate indeterminate form

E. You will not use L'Hôpital's rule because, since the top has limit zero, the whole function has limit 0






ans

# Votey McVoteface

Suppose you want to use L'Hôpital's rule to evaluate $\lim_{x \to a} \frac{f(x)}{g(x)}$, which has the form $\frac{0}{0}$. How does the quotient rule fit into this problem?


A. You should use the quotient rule because the function you are differentiating is a quotient.

B. You will not use the quotient rule because you differentiate the numerator and the denominator separately

C. You may use the quotient rule because perhaps $f(x)$ or $g(x)$ is itself in the form of a quotient

D. You will not use L'Hôpital's rule because $\frac{0}{0}$ is not an appropriate indeterminate form

E. You will not use L'Hôpital's rule because, since the top has limit zero, the whole function has limit 0



# More Questions

Which of the following is NOT an indeterminate form?

A. $\frac{\infty}{\infty}$ for example, $\lim_{x\to\infty} \frac{e^x}{x^2}$

B. $\frac{0}{0}$ for example, $\lim_{x\to 0} \frac{e^x - 1}{x}$

C. $\frac{0}{\infty}$ for example, $\lim_{x\to 0^+} \frac{x}{\log x}$

D. $0 \cdot \infty$ for example, $\lim_{x\to\infty} x(\arctan(x) - \pi/2)$

E. all of the above are indeterminate forms

Q



ans

# More Questions

Which of the following is NOT an indeterminate form?

A. $\frac{\infty}{\infty}$ for example, $\lim_{x\to\infty} \frac{e^x}{x^2}$

B. $\frac{0}{0}$ for example, $\lim_{x\to 0} \frac{e^x - 1}{x}$

C. $\frac{0}{\infty}$ for example, $\lim_{x\to 0^+} \frac{x}{\log x} = 0$

D. $0 \cdot \infty$ for example, $\lim_{x\to\infty} x(\arctan(x) - \pi/2)$

E. all of the above are indeterminate forms

A

# I HAVE SO MANY QUESTIONS

Which of the following is NOT an indeterminate form?

A. $1^{\infty}$ for example, $\lim_{x\to\infty}\left(\frac{x+1}{x}\right)^{x}$

B. $0^{\infty}$ for example, $\lim_{x\to\infty}\left(\frac{1}{x}\right)^{x}$

C. $\infty^{0}$ for example, $\lim_{x\to\infty} x^{\frac{1}{x}}$

D. $0^{0}$ for example, $\lim_{x\to 0^{+}} x^{x}$

E. all of the above are indeterminate forms

F. none of the above are indeterminate forms

Q



ans

# I HAVE SO MANY QUESTIONS

Which of the following is NOT an indeterminate form?

A. $1^{\infty}$ for example, $\lim_{x\to\infty}\left(\frac{x+1}{x}\right)^{x}$

B. $0^{\infty}$ for example, $\lim_{x\to\infty}\left(\frac{1}{x}\right)^{x}=0$

C. $\infty^{0}$ for example, $\lim_{x\to\infty} x^{\frac{1}{x}}$

D. $0^{0}$ for example, $\lim_{x\to 0^{+}} x^{x}$

E. all of the above are indeterminate forms

F. none of the above are indeterminate forms

A



# Exponential Indeterminate Forms

$$\lim_{x \to \infty} x^{1/x}$$

# Exponential Indeterminate Forms

$$\lim_{x\to\infty} x^{1/x}$$

$$\begin{aligned}
\lim_{x\to\infty} x^{1/x} &= \lim_{x\to\infty} e^{\left(\log\left(x^{1/x}\right)\right)} \\
&= \lim_{x\to\infty} e^{\left(\frac{\log x}{x}\right)} \\
&= e^{\left(\lim_{x\to\infty} \frac{\log x}{x}\right)} \\
&\overset{\text{l'H}}{=} e^{\left(\lim_{x\to\infty} \frac{1}{x}\right)} \\
&= e^0 = \boxed{1}
\end{aligned}$$

# Exponential Indeterminate Forms

$$\lim_{x\to\infty}\left(1+\frac{2}{x}\right)^{3x}$$

# Exponential Indeterminate Forms

$$\lim_{x\to\infty}\left(1+\frac{2}{x}\right)^{3x}$$

First we calculate:

$$\begin{aligned}\lim_{x\to\infty}\log\left(\left(1+\frac{2}{x}\right)^{3x}\right) &= \lim_{x\to\infty} 3x\log\left(1+\frac{2}{x}\right)\\ &= \lim_{x\to\infty}\frac{3\log\left(1+\frac{2}{x}\right)}{x^{-1}}\\ &\overset{\text{l'H}}{=} \lim_{x\to\infty}\frac{3\left(\frac{-2x^{-2}}{1+2/x}\right)}{-x^{-2}}\\ &= \lim_{x\to\infty}\frac{6}{1+2/x} = 6\end{aligned}$$

So, now:

$$\lim_{x\to\infty}\left(1+\frac{2}{x}\right)^{3x} = \boxed{e^6}$$

Evaluate:

$$\lim_{x\to\infty} \frac{\log x}{\log \sqrt{x}}$$

$$\lim_{x\to\infty} (\log x)^{\sqrt{x}}$$

$$\lim_{x\to 0} \frac{\arcsin x}{x}$$



ans

Evaluate:

$$\lim_{x\to\infty} \frac{\log x}{\log \sqrt{x}}$$

Easier to simplify first.

$$\lim_{x\to\infty} (\log x)^{\sqrt{x}}$$

Not an indeterminate form: huge number to a huge power. Limit is infinity.

$$\lim_{x\to 0} \frac{\arcsin x}{x}$$

L'Hôpital: $\lim_{x\to 0} \dfrac{\frac{1}{\sqrt{1-x^2}}}{1} = 1$

# More Examples

$$\lim_{x\to\infty} \sqrt{2x^2+1} - \sqrt{x^2+x}$$

$$\lim_{x\to 0} \sqrt[x^2]{\sin^2 x}$$

$$\lim_{x\to 0} \sqrt[x^2]{\cos x}$$

$$\lim_{x \to \infty} \sqrt{2x^2 + 1} - \sqrt{x^2 + x}$$

$\lim_{x\to\infty} \sqrt{2x^2+1} - \sqrt{x^2+x}$ has the indeterminate form $\infty - \infty$. To get a better idea of what's going on, let's rationalize.

$$\lim_{x\to\infty} \sqrt{2x^2+1} - \sqrt{x^2+x}$$

$$= \lim_{x\to\infty} \left(\sqrt{2x^2+1} - \sqrt{x^2+x}\right)\left(\frac{\sqrt{2x^2+1} + \sqrt{x^2+x}}{\sqrt{2x^2+1} + \sqrt{x^2+x}}\right)$$

$$= \lim_{x\to\infty} \frac{(2x^2+1) - (x^2+x)}{\sqrt{2x^2+1} + \sqrt{x^2+x}} = \lim_{x\to\infty} \frac{x^2 - x + 1}{\sqrt{2x^2+1} + \sqrt{x^2+x}}$$

Here, we have the indeterminate form $\frac{\infty}{\infty}$, so L'Hôpital's Rule applies. However, if we try to use it here, we quickly get a huge mess. Instead, remember how we dealt with these kinds of limits in the past: factor out the highest power of the denominator, which is $x$.

$$\lim_{x\to\infty}\frac{x^2-x+1}{\sqrt{2x^2+1}+\sqrt{x^2+x}}=\lim_{x\to\infty}\frac{x\left(x-1+\frac{1}{x}\right)}{\sqrt{x^2(2+\frac{1}{x^2})}+\sqrt{x^2(1+\frac{1}{x})}}$$

$$=\lim_{x\to\infty}\frac{x\left(x-1+\frac{1}{x}\right)}{x\left(\sqrt{2+\frac{1}{x^2}}+\sqrt{1+\frac{1}{x}}\right)}$$

$$=\lim_{x\to\infty}\frac{x-1+\frac{1}{x}}{\underbrace{\sqrt{2+\frac{1}{x^2}}+\sqrt{1+\frac{1}{x}}}_{\substack{\text{num}\to\infty\\ \text{den}\to\sqrt{2}+1}}}$$

$$=\infty$$

$$\lim_{x \to 0} \sqrt[x^2]{\sin^2 x}$$

A

$$\lim_{x\to 0} \sqrt[x^2]{\sin^2 x}$$

$\lim_{x\to 0} \sin^2 x = 0$, and $\lim_{x\to 0} \frac{1}{x^2} = \infty$, so we have the form $0^\infty$. (Note that $\sin^2 x$ is positive, so our root is defined.) This is not an indeterminate form: $\lim_{x\to 0} \sqrt[x^2]{\sin^2 x} = 0$.

$$\lim_{x \to 0} \sqrt[x^2]{\cos x}$$

$$\lim_{x\to 0} \sqrt[x^2]{\cos x}$$

$\lim_{x\to 0} \cos x = 1$ and $\lim_{x\to 0} \frac{1}{x^2} = \infty$, so $\lim_{x\to 0} (\cos x)^{\frac{1}{x^2}}$ has the indeterminate form $1^\infty$. We want to use L'Hôpital, but we need to get our function into a fractional indeterminate form. So, we'll use a logarithm.

$$y := (\cos x)^{\frac{1}{x^2}}$$

$$\log y = \log\left((\cos x)^{\frac{1}{x^2}}\right) = \frac{1}{x^2}\log(\cos x) = \frac{\log\cos x}{x^2}$$

$$\lim_{x\to 0} \log y = \lim_{x\to 0} \underbrace{\frac{\log\cos x}{x^2}}_{\substack{\text{num}\to 0\\ \text{den}\to 0}} = \lim_{x\to 0} \frac{\frac{-\sin x}{\cos x}}{2x} = \lim_{x\to 0} \underbrace{\frac{-\tan x}{2x}}_{\substack{\text{num}\to 0\\ \text{den}\to 0}} = \lim_{x\to 0} \frac{-\sec^2 x}{2}$$

$$= \lim_{x\to 0} \frac{-1}{2\cos^2 x} = -\frac{1}{2}$$

Therefore, $\lim_{x\to 0} y = \lim_{x\to 0} e^{\log y} = e^{-1/2} = \frac{1}{\sqrt{e}}$

Sketch the graph of $f(x) = x \log x$.
Note: when you want to know $\lim_{x \to 0} f(x)$, you'll need to use L'Hôpital.

Evaluate $\lim_{x \to 0^+} (\csc x)^x$

$$f(x) = x \log x$$

$$f(x) = x \log x$$


- Domain: $x > 0$
- HA: none
- Intercepts: $(1, 0)$
- $y = \log x$ has a VA at $x = 0$. For our function, let's see what its behaviour is near 0:

$$\lim_{x \to 0^+} x \log x = \lim_{x \to 0^+} \frac{\log x}{1/x} \qquad \text{form } \frac{\infty}{\infty}$$

$$\overset{\text{l'H}}{=} \lim_{x \to 0^+} \frac{1/x}{-1/x^2} = \lim_{x \to 0^+} -x = 0$$

- $f'(x) = 1 + \log x$;
  CP at $x = \frac{1}{e}$ (and $y = -\frac{1}{e}$);
  As $x$ gets close to 0, $f'(x)$ goes to negative infinity, so near $x = 0$ our curve looks vertical.
  Decreasing on $(0, \frac{1}{e})$ and increasing on $(1/e, \infty)$
- $f''(x) = \frac{1}{x}$; concave up on entire domain

A


y
x

Evaluate $\lim_{x \to 0^+} (\csc x)^x$

A



Evaluate $\lim_{x\to 0^+} (\csc x)^x$

Indeterminate form: $\infty^0$

$$\begin{aligned} y &= (\csc x)^x \\ \log y &= x \log(\csc x) \\ \lim_{x\to 0^+} \log y &= \lim_{x\to 0^+} x \log(\csc x) \\ &= \lim_{x\to 0^+} x \log\left(\frac{1}{\sin x}\right) \\ &= \lim_{x\to 0^+} -x \log(\sin x) \end{aligned}$$

indeterminate form $0 \cdot \infty$

$$\begin{aligned} &= \lim_{x\to 0^+} \frac{\log(\sin x)}{-1/x} \\ &\overset{\text{l'H}}{=} \lim_{x\to 0^+} \frac{\frac{\cos x}{\sin x}}{-1/x^2} \\ &= \lim_{x\to 0^+} \frac{-x^2}{\tan x} \end{aligned}$$

indeterminate form $\frac{0}{0}$

$$\begin{aligned} &\overset{\text{l'H}}{=} \lim_{x\to 0^+} \frac{-2x}{\sec^2 x} \\ &= \frac{0}{1} = 0 \end{aligned}$$

conclusion

$$\lim_{x\to 0^+} \log y = 0$$

$$\lim_{x\to 0^+} y = e^0 = \boxed{1}$$

# Included Work

 ‘Notebook’ by Iconic is licensed under CC BY 3.0 (accessed 9 June 2021), 41, 49