


# TABLE OF CONTENTS



## WARMUP

Evaluate  $\int_3^7 \frac{1}{\sqrt{x^2 + 2x + 1}} dx$ .

Evaluate  $\int \frac{1}{\sqrt{x^2 + 1}} dx$ .

# CHECK OUR WORK

Let's verify that  $\int \frac{1}{\sqrt{x^2 + 1}} = \log \left| \sqrt{x^2 + 1} + x \right| + C.$

Seems improbable, right?

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
  - ▶  $x = \sin \theta$ ,  $1 - \sin^2 \theta = \cos^2 \theta$  changes  $\sqrt{1 - x^2}$  into

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
  - ▶  $x = \sin \theta$ ,  $1 - \sin^2 \theta = \cos^2 \theta$  changes  $\sqrt{1 - x^2}$  into  $\sqrt{\cos^2 \theta} = |\cos \theta|$



# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
  - ▶  $x = \sin \theta$ ,  $1 - \sin^2 \theta = \cos^2 \theta$  changes  $\sqrt{1 - x^2}$  into  $\sqrt{\cos^2 \theta} = |\cos \theta|$
  - ▶  $x = \tan \theta$ ,  $1 + \tan^2 \theta = \sec^2 \theta$  changes  $\sqrt{1 + x^2}$  into

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
  - ▶  $x = \sin \theta$ ,  $1 - \sin^2 \theta = \cos^2 \theta$  changes  $\sqrt{1 - x^2}$  into  $\sqrt{\cos^2 \theta} = |\cos \theta|$
  - ▶  $x = \tan \theta$ ,  $1 + \tan^2 \theta = \sec^2 \theta$  changes  $\sqrt{1 + x^2}$  into  $\sqrt{\sec^2 \theta} = |\sec \theta|$

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
  - ▶  $x = \sin \theta$ ,  $1 - \sin^2 \theta = \cos^2 \theta$  changes  $\sqrt{1 - x^2}$  into  $\sqrt{\cos^2 \theta} = |\cos \theta|$
  - ▶  $x = \tan \theta$ ,  $1 + \tan^2 \theta = \sec^2 \theta$  changes  $\sqrt{1 + x^2}$  into  $\sqrt{\sec^2 \theta} = |\sec \theta|$
  - ▶  $x = \sec \theta$ ,  $\sec^2 \theta - 1 = \tan^2 \theta$  changes  $\sqrt{x^2 - 1}$  into

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
  - ▶  $x = \sin \theta$ ,  $1 - \sin^2 \theta = \cos^2 \theta$  changes  $\sqrt{1 - x^2}$  into  $\sqrt{\cos^2 \theta} = |\cos \theta|$
  - ▶  $x = \tan \theta$ ,  $1 + \tan^2 \theta = \sec^2 \theta$  changes  $\sqrt{1 + x^2}$  into  $\sqrt{\sec^2 \theta} = |\sec \theta|$
  - ▶  $x = \sec \theta$ ,  $\sec^2 \theta - 1 = \tan^2 \theta$  changes  $\sqrt{x^2 - 1}$  into  $\sqrt{\tan^2 \theta} = |\tan \theta|$

# METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form:  $\sqrt{\text{quadratic}}$ , and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
  - ▶  $x = \sin \theta$ ,  $1 - \sin^2 \theta = \cos^2 \theta$  changes  $\sqrt{1 - x^2}$  into  $\sqrt{\cos^2 \theta} = |\cos \theta|$
  - ▶  $x = \tan \theta$ ,  $1 + \tan^2 \theta = \sec^2 \theta$  changes  $\sqrt{1 + x^2}$  into  $\sqrt{\sec^2 \theta} = |\sec \theta|$
  - ▶  $x = \sec \theta$ ,  $\sec^2 \theta - 1 = \tan^2 \theta$  changes  $\sqrt{x^2 - 1}$  into  $\sqrt{\tan^2 \theta} = |\tan \theta|$
- ▶ After integrating, convert back to the original variable (possibly using a triangle—more details later)

# FOCUS ON THE ALGEBRA

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

Choose a trigonometric substitution that will allow the square root to cancel out of the following expressions:

▶  $\sqrt{x^2 - 1}$

▶  $\sqrt{x^2 + 1}$

▶  $\sqrt{1 - x^2}$



## FOCUS ON THE ALGEBRA

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

Choose a trigonometric substitution that will allow the square root to cancel out of the following expressions:

►  $\sqrt{x^2 + 7}$

►  $\sqrt{3 - 2x^2}$



## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Consider the substitution  $x = \sin \theta$ ,  $dx = \cos \theta \, d\theta$  for the integral:

$$\int_0^1 \sqrt{1-x^2} \, dx$$

When  $x = 0$  (lower limit of integration), what is  $\theta$ ?

When  $x = 1$  (upper limit of integration), what is  $\theta$ ?





# CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

More generally, suppose  $a$  is a positive constant and we use the substitution  $x = a \sin \theta$  for the term  $\sqrt{a^2 - x^2}$ .

# CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

More generally, suppose  $a$  is a positive constant and we use the substitution  $x = a \sin \theta$  for the term  $\sqrt{a^2 - x^2}$ .

►  $\theta = \arcsin\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

# CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

More generally, suppose  $a$  is a positive constant and we use the substitution  $x = a \sin \theta$  for the term  $\sqrt{a^2 - x^2}$ .


- ▶  $\theta = \arcsin\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶  $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|$

## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

More generally, suppose  $a$  is a positive constant and we use the substitution  $x = a \sin \theta$  for the term  $\sqrt{a^2 - x^2}$ .

- ▶  $\theta = \arcsin\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶  $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|$
- ▶ On the interval  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $\cos \theta \geq 0$ , so  $|\cos \theta| = \cos \theta$




## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

More generally, suppose  $a$  is a positive constant and we use the substitution  $x = a \sin \theta$  for the term  $\sqrt{a^2 - x^2}$ .

- ▶  $\theta = \arcsin\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶  $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|$
- ▶ On the interval  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $\cos \theta \geq 0$ , so  $|\cos \theta| = \cos \theta$



- ▶ So, in general, when we use the substitution  $x = \sin \theta$  with trigonometric substitution, we can expect  $|\cos \theta| = \cos \theta$ .

# CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Now, consider the substitution  $x = a \tan \theta$  for  $\sqrt{a^2 + x^2}$ , where  $a$  is a positive constant.

## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Now, consider the substitution  $x = a \tan \theta$  for  $\sqrt{a^2 + x^2}$ , where  $a$  is a positive constant.

▶  $\theta = \arctan\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

# CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Now, consider the substitution  $x = a \tan \theta$  for  $\sqrt{a^2 + x^2}$ , where  $a$  is a positive constant.

▶  $\theta = \arctan\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

▶  $\sqrt{a^2 + x^2} = \sqrt{a^2 + a^2 \tan^2 \theta} = \sqrt{a^2 \sec^2 \theta} = \frac{a}{|\cos \theta|}$




## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Now, consider the substitution  $x = a \tan \theta$  for  $\sqrt{a^2 + x^2}$ , where  $a$  is a positive constant.

- ▶  $\theta = \arctan\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶  $\sqrt{a^2 + x^2} = \sqrt{a^2 + a^2 \tan^2 \theta} = \sqrt{a^2 \sec^2 \theta} = \frac{a}{|\cos \theta|}$
- ▶ On the interval  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $\cos \theta \geq 0$ , so  $|\cos \theta| = \cos \theta$  and  $|\sec \theta| = \sec \theta$ .




## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Now, consider the substitution  $x = a \tan \theta$  for  $\sqrt{a^2 + x^2}$ , where  $a$  is a positive constant.

- ▶  $\theta = \arctan\left(\frac{x}{a}\right)$ , so  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶  $\sqrt{a^2 + x^2} = \sqrt{a^2 + a^2 \tan^2 \theta} = \sqrt{a^2 \sec^2 \theta} = \frac{a}{|\cos \theta|}$
- ▶ On the interval  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $\cos \theta \geq 0$ , so  $|\cos \theta| = \cos \theta$  and  $|\sec \theta| = \sec \theta$ .



- ▶ So, in general, when we use the substitution  $x = \tan \theta$  with trigonometric substitution, we can expect  $|\sec \theta| = \sec \theta$ .

# CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Finally, consider the substitution  $x = a \sec \theta$  for  $\sqrt{x^2 - a^2}$ , where  $a$  is a positive constant.

## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution  $x = a \sec \theta$  for  $\sqrt{x^2 - a^2}$ , where  $a$  is a positive constant.

▶  $\sec \theta = \frac{x}{a}$ , so  $\cos \theta = \frac{a}{x}$ , so  $\theta = \arccos\left(\frac{a}{x}\right)$ . Then  $0 \leq \theta \leq \pi$

## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution  $x = a \sec \theta$  for  $\sqrt{x^2 - a^2}$ , where  $a$  is a positive constant.

- ▶  $\sec \theta = \frac{x}{a}$ , so  $\cos \theta = \frac{a}{x}$ , so  $\theta = \arccos\left(\frac{a}{x}\right)$ . Then  $0 \leq \theta \leq \pi$
- ▶  $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$

## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution  $x = a \sec \theta$  for  $\sqrt{x^2 - a^2}$ , where  $a$  is a positive constant.

- ▶  $\sec \theta = \frac{x}{a}$ , so  $\cos \theta = \frac{a}{x}$ , so  $\theta = \arccos\left(\frac{a}{x}\right)$ . Then  $0 \leq \theta \leq \pi$
- ▶  $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:

## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution  $x = a \sec \theta$  for  $\sqrt{x^2 - a^2}$ , where  $a$  is a positive constant.

- ▶  $\sec \theta = \frac{x}{a}$ , so  $\cos \theta = \frac{a}{x}$ , so  $\theta = \arccos\left(\frac{a}{x}\right)$ . Then  $0 \leq \theta \leq \pi$
- ▶  $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:
  - ▶ For  $\sqrt{x^2 - a^2}$  to be defined, we need  $x^2 \geq a^2$ . I.e.  $x \geq a$  or  $x \leq -a$ .

## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution  $x = a \sec \theta$  for  $\sqrt{x^2 - a^2}$ , where  $a$  is a positive constant.

- ▶  $\sec \theta = \frac{x}{a}$ , so  $\cos \theta = \frac{a}{x}$ , so  $\theta = \arccos\left(\frac{a}{x}\right)$ . Then  $0 \leq \theta \leq \pi$
- ▶  $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:
  - ▶ For  $\sqrt{x^2 - a^2}$  to be defined, we need  $x^2 \geq a^2$ . I.e.  $x \geq a$  or  $x \leq -a$ .
  - ▶ When  $x \geq a$ , we have  $0 \leq \theta < \frac{\pi}{2}$ ,  $\tan \theta \geq 0$ , so  $|\tan \theta| = \tan \theta$ .





## CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution  $x = a \sec \theta$  for  $\sqrt{x^2 - a^2}$ , where  $a$  is a positive constant.

- ▶  $\sec \theta = \frac{x}{a}$ , so  $\cos \theta = \frac{a}{x}$ , so  $\theta = \arccos\left(\frac{a}{x}\right)$ . Then  $0 \leq \theta \leq \pi$
- ▶  $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:
  - ▶ For  $\sqrt{x^2 - a^2}$  to be defined, we need  $x^2 \geq a^2$ . I.e.  $x \geq a$  or  $x \leq -a$ .
  - ▶ When  $x \geq a$ , we have  $0 \leq \theta < \frac{\pi}{2}$ ,  $\tan \theta \geq 0$ , so  $|\tan \theta| = \tan \theta$ .
  - ▶ When  $x \leq -a$ , we have  $\frac{\pi}{2} < \theta \leq \pi$ ,  $\tan \theta < 0$ , so  $|\tan \theta| = -\tan \theta$ .



# ABSOLUTE VALUES

From now on, we will assume:

- ▶ With the substitution  $x = a \sin \theta$  for  $\sqrt{a^2 - x^2}$ ,  $|\cos \theta| = \cos \theta$
- ▶ With the substitution  $x = a \tan \theta$  for  $\sqrt{a^2 + x^2}$ ,  $|\sec \theta| = \sec \theta$

## Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate  $\int_0^1 (1 + x^2)^{-3/2} dx$

Evaluate  $\int_0^1 (1+x^2)^{-3/2} dx$

## Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate  $\int \sqrt{1 - 4x^2} \, dx$

Evaluate  $\int \sqrt{1 - 4x^2} \, dx$

# CHECK OUR WORK

In the last example, we computed

$$\int \sqrt{1 - 4x^2} \, dx = \frac{1}{4} (\arcsin(2x) + 2x\sqrt{1 - 4x^2}) + C.$$

To check, we differentiate.

## Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate  $\int \frac{1}{\sqrt{x^2 - 1}} dx$



Evaluate  $\int \frac{1}{\sqrt{x^2 - 1}} dx$

# CHECK OUR WORK

Let's check our result,  $\int \frac{1}{\sqrt{x^2 - 1}} dx = \log \left| x + \sqrt{x^2 - 1} \right| + C.$

# COMPLETING THE SQUARE

Choose a trigonometric substitution to simplify  $\sqrt{3 - x^2 + 2x}$ .

Identities have two “parts” that turn into one part:

▶  $1 - \sin^2 \theta = \cos^2 \theta$

▶  $1 + \tan^2 \theta = \sec^2 \theta$

▶  $\sec^2 \theta - 1 = \tan^2 \theta$

But our quadratic expression has *three* parts.

# COMPLETING THE SQUARE

Choose a trigonometric substitution to simplify  $\sqrt{3 - x^2 + 2x}$ .

Identities have two “parts” that turn into one part:

▶  $1 - \sin^2 \theta = \cos^2 \theta$

▶  $1 + \tan^2 \theta = \sec^2 \theta$

▶  $\sec^2 \theta - 1 = \tan^2 \theta$

But our quadratic expression has *three* parts.

Fact:  $3 - x^2 + 2x = 4 - (x - 1)^2$

# COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write  $3 - x^2 + 2x$  in the form  $c - (x + b)^2$  for constants  $b, c$ .

# COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write  $3 - x^2 + 2x$  in the form  $c - (x + b)^2$  for constants  $b, c$ .

1. Find  $b$ :

# COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write  $3 - x^2 + 2x$  in the form  $c - (x + b)^2$  for constants  $b, c$ .

1. Find  $b$ :  $-2bx = 2x$ , so  $b = -1$

# COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write  $3 - x^2 + 2x$  in the form  $c - (x + b)^2$  for constants  $b, c$ .

1. Find  $b$ :  $-2bx = 2x$ , so  $b = -1$

2. Solve for  $c$ :



# COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write  $3 - x^2 + 2x$  in the form  $c - (x + b)^2$  for constants  $b, c$ .

1. Find  $b$ :  $-2bx = 2x$ , so  $b = -1$

2. Solve for  $c$ :  $3 = c - b^2 = c - 1$ , so  $c = 4$

# COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write  $3 - x^2 + 2x$  in the form  $c - (x + b)^2$  for constants  $b, c$ .

1. Find  $b$ :  $-2bx = 2x$ , so  $b = -1$

2. Solve for  $c$ :  $3 = c - b^2 = c - 1$ , so  $c = 4$

3. All together:

# COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write  $3 - x^2 + 2x$  in the form  $c - (x + b)^2$  for constants  $b, c$ .

1. Find  $b$ :  $-2bx = 2x$ , so  $b = -1$

2. Solve for  $c$ :  $3 = c - b^2 = c - 1$ , so  $c = 4$

3. All together:  $3 - x^2 + 2x = 4 - (x - 1)^2$

Evaluate  $\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} dx$ .

Identities have two “parts” that turn into one part:

- ▶  $1 - \sin^2 \theta = \cos^2 \theta$
- ▶  $1 + \tan^2 \theta = \sec^2 \theta$
- ▶  $\sec^2 \theta - 1 = \tan^2 \theta$

One of those parts is a constant, and one is squared.

Evaluate  $\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} dx = \int \frac{(x - 3)^2}{\sqrt{9 - (x - 3)^2}} dx.$

# CHECK OUR WORK

Let's verify that

$$\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} = \frac{9}{2} \left( \arcsin \left( \frac{x-3}{3} \right) - \frac{x-3}{3} \cdot \frac{\sqrt{6x-x^2}}{3} \right) + C :$$

## Included Work



'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021, modified), 14–16



'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021), 35, 37, 40, 52