
# TABLE OF CONTENTS



Fact:

$$\frac{\mathrm{d}}{\mathrm{d}x}\left\{\sin\left(x^2+x\right)\right\} =$$

Fact:

$$\frac{\mathrm{d}}{\mathrm{d}x}\left\{\sin\left(x^2+x\right)\right\} = (2x+1)\cos(x^2+x)$$

Fact:

$$\frac{\mathrm{d}}{\mathrm{d}x}\left\{\sin\left(x^2+x\right)\right\} = (2x+1)\cos(x^2+x)$$

**Related Fact:** 

$$\int (2x+1)\cos(x^2+x)\,\mathrm{d}x =$$

Fact:

$$\frac{\mathrm{d}}{\mathrm{d}x}\left\{\sin\left(x^2+x\right)\right\} = (2x+1)\cos(x^2+x)$$

**Related Fact:** 

$$\int (2x+1)\cos(x^2+x) \, dx = \sin(x^2+x) + C$$

Fact:

$$\frac{\mathrm{d}}{\mathrm{d}x}\left\{\sin\left(x^2+x\right)\right\} = (2x+1)\cos(x^2+x)$$

**Related Fact:** 

$$\int (2x+1)\cos(x^2+x) \, dx = \sin(x^2+x) + C$$

Hard to guess the antiderivative without seeing the derivative first!

#### Chain Rule:

$$\frac{\mathrm{d}}{\mathrm{d}x} \left\{ \sin \left( \underbrace{x^2 + x} \right) \right\} =$$
inside function

#### Chain Rule:

$$\frac{\mathrm{d}}{\mathrm{d}x} \left\{ \sin\left(\frac{x^2 + x}{x^2 + x}\right) \right\} = \left(\frac{2x + 1}{x}\right) \cos\left(\frac{x^2 + x}{x^2 + x}\right)$$
inside function
inside function

#### Chain Rule:

$$\frac{d}{dx} \left\{ \sin \left( \frac{x^2 + x}{x^2 + x} \right) \right\} = \left( \frac{2x + 1}{x^2 + x} \right) \cos \left( \frac{x^2 + x}{x^2 + x} \right)$$
inside function
inside function

Hallmark of the chain rule: an "inside" function, with that function's derivative multiplied.

# **SOLVE BY INSPECTION**

$$\int 2xe^{x^2+1}\,\mathrm{d}x$$

$$\int \frac{1}{x} \cos(\log x) \, \mathrm{d}x$$

$$\int 3(\sin x + 1)^2 \cos x \, \mathrm{d}x$$

(Look for an "inside" function, with its derivative multiplied.)

### Chain Rule:

$$\frac{\mathrm{d}}{\mathrm{d}x}\left\{f(u(x))\right\} = f'(u(x)) \cdot u'(x)$$

(Here, u(x) is our "inside function")

#### Antiderivative Fact:

$$\int f'(u(x)) \cdot u'(x) \, dx = f(u(x)) + C$$

### **Antiderivative Fact:**

$$\int f'(u(x)) \cdot u'(x) \, dx = f(u(x)) + C$$

### **Antiderivative Fact:**

$$\int f'(u(x)) \cdot u'(x) \, \mathrm{d}x = f(u(x)) + C$$

Shorthand: call u(x) simply u; since  $\frac{du}{dx} = u'(x)$ , call u'(x) dx simply du.

#### **Antiderivative Fact:**

$$\int f'(u(x)) \cdot u'(x) \, \mathrm{d}x = f(u(x)) + C$$

Shorthand: call u(x) simply u; since  $\frac{du}{dx} = u'(x)$ , call u'(x) dx simply du.

$$\int f'(u(x)) \cdot u'(x) \, \mathrm{d}x = \int f'(u) \, \mathrm{d}u \Big|_{u=u(x)} = f(u(x)) + C$$

This is the substitution rule.

We saw these integrals before, and solved them by inspection. Now try using the language of substitution.

$$\int 2xe^{x^2+1}\,\mathrm{d}x$$

$$\int \frac{1}{x} \cos(\log x) \, \mathrm{d}x$$

$$\int 3(\sin x + 1)^2 \cos x \, \mathrm{d}x$$

$$\int (3x^2)\sin(x^3+1)\,\mathrm{d}x =$$

$$\int (3x^2)\sin(x^3+1)\,\mathrm{d}x =$$

"Inside" function:  $x^3 + 1$ .



$$\int (3x^2)\sin(x^3+1)\,\mathrm{d}x =$$

"Inside" function:  $x^3 + 1$ . Its derivative:  $3x^2$ 

$$\int (3x^2)\sin(x^3+1)\,\mathrm{d}x =$$



$$\int (3x^2) \sin(x^3 + 1) dx = \int \sin(u) du \Big|_{u = x^3 + 1}$$

$$\int (3x^2) \sin(x^3 + 1) dx = \int \sin(u) du \Big|_{u=x^3+1}$$
$$= -\cos(u) + C|_{u=x^3+1}$$

$$\int (3x^2) \sin(x^3 + 1) dx = \int \sin(u) du \Big|_{u=x^3+1}$$

$$= -\cos(u) + C|_{u=x^3+1}$$

$$= \cos(x^3 + 1) + C$$

$$\int (3x^2) \sin(x^3 + 1) dx = \int \sin(u) du \Big|_{u=x^3+1}$$

$$= -\cos(u) + C|_{u=x^3+1}$$

$$= \cos(x^3 + 1) + C$$

Warning 1: We don't just change dx to du. We need to couple dx with the derivative of our inside function.

After all, we're undoing the chain rule! We need to have an "inside derivative."

$$\int (3x^2) \sin(x^3 + 1) dx = \int \sin(u) du \Big|_{u=x^3+1}$$

$$= -\cos(u) + C|_{u=x^3+1}$$

$$= \cos(x^3 + 1) + C$$

Warning 2: The final answer is a function of *x*.

We used the substitution rule to conclude

$$\int (3x^2)\sin(x^3+1)\,\mathrm{d}x = -\cos(x^3+1) + C$$

We can check that our antiderivative is correct by differentiating.

We saw:

$$\int 3x^2 \sin(x^3 + 1) \, \mathrm{d}x = -\cos(x^3 + 1) + C$$

So, we can evaluate:

$$\int_0^1 3x^2 \sin(x^3 + 1) \, \mathrm{d}x$$

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$



$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$

$$\uparrow \qquad \qquad \downarrow$$

$$x = \frac{\pi}{2}$$

$$x = \frac{\pi}{4}$$


$$\int_{1/\sqrt{2}}^{1} \frac{1}{u^3} du$$

$$\downarrow u = \sin\left(\frac{\pi}{2}\right) = 1$$

$$u = \sin\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}$$


$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$


$$\int_{\pi/4}^{\pi/2} \frac{1}{u^3} \, \mathrm{d}u$$



$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$









$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$








$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$

$$\int_{x=\pi/4}^{x=\pi/2} \frac{1}{u^3} \, \mathrm{d}u$$



$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$

$$\uparrow \qquad \qquad \qquad \downarrow$$

$$x = \frac{\pi}{2}$$

$$x = \frac{\pi}{4}$$

$$\int_{x=\pi/4}^{x=\pi/2} \frac{1}{u^3} \, \mathrm{d}u$$

$$x = \frac{\pi}{2}$$

$$x = \frac{\pi}{4}$$

$$u = \sin\left(\frac{\pi}{2}\right) = 1$$

$$u = \sin\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}$$

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$

Let  $u = \sin x$ ,  $du = \cos x dx$ . Note the limits (or bounds) of integration  $\pi/4$  and  $\pi/2$  are values of x, not u: they follow the differential, unless otherwise specified.

not standard, but OK

#### NOTATION: LIMITS OF INTEGRATION

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$

Let  $u = \sin x$ ,  $du = \cos x \, dx$ . Note the limits (or bounds) of integration  $\pi/4$  and  $\pi/2$  are values of x, not u: they follow the differential, unless otherwise specified.

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, dx = \int_{x=\pi/4}^{x=\pi/2} \frac{1}{u^3} \, du = \int_{1/\sqrt{2}}^{1} \frac{1}{u^3} \, du$$

$$\uparrow \qquad \qquad \downarrow$$

$$x = \frac{\pi}{2}$$

$$x = \frac{\pi}{4}$$

$$x = \frac{\pi}{2}$$

$$x = \frac{\pi}{4}$$

$$u = \sin(\frac{\pi}{2}) = 1$$

$$u = \sin(\frac{\pi}{4}) = \frac{1}{\sqrt{2}}$$

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x$$

Let  $u = \sin x$ ,  $du = \cos x dx$ .

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} \, \mathrm{d}x = \int_{1/\sqrt{2}}^1 \frac{1}{u^3} \, \mathrm{d}u$$

## TRUE OR FALSE?

1. Using  $u = x^2$ ,

$$\int e^{x^2} \, \mathrm{d}x = \int e^u \, \mathrm{d}u$$

2. Using  $u = x^2 + 1$ ,

$$\int_0^1 x \sin(x^2 + 1) \, \mathrm{d}x = \int_0^1 \frac{1}{2} \sin u \, \, \mathrm{d}u$$

Evaluate 
$$\int_0^1 x^7 (x^4 + 1)^5 dx$$
.

Time permitting, more examples using the substitution rule

Evaluate  $\int \sin x \cos x \, dx$ .

We can check that 
$$\int \sin x \cos x \, dx = \frac{1}{2} \sin^2 x + C$$
 by differentiating.

We can check that 
$$\int \sin x \cos x \, dx = -\frac{1}{2} \cos^2 x + C$$
 by differentiating.

Evaluate 
$$\int \frac{\log x}{3x} dx$$
.



We can check that 
$$\int \frac{\log x}{3x} dx = \frac{1}{6} \log^2 x + C$$
 by differentiating.

Evaluate 
$$\int \frac{e^x}{e^x + 15} dx$$
.

Evaluate 
$$\int x^4(x^5+1)^8 dx$$
.



We can check that 
$$\int \frac{e^x}{e^x + 15} dx = \log |e^x + 15| + C$$
 by differentiating.

We can check that 
$$\int x^4(x^5+1)^8 dx = \frac{1}{45}(x^5+1)^9 + C$$
 by differentiating.

Evaluate  $\int_4^8 \frac{s}{s-3} ds$ . Be careful to use correct notation.

Evaluate 
$$\int x^9 (x^5 + 1)^8 dx$$
.

We can check that 
$$\int x^9 (x^5 + 1)^8 dx = \frac{1}{5} \left[ \frac{(x^5 + 1)^{10}}{10} - \frac{(x^5 + 1)^9}{9} \right] + C$$
 by differentiating.

# PARTICULARLY TRICKY SUBSTITUTION

Evaluate 
$$\int \frac{1}{e^x + e^{-x}} dx$$
.

We can check that 
$$\int \frac{1}{e^x + e^{-x}} dx = \arctan(e^x) + C$$
 by differentiating.

#### Included Work

'Notebook' by Iconic is licensed under CC BY 3.0 (accessed 9 June 2021, modified), 15, 46