

#### TABLE OF CONTENTS




A sequence is a list of numbers A series is the sum of such a list.



Square of Area 1

Size of Tiles:



$$\frac{1}{2}$$

Size of Tiles:

$$\frac{1}{2}$$
,  $\frac{1}{2^2}$ 

$$\frac{1}{2} + \frac{1}{2^2}$$

Size of Tiles:

$$\frac{1}{2}$$
,  $\frac{1}{2^2}$ ,  $\frac{1}{2^3}$ 

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3}$$

Size of Tiles:

$$\frac{1}{2}$$
,  $\frac{1}{2^2}$ ,  $\frac{1}{2^3}$ ,  $\frac{1}{2^4}$ 


$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4}$$

Size of Tiles:

$$\frac{1}{2}$$
,  $\frac{1}{2^2}$ ,  $\frac{1}{2^3}$ ,  $\frac{1}{2^4}$ ,  $\frac{1}{2^5}$ 

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5}$$

Size of Tiles:




$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6}$$

Size of Tiles:

$$\frac{1}{2}$$
,  $\frac{1}{2^2}$ ,  $\frac{1}{2^3}$ ,  $\frac{1}{2^4}$ ,  $\frac{1}{2^5}$ ,  $\frac{1}{2^6}$ ,  $\frac{1}{2^7}$ 

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7}$$

Size of Tiles:




Covered Area:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8}$$

 $\Box$ 

Size of Tiles:



Covered Area:


$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9}$$

12/140

Size of Tiles:

Sequence

 $\frac{1}{2} \ , \frac{1}{2^2} \ , \frac{1}{2^3} \ , \frac{1}{2^4} \ , \frac{1}{2^5} \ , \frac{1}{2^6} \ , \frac{1}{2^7} \ , \frac{1}{2^8} \ , \frac{1}{2^9} \ , \ \cdots$ 



Covered Area:


$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$$

13/140

Size of Tiles:

Sequence

**List** of numbers, approaching



$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$$

Size of Tiles:

List of numbers,

approaching **zero**.

$$\frac{1}{2}$$
,  $\frac{1}{2^2}$ ,  $\frac{1}{2^3}$ ,  $\frac{1}{2^4}$ ,  $\frac{1}{2^5}$ ,  $\frac{1}{2^6}$ ,  $\frac{1}{2^7}$ ,  $\frac{1}{2^8}$ ,  $\frac{1}{2^9}$ , ...

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$$


Size of Tiles:

Sequence

List of numbers,

approaching zero.

Series



Covered Area:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$$

16/140

Size of Tiles:

Sequence


**List** of numbers, approaching **zero**.

Series

**Sum** of numbers,

approaching

$$\frac{1}{2} \ , \frac{1}{2^2} \ , \frac{1}{2^3} \ , \frac{1}{2^4} \ , \frac{1}{2^5} \ , \frac{1}{2^6} \ , \frac{1}{2^7} \ , \frac{1}{2^8} \ , \frac{1}{2^9} \ , \ \dots$$



$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$$

Size of Tiles:

Sequence


**List** of numbers, approaching **zero**.

Series

Sum of numbers,

approaching one.

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$$



18/140

#### QUICK REVIEW: SIGMA NOTATION

Recall:

$$\sum_{n=1}^{5} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2}$$

#### **QUICK REVIEW: SIGMA NOTATION**

Recall:

$$\sum_{n=1}^{5} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2}$$

We informally interpret:

$$\sum_{n=1}^{\infty} \frac{1}{n^2} =$$

#### QUICK REVIEW: SIGMA NOTATION

Recall:

$$\sum_{n=1}^{5} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2}$$

We informally interpret:

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \frac{1}{7^2} + \frac{1}{8^2} + \frac{1}{9^2} + \frac{1}{10^2} + \cdots$$

(a more rigorous definition will be discussed soon)

Let  $a_n$  and  $b_n$  be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (C \cdot a_n) =$$

A. 
$$\sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$$

$$B. \sum_{n=1}^{\infty} C + \sum_{n=1}^{\infty} a_n$$

C. 
$$C\sum_{n=1}^{\infty} a_n$$

D. 
$$a_n \sum_{i=1}^{\infty} C$$

E. none of the above

Let  $a_n$  and  $b_n$  be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n + b_n) =$$

A. 
$$\sum_{n=1}^{\infty} a_n \cdot \sum_{n=1}^{\infty} b_n$$

$$B. \sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n$$

$$C. \ a_n + \sum_{n=1}^{\infty} b_n$$

D. 
$$a_n \sum_{n=1}^{\infty} b_n$$

E. none of the above



Let  $a_n$  and  $b_n$  be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n)^C =$$

A. 
$$\sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$$

B. 
$$\left(\sum_{n=1}^{\infty} a_n\right)^C$$

C. 
$$C^n \sum_{n=1}^{\infty} a_n$$

D. 
$$\sum_{n=1}^{\infty} C(a_n)^{C-1}$$

E. none of the above

$$1-1+1-1+1-1+1-1+1-1+1-1+\cdots$$

$$\underbrace{1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + \dots}_{0} + \underbrace{1 - 1 + 1 - 1 + 1 - 1 + \dots}_{0} + \underbrace{1 - 1 + 1 - 1 + 1 - 1 + \dots}_{0}$$

$$\underbrace{1 - 1 + 1}_{1} \underbrace{-1 + 1}_{0} \underbrace{-1 + 1}_{0}$$

What does it really mean to add up infinitely many things?

$$1-1+1-1+1-1+1-1+1-1+1-1+\cdots$$

We need an unambiguous definition.


SEQUENCE OF PARTIAL SUMS





SEQUENCE OF PARTIAL SUMS

 $S_1 = 0.2000$ 




SEQUENCE OF PARTIAL SUMS

 $S_1 = 0.2000$ 

 $S_2 = 0.2400$ 






SEQUENCE OF PARTIAL SUMS



$$S_2 = 0.2400$$
  
 $S_3 = 0.2480$ 





SEQUENCE OF PARTIAL SUMS


 $S_1 = 0.2000$ 

 $S_2 = 0.2400$ 

 $S_3=0.2480$ 

\_ \_\_\_\_

 $S_4 = 0.2496$ 



SEQUENCE OF PARTIAL SUMS

 $S_1 = 0.2000$ 

 $S_2 = 0.2400$ 

 $S_3=0.2480$ 

 $S_4 = 0.2496$ 

 $S_5 = 0.2499$ 



**Partial sums** let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

$$a_1 = \frac{1}{5} = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

$$a_{1} = \frac{1}{5} = 0.2$$

$$a_{2} = \frac{1}{5^{2}} = 0.04$$

$$a_{3} = \frac{1}{5^{3}} = 0.008$$

$$a_{4} = \frac{1}{5^{4}} = 0.0016$$

$$a_{5} = \frac{1}{5^{5}} = 0.00032$$



$$a_1 = \frac{1}{5} = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

$$a_{1} = \frac{1}{5} = 0.2$$

$$a_{2} = \frac{1}{5^{2}} = 0.04$$

$$a_{3} = \frac{1}{5^{3}} = 0.008$$

$$a_{4} = \frac{1}{5^{4}} = 0.0016$$

$$a_{5} = \frac{1}{5^{5}} = 0.00032$$

$$S_{1} = 0.2$$

$$S_{2} = 0.24$$

$$a_{1} = \frac{1}{5} = 0.2$$

$$a_{2} = \frac{1}{5^{2}} = 0.04$$

$$a_{3} = \frac{1}{5^{3}} = 0.008$$

$$a_{4} = \frac{1}{5^{4}} = 0.0016$$

$$a_{5} = \frac{1}{5^{5}} = 0.00032$$

$$S_{1} = 0.2$$

$$S_{2} = 0.24$$

$$a_{1} = \frac{1}{5} = 0.2$$

$$a_{2} = \frac{1}{5^{2}} = 0.04$$

$$a_{3} = \frac{1}{5^{3}} = 0.008$$

$$a_{4} = \frac{1}{5^{4}} = 0.0016$$

$$a_{5} = \frac{1}{5^{5}} = 0.00032$$

$$S_{1} = 0.2$$

$$S_{2} = 0.24$$

$$S_{3} = 0.248$$

$$a_1 = \frac{1}{5} = 0.2$$
  $S_1 = 0.2$   $S_2 = 0.24$   $S_3 = 0.24$   $S_4 = 0.008$   $S_5 = 0.248$   $S_6 = 0.0016$   $S_7 = 0.00032$ 

$$a_1 = \frac{1}{5} = 0.2$$
  $S_1 = 0.2$   $a_2 = \frac{1}{5^2} = 0.04$   $S_2 = 0.24$   $a_3 = \frac{1}{5^3} = 0.008$   $S_3 = 0.248$   $a_4 = \frac{1}{5^4} = 0.0016$   $S_4 = 0.2496$   $a_5 = \frac{1}{5^5} = 0.00032$ 

$$a_1 = \frac{1}{5} = 0.2$$
  $S_1 = 0.2$   $a_2 = \frac{1}{5^2} = 0.04$   $S_2 = 0.24$   $a_3 = \frac{1}{5^3} = 0.008$   $S_3 = 0.248$   $a_4 = \frac{1}{5^4} = 0.0016$   $S_4 = 0.2496$   $a_5 = \frac{1}{5^5} = 0.00032$ 

$$a_1 = \frac{1}{5} = 0.2$$
  $S_1 = 0.2$   $S_2 = 0.24$   $S_3 = 0.248$   $S_4 = 0.248$   $S_4 = 0.2496$   $S_5 = 0.24992$ 


We define 
$$\sum_{n=1}^{\infty} a_n = \lim_{N \to \infty} \sum_{n=1}^{N} a_n = \lim_{N \to \infty} S_N$$
.

$$a_1 = \frac{1}{5} = 0.2$$
  $S_1 = 0.2$   $a_5 = \frac{1}{5^5} = 0.00032$   $S_5 = 0.24992$   $a_2 = \frac{1}{5^2} = 0.04$   $S_2 = 0.24$   $a_6 = \frac{1}{5^6} = 0.000064$   $S_6 = 0.249984$   $a_3 = \frac{1}{5^3} = 0.008$   $S_3 = 0.248$   $a_7 = \frac{1}{5^7} = 0.0000128$   $S_7 = 0.2499968$   $a_4 = \frac{1}{5^4} = 0.0016$   $S_4 = 0.2496$   $a_8 = \frac{1}{5^8} = 0.00000256$   $S_8 = 0.24999936$ 


From the sequence of partial sums, we guess

$$\sum_{n=1}^{\infty} = \lim_{N \to \infty} S_N =$$











$$S_1 = a_1$$
$$S_2 = a_1 + a_2$$






$$S_2 = a_1 + a_2$$
 $S_3 = a_1 + a_2 + a_3$ 
 $a_4 \quad a_5 \quad a_6$ 





$$S_2 = a_1 + a_2$$
 $S_3 = a_1 + a_2 + a_3$ 
 $S_4 = a_1 + \dots + a_4$ 



 $a_1 + \cdots + a_5$ 







Suppose 
$$\sum_{n=1}^{\infty} a_n$$
 has partial sums  $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$ .

ightharpoonup Evaluate  $\sum_{n=1}^{100} a_n$ .


ightharpoonup Evaluate  $\sum_{n=1}^{\infty} a_n$ .





Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.






Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.






Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.






Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.





Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.


Suppose 
$$\sum_{n=1}^{\infty} a_n$$
 has partial sums  $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$ .

▶ Find  $a_1$ .

► Give an explicit expression for  $a_n$ , when n > 1.

$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$






$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$

 $S_1 = 1/(1+1)$ 






$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$

$$S_1=1/(1+1)$$

$$S_2 = 2/(2+1)$$






64/140

$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$

$$S_1 = 1/(1+1)$$

$$S_2 = 2/(2+1)$$

$$S_3 = 3/(3+1)$$



65/140



$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$


$$S_1 = 1/(1+1)$$

$$S_2 = 2/(2+1)$$
  
 $S_3 = 3/(3+1)$ 

 $a_5$ 

$$a_6$$
  $a_7$   $a_8$ 

$$S_4 = 4/(4+1)$$



66/140

#### Definition

The  $N^{\text{th}}$  partial sum of the series  $\sum_{n=1}^{\infty} a_n$  is the sum of its first N terms

$$S_N = \sum_{n=1}^N a_n.$$

The partial sums form a sequence  $\{S_N\}_{N=1}^{\infty}$ . If this sequence of partial sums converges  $S_N \to S$  as  $N \to \infty$  then we say that the series  $\sum_{n=1}^{\infty} a_n$  converges to S and we write

$$\sum_{n=1}^{\infty} a_n = S$$

If the sequence of partial sums diverges, we say that the series diverges.

### Geometric Series

Let *a* and *r* be two fixed real numbers with  $a \neq 0$ . The series

$$a + ar + ar^2 + ar^3 + \cdots$$

is called the **geometric series** with first term a and ratio r.

We call *r* the *ratio* because it is the quotient of consecutive terms:

$$\frac{ar^{n+1}}{ar^n} = r$$

#### Geometric Series

Let *a* and *r* be two fixed real numbers with  $a \neq 0$ . The series

$$a + ar + ar^2 + ar^3 + \cdots$$

is called the **geometric series** with first term *a* and ratio *r*.

We call *r* the *ratio* because it is the quotient of consecutive terms:

$$\frac{ar^{n+1}}{ar^n} = r$$

Another useful way of identifying geometric series is to determine whether all pairs of consecutive terms have the same ratio.

- Geometric:  $1 + \frac{1}{5} + \frac{1}{5^2} + \frac{1}{5^3} + \frac{1}{5^4} + \cdots$
- Geometric:  $\sum_{n=0}^{\infty} \frac{1}{2^n}$
- ► Not geometric:  $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \cdots$

### Consider the partial sum $S_N$ of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \dots + ar^N$$

#### Consider the partial sum $S_N$ of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \dots + ar^N$$
  
$$rS_N =$$

#### Consider the partial sum $S_N$ of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \dots + ar^N$$
  
$$rS_N = ar + ar^2 + ar^3 + \dots + ar^N + ar^{N+1}$$

## Consider the partial sum $S_N$ of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \dots + ar^N$$

$$rS_N = ar + ar^2 + ar^3 + \dots + ar^N + ar^{N+1}$$

$$rS_N - S_N =$$

## Consider the partial sum $S_N$ of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \dots + ar^N$$

$$rS_N = ar + ar^2 + ar^3 + \dots + ar^N + ar^{N+1}$$

$$rS_N - S_N = -a + ar^{N+1}$$

Consider the partial sum  $S_N$  of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \dots + ar^N$$
  
 $rS_N = ar + ar^2 + ar^3 + \dots + ar^N + ar^{N+1}$   
 $rS_N - S_N = -a + ar^{N+1}$   
 $S_N(r-1) = ar^{N+1} - a$ 

If  $r \neq 1$ , then

$$S_N = \frac{ar^{N+1} - a}{r - 1} = a\frac{r^{N+1} - 1}{r - 1}$$

## Geometric Series and Partial Sums

Let *a* and *r* be constants with  $a \neq 0$ , and let *N* be a natural number.

► If 
$$r \neq 1$$
, then  $a + ar + ar^2 + ar^3 + \dots + ar^N = a \frac{r^{N+1} - 1}{r - 1}$ .

► If 
$$r = 1$$
, then  $a + ar + ar^2 + ar^3 + \cdots + ar^N =$ 

$$If |r| < 1, then \sum_{n=0}^{\infty} ar^n =$$

$$If r = 1, then \sum_{n=0}^{\infty} ar^n$$

$$If r = -1, then \sum_{n=0}^{\infty} ar^n$$


• If 
$$|r| > 1$$
, then  $\sum_{n=0}^{\infty} ar^n$ 

$$\sum_{n=0}^{\infty} ar^n, r=1, a\neq 0$$











$$S_0 = a$$

$$S_1 = 2a$$





$$S_0 = a$$

$$S_1 = 2a$$








$$S_0 = a$$

$$S_1 = 2a$$

$$S_2 = 3a$$

$$S_3 = 4a$$








$$S_0 = a$$


$$S_1 = 2a$$


$$S_2 = 3a$$

$$S_3 = 4a$$

$$S_4 = 5a$$







$$S_0 = a$$

$$S_1 = 2a$$


$$S_2 = 3a$$

$$S_3 = 4a$$


$$S_4 = 5a$$

$$S_5=6a$$






$$\sum_{n=0}^{\infty} ar^n, r=-1, a\neq 0$$







$$\sum_{n=0}^{\infty} ar^n, r=-1, a\neq 0$$



$$S_0 = a$$

$$S_1 = 0$$




n=0

а


$$S_0 = a$$

$$S_1 = 0$$

$$S_2 = a$$



$$\sum_{n=0}^{\infty} ar^n$$
,  $r = -1$ ,  $a \neq 0$ 



$$S_0 = a$$


$$S_1 = 0$$

$$S_2 = a$$

$$S_3 = 0$$



$$\sum_{n=0}^{\infty} ar^n, r=-1, a\neq 0$$



$$S_0 = a$$


$$S_1 = 0$$

$$S_2 = a$$

$$S_3 = 0$$

$$S_4 = a$$

n=0



$$S_0 = a$$

$$S_1 = 0$$


$$S_2 = a$$

$$S_3 = 0$$

$$S_4 = a$$

$$S_5 = 0$$

$$\sum_{n=0}^{\infty} ar^n, r=-1, a\neq 0$$



$$S_0 = a$$

$$S_1 = 0$$


$$S_2 = a$$

$$S_3 = 0$$

$$S_4 = a$$

$$S_5 = 0$$

 $\sum_{n=0}^{3.2 \, \text{Series}} ar^n, \, r>1, \, a\neq 0$ 








 $S_0 = a$ 








$$ar^2 \qquad ar^3 \qquad ar^4$$

$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$










$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$








$$S_0 = a$$


$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$








$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$




$$S_0 = a$$


$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

















$$S_0 = a$$
$$S_1 = a \frac{r^2 - 1}{r - 1}$$











$$S_0 = a$$


$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$








$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$



$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$


$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$

$$S_5 = a \frac{r^6 - 1}{r - 1}$$



$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$



$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$

$$S_5 = a \frac{r^6 - 1}{r - 1}$$

## GEOMETRIC SERIES

New bitcoins are produced when a particular type of computational problem is solved. Every time 210,000 solutions are found, the number of bitcoins each solution can produce is cut in half.

- ► Each of the first 210,000 solutions can produce 50 bitcoins.
- ► Each of the next 210,000 solutions can produce  $\frac{50}{2}$  bitcoins.
- ► Each of the next 210,000 solutions can produce  $\frac{50}{2^2}$  bitcoins.
- ► Each of the next 210,000 solutions can produce  $\frac{50}{2^3}$  bitcoins.

Assume that this continues forever, and that bitcoins are infinitely divisible. How many bitcoins can possibly be produced?


 $<sup>^{1}</sup>$ Actually the smallest allowed division of a bitcoin is  $10^{-8}$ .

# GEOMETRIC SERIES

New bitcoins are produced when a particular type of computational problem is solved. Every time 210,000 solutions are found, the number of bitcoins each solution can produce is cut in half. How many bitcoins can possibly be produced?


$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n}\right) = 21\,000\,000$$





$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n}\right) = 21\,000\,000$$

 $S_0 = 10\,500\,000$ 



$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n}\right) = 21\,000\,000$$

 $S_0 = 10\,500\,000$ 

 $S_1 = 15\,750\,000$ 








15 750 000


$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n}\right) = 21\,000\,000$$





$$S_1 = 15\,750\,000$$





$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n}\right) = 21\,000\,000$$

 $S_0 = 10\,500\,000$ 


 $S_1 = 15\,750\,000$ 

 $S_2 = 18\,375\,000$ 


 $S_3 = 19687500$ 







$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n}\right) = 21\,000\,000$$



$$S_0 = 10\,500\,000$$

$$S_1 = 15\,750\,000$$

$$S_2 = 18\,375\,000$$

$$S_3 = 19687500$$

$$S_4 = 20343750$$

#### Arithmetic of Series

Let *S*, *T*, and *C* be real numbers. Let the two series  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  converge to *S* and *T* respectively. Then

$$\sum_{n=1}^{\infty} [a_n + b_n] = S + T$$

$$\sum_{n=1}^{\infty} [a_n - b_n] = S - T$$

$$\sum_{n=1}^{\infty} [Ca_n] = CS$$

## Geometric Series and Partial Sums

Let a and r be fixed numbers, and let N be a positive integer. Then

$$\sum_{n=0}^{N} ar^{n} = \begin{cases} a \cdot \frac{1 - r^{N+1}}{1 - r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate 
$$\sum_{n=0}^{\infty} \left( \frac{2}{3^n} + \frac{4}{5^n} \right)$$

$$\sum_{n=0}^{\infty} \left( \frac{2}{3^n} + \frac{4}{5^n} \right) =$$

## Geometric Series and Partial Sums

Let *a* and *r* be fixed numbers, and let *N* be a positive integer. Then

$$\sum_{n=0}^{N} ar^{n} = \begin{cases} a \cdot \frac{1 - r^{N+1}}{1 - r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate 
$$\sum_{n=6}^{\infty} \left( \frac{3^{n-1}}{5^{2n}} \right)$$



$$\sum_{n=6}^{\infty} \left( \frac{3^{n-1}}{5^{2n}} \right) =$$

#### Geometric Series and Partial Sums

Let *a* and *r* be fixed numbers, and let *N* be a positive integer. Then

$$\sum_{n=0}^{N} ar^n = \begin{cases} a \cdot \frac{1-r^{N+1}}{1-r} & \text{if } r \neq 1\\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate 
$$\sum_{n=0}^{\infty} \left( \frac{2^{2n}}{3^n} \right)$$

$$\sum_{n=0}^{\infty} \left( \frac{2^{2n}}{3^n} \right) =$$

$$\sum_{n=0}^{\infty} \left( \frac{2^{2n}}{3^n} \right) \text{ diverges}$$






$$\sum_{n=0}^{\infty} \left( \frac{2^{2n}}{3^n} \right) \text{ diverges}$$

 $S_0 = 1.0000$ 






$$\sum_{n=0}^{\infty} \left( \frac{2^{2n}}{3^n} \right) \text{ diverges}$$

 $S_0 = 1.0000$ 

 $S_1 = 2.3333$ 





$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n}\right) \text{ diverges}$$



 $\frac{64}{27}$ 

 $\frac{256}{81}$ 



 $S_0=1.0000$ 

 $S_1 = 2.3333$ 

 $S_2=4.1111$ 



$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n}\right) \text{ diverges}$$





256 81


 $\frac{1024}{243}$ 

$$S_0 = 1.0000$$

$$S_1 = 2.3333$$

$$S_2=4.1111$$

$$S_3=6.4815$$



$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n}\right) \text{ diverges}$$



 $\frac{1024}{243}$ 




$$S_1 = 2.3333$$

$$S_2 = 4.1111$$

$$S_3=6.4815$$

$$S_4=9.6420$$



$$\sum_{n=0}^{\infty} \left( \frac{2^{2n}}{3^n} \right) \text{ diverges}$$

 $S_0=1.0000$ 


 $S_1 = 2.3333$ 

 $S_2 = 4.1111$ 

 $S_3 = 6.4815$ 

 $S_4 = 9.6420$ 

 $S_5=13.856$ 



# TELESCOPING SUMS

Evaluate 
$$\sum_{n=1}^{800} \left( \frac{1}{n} - \frac{1}{n+1} \right).$$

Evaluate 
$$\sum_{n=1}^{\infty} \left( \frac{1}{n} - \frac{1}{n+1} \right)$$
.

Evaluate 
$$\sum_{n=1}^{1000} \log \left( \frac{n+1}{n} \right)$$
.

Evaluate 
$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
.

Evaluate 
$$\sum_{n=1}^{1000} \log \left( \frac{n+1}{n} \right)$$
.

Evaluate 
$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
.

$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right) \text{ diverges}$$
















$$\log \frac{3}{2}$$

 $\log 2 \qquad \log \frac{3}{2} \quad \log \frac{4}{3} \, \log \frac{5}{4} \, \log \frac{6}{5} \, \log \frac{7}{6} \, \log \frac{8}{7}$ 



$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
 diverges


 $S_1=0.6931$ 



$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
 diverges

 $S_1=0.6931$ 

 $S_2 = 1.0986$ 



$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
 diverges

 $S_1=0.6931$ 

 $S_2 = 1.0986$ 

 $S_3 = 1.3863$ 










 $\log \tfrac{5}{4} \, \log \tfrac{6}{5} \, \log \tfrac{7}{6} \, \log \tfrac{8}{7}$ 



$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
 diverges


 $S_1 = 0.6931$ 

 $S_2 = 1.0986$ 

 $S_3 = 1.3863$ 

 $\log \frac{6}{5} \log \frac{7}{6} \log \frac{8}{7}$ 

 $S_4 = 1.6094$ 



$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
 diverges


 $S_1 = 0.6931$ 

 $S_2 = 1.0986$ 

 $S_3 = 1.3863$ 

 $S_4 = 1.6094$ 

 $S_5 = 1.7917$ 



$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
 diverges

 $S_1 = 0.6931$ 


 $S_2 = 1.0986$ 

 $S_3 = 1.3863$ 


 $S_4 = 1.6094$ 

 $S_5 = 1.7917$ 

 $S_6 = 1.9459$ 



$$\sum_{n=1}^{\infty} \log \left( \frac{n+1}{n} \right)$$
 diverges



$$S_1 = 0.6931$$

$$S_2 = 1.0986$$

$$S_3 = 1.3863$$

$$S_4 = 1.6094$$

$$S_5 = 1.7917$$

$$S_6 = 1.9459$$

$$S_7 = 2.0794$$

#### Included Work

P'Balloon' by Simon Farkas is licensed under CC-BY (accessed November 2022, edited), 85-92, 100-107 'Waage/Libra' by B. Lachner is in the public domain (accessed April 2021, edited), 30–35, 48–54, 62–66, 77–107, 110–115, 123–129, 133–140 Weight' by Kris Brauer is licensed under CC-BY(accessed May 2021), 30–35, 48–54, 62-66, 77-107, 110-115, 123-129, 133-140 60 'bov' by Xinh Studio is licensed under CC BY 3.0 (accessed 6 June 2023), 56–60 cookies' by Azam Ishaq is licensed under CC BY 3.0 (accessed 6 June 2023), 56–60 'cookies' by Vectors Point is licensed under CC BY 3.0 (accessed 6 June 2023), 56 - 60office desk' by Abdul Baasith is licensed under CC BY 3.0 (accessed 6 June 2023), 56-60 Man' by Xinh Studio is licensed under CC BY 3.0 (accessed 6 June 2023), 56–60 Notebook' by Iconic is licensed under CC BY 3.0 (accessed 9 June 2021, modified), 130 Notebook' by Iconic is licensed under CC BY 3.0 (accessed 9 June 2021), 108, 117, 119, 121