
## TABLE OF CONTENTS



## REVERSE THE PRODUCT RULE

### Product Rule:

$$\frac{\mathrm{d}}{\mathrm{d}x}\big\{u(x)\cdot v(x)\big\} = u'(x)\cdot v(x) + u(x)\cdot v'(x)$$

## REVERSE THE PRODUCT RULE

#### Product Rule:

$$\frac{\mathrm{d}}{\mathrm{d}x}\big\{u(x)\cdot v(x)\big\} = u'(x)\cdot v(x) + u(x)\cdot v'(x)$$

#### Related fact:

$$\int \left[ u'(x) \cdot v(x) + u(x) \cdot v'(x) \right] dx = u(x) \cdot v(x) + C$$

## REVERSE THE PRODUCT RULE

Product Rule:

$$\frac{\mathrm{d}}{\mathrm{d}x}\big\{u(x)\cdot v(x)\big\} = u'(x)\cdot v(x) + u(x)\cdot v'(x)$$

Related fact:

$$\int \left[ u'(x) \cdot v(x) + u(x) \cdot v'(x) \right] dx = u(x) \cdot v(x) + C$$

Rearrange:

$$\implies \int \left[ u'(x)v(x) \right] dx + \int \left[ u(x)v'(x) \right] dx = u(x)v(x) + C$$

$$\implies \int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

## INTEGRATION BY PARTS

$$\int \left[ u(x)v'(x) \right] \mathrm{d}x = u(x)v(x) - \int \left[ v(x)u'(x) \right] \mathrm{d}x$$
 Example: 
$$\int xe^x \mathrm{d}x$$



In the previous slide, we evaluated

$$\int xe^x dx = xe^x - e^x + C$$

for some constant *C*. We can check that this is correct by differentiating.

## INTEGRATION BY PARTS (IBP): A CLOSER LOOK

$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\underbrace{\int xe^{x} dx}_{\text{How to integrate?}} = x(e^{x}) - \underbrace{1 \int e^{x} dx}_{\text{Easy to integrate!}} + C$$

We start and end with an integral. IBP is only useful if the new integral is somehow an improvement.

We differentiate the function we choose as u(x), and antidifferentiate the function we choose as v'(x)

# Choosing u(x) and v(x)

$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\int \left[ x \sin x \right] dx =$$



# CHOOSING u(x) AND v(x)

$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\int \left[ x \sin x \right] dx =$$

### Option A:

$$u(x) = x$$
 $v'(x) = \sin x$ 
 $u(x) = \sin x$ 
 $v'(x) = x$ 

$$u(x) = \sin x$$
$$v'(x) = x$$



To check our work, we can calculate  $\frac{d}{dx} \left\{ -x \cos x + \sin x + C \right\}$ . It should work out to be  $x \sin x$ .

# Choosing u(x) and v(x)

$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\int \left[ x^2 \log x \right] dx =$$



# CHOOSING u(x) AND v(x)

$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\int \left[ x^2 \log x \right] dx =$$

Option A:

$$u(x) = x^2$$

$$f(x) = x$$
  
 $f(x) = \log x$ 

Option B:

$$\begin{array}{c|c} u(x) = x^2 & u(x) = \log x \\ v'(x) = \log x & v'(x) = x^2 \end{array}$$



To check our work, we can calculate  $\frac{d}{dx} \left\{ \frac{1}{3} x^3 \log x - \frac{1}{9} x^3 + C \right\}$ . It should work out to be  $x^2 \log x$ .

$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\int \left[ \frac{1}{2}xe^{6x} \right] dx =$$



$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\int \left[ \frac{1}{2}xe^{6x} \right] dx =$$

$$\begin{array}{c|c}
u(x) = \frac{1}{2}x & u(x) = e^{6x} \\
v'(x) = e^{6x} & v'(x) = \frac{1}{2}x
\end{array}$$

### Option B:

$$u(x) = e^{6x}$$
$$v'(x) = \frac{1}{2}x$$



We check that 
$$\int \left[\frac{1}{2}xe^{6x}\right] dx = \frac{1}{12}xe^{6x} - \frac{1}{72}e^{6x} + C$$
 by differentiating.

## **MNEMONIC**

$$\int \left[ u(x)v'(x) \right] dx = u(x)v(x) - \int \left[ v(x)u'(x) \right] dx + C$$

$$\int u \, dv = uv - \int v \, du + C$$

#### We abbreviate:

- ightharpoonup u(x) 
  ightharpoonup u
- $ightharpoonup u'(x) dx \rightarrow du$
- ightharpoonup v(x) 
  ightharpoonup v
- $ightharpoonup v'(x) dx \rightarrow dv$

## CHOOSING u, dv in your head

Choose u and dv to evaluate the integral below:

$$\int (3t+5)\cos(t/4)\mathrm{d}t$$

## Choosing u, dv in your head

Choose u and dv to evaluate the integral below:

$$\int (3t+5)\cos(t/4)\mathrm{d}t$$

Thoughts: 
$$\int u \, dv = uv - \int v \, du$$
*u* gets differentiated, and d*v* gets antidifferentiated.

## Choosing u, dv in your head

Choose u and dv to evaluate the integral below:

$$\int (3t+5)\cos(t/4)\mathrm{d}t$$

Thoughts: 
$$\int u \, dv = uv - \int v \, du$$
*u* gets differentiated, and d*v* gets antidifferentiated.

▶ If we differentiate OR antidifferentiate cos(t/4) we get almost the same thing (they only differ by a constant).

## CHOOSING u, dv in your head

Choose u and dv to evaluate the integral below:

$$\int (3t+5)\cos(t/4)\mathrm{d}t$$

Thoughts:  $\int u \, dv = uv - \int v \, du$ u gets differentiated, and dv gets antidifferentiated.

- ▶ If we differentiate OR antidifferentiate cos(t/4) we get almost the same thing (they only differ by a constant).
- ▶ If we differentiate 3t + 5, we get something simpler: a constant. If we antidifferentiate 3t + 4, we get something more complicated—a quadratic.

## Choosing u, dv in your head

Choose u and dv to evaluate the integral below:

$$\int (3t+5)\cos(t/4)\mathrm{d}t$$

Thoughts:  $\int u \, dv = uv - \int v \, du$  *u* gets differentiated, and d*v* gets antidifferentiated.

- ▶ If we differentiate OR antidifferentiate cos(t/4) we get almost the same thing (they only differ by a constant).
- ▶ If we differentiate 3t + 5, we get something simpler: a constant. If we antidifferentiate 3t + 4, we get something more complicated–a quadratic.

Simpler is better: we want to differentiate 3t + 5, which means we choose it to be u. Then  $dv = \cos(t/4)dt$ .

$$\int u dv = uv - \int v du + C$$

- $ightharpoonup \int x^2 e^x dx$

$$\int u dv = uv - \int v du + C$$

$$\int u dv = uv - \int v du + C$$



$$\int u dv = uv - \int v du + C$$

## **DEFINITE INTEGRATION BY PARTS**

Method 1: Antidifferentiate first, then plug in limits of integration.

Method 2: Plug as you go.

Evaluate 
$$\int_{1}^{e} \log^{2} x \, dx$$

## Special Technique: v'(x) = 1

$$\int u \, \mathrm{d}v = uv - \int v \, \mathrm{d}u + C$$

Evaluate  $\int \log x \, dx$  using integration by parts.

Let's check that 
$$\int \log x \, dx = x \log x - x + C$$
.

$$\int u \, \mathrm{d}v = uv - \int v \, \mathrm{d}u + C$$

Evaluate  $\int \arctan x \, dx$  using integration by parts.

Let's check that 
$$\int \arctan x \, dx = x \arctan x - \frac{1}{2} \log |1 + x^2| + C.$$

Setting dv = 1 dx is a very specific technique. You'll probably only see it in situations integrating logarithms and inverse trigonometric functions.

$$\int \log x \, dx$$
,  $\int \arcsin x \, dx$ ,  $\int \arccos x \, dx$ ,  $\int \arctan x \, dx$ , etc

Evaluate  $\int e^x \cos x \, dx$  using integration by parts.

Evaluate  $\int e^x \cos x \, dx$  using integration by parts.

Let  $u = e^x$  and  $dv = \cos x \, dx$ . Then  $du = e^x \, dx$  and  $v = \sin x$ :

$$\int e^x \cos x \, dx = e^x \sin x - \int e^x \sin x \, dx$$

Let  $u = e^x$  and  $dv = \sin x \, dx$ . Then  $du = e^x \, dx$  and  $v = -\cos x$ :

$$= e^x \sin x - \left[ -e^x \cos x - \int -e^x \cos x \, dx \right]$$

$$= e^x \sin x + e^x \cos x - \int e^x \cos x \, dx$$

$$\int e^x \cos x \, dx = e^x \sin x + e^x \cos x - \int e^x \cos x \, dx$$

$$2 \int e^x \cos x \, dx = e^x \sin x + e^x \cos x + C$$

$$\int e^x \cos x \, dx = \frac{1}{2} \left( e^x \sin x + e^x \cos x \right) + C$$

### INTEGRATING AROUND IN A CIRCLE

We can use this technique to antidifferentiate products of two functions that almost, but don't quite, stay the same under (anti)differentiation.

Use integration by parts a number of times, ending up with an expression involving (a scalar multiple of) the original integral.

To do this, be consistent with your choice of u and dv.

Evaluate  $\int e^x \cos x \, dx$  using integration by parts.

Let  $u = e^x$  and  $dv = \cos x \, dx$ . Then  $du = e^x \, dx$  and  $v = \sin x$ :

$$\int e^x \cos x \, dx = e^x \sin x - \int e^x \sin x \, dx$$

Let  $u = e^x$  and  $dv = \sin x \, dx$ . Then  $du = e^x \, dx$  and  $v = -\cos x$ :

$$= e^x \sin x - \left[ -e^x \cos x - \int -e^x \cos x \, dx \right]$$

$$= e^x \sin x + e^x \cos x - \int e^x \cos x \, dx$$

$$\int e^x \cos x \, dx = e^x \sin x + e^x \cos x - \int e^x \cos x \, dx$$

$$2 \int e^x \cos x \, dx = e^x \sin x + e^x \cos x + C$$

$$\int e^x \cos x \, dx = \frac{1}{2} \left( e^x \sin x + e^x \cos x \right) + C$$

Evaluate  $\int \cos(\log x) \, dx$ .

We check that  $\int \cos(\log x) dx = \frac{x}{2} [\cos(\log x) + \sin(\log x)] + C$  by differentiating.

Evaluate  $\int e^{2x} \sin x \, dx$  using integration by parts.

We can check our work by differentiating  $\frac{1}{5}e^{2x}[2\sin x - \cos x] + C$ . We should end up with  $e^{2x}\sin x$ .

#### Included Work

Notebook' by Iconic is licensed under CC BY 3.0 (accessed 9 June 2021), 23