


# TABLE OF CONTENTS



# REVIEW: AREA UNDER A CURVE


Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum
  - ▶ **Conceptually** easy – cut into rectangles




# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ **Conceptually** easy – cut into rectangles

- ▶ **Computationally** rough



$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$

# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ Conceptually easy – cut into rectangles

- ▶ Computationally rough

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$



- ▶ Use Geometry

# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ Conceptually easy – cut into rectangles


- ▶ Computationally rough

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$



- ▶ Use Geometry

- ▶ Computationally nice when it's available!  
(Circles, triangles, symmetry, etc.)



# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ Conceptually easy – cut into rectangles

- ▶ Computationally rough


$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$



- ▶ Use Geometry

- ▶ Computationally nice when it's available!  
(Circles, triangles, symmetry, etc.)

- ▶ Often not available – most functions don't make such nice shapes.



# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ Conceptually easy – cut into rectangles

- ▶ Computationally rough


$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$



- ▶ Use Geometry

- ▶ Computationally nice when it's available!  
(Circles, triangles, symmetry, etc.)

- ▶ Often not available – most functions don't make such nice shapes.



- ▶ Up next: Fundamental Theorem of Calculus



# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ Conceptually easy – cut into rectangles

- ▶ Computationally rough


$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$



- ▶ Use Geometry

- ▶ Computationally nice when it's available!  
(Circles, triangles, symmetry, etc.)

- ▶ Often not available – most functions don't make such nice shapes.



- ▶ Up next: Fundamental Theorem of Calculus

- ▶ Conceptually less obvious – we'll spend about a day explaining why it works

# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ **Conceptually** easy – cut into rectangles

- ▶ **Computationally** rough


$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$



- ▶ Use Geometry

- ▶ **Computationally** nice when it's available!  
(Circles, triangles, symmetry, etc.)

- ▶ Often not available – most functions don't make such nice shapes.



- ▶ Up next: Fundamental Theorem of Calculus

- ▶ **Conceptually** less obvious – we'll spend about a day explaining why it works

- ▶ **Computationally** generally nicer than Riemann sums

# REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

- ▶ Limit of a Riemann Sum

- ▶ **Conceptually** easy – cut into rectangles

- ▶ **Computationally** rough


$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}$$



- ▶ Use Geometry

- ▶ **Computationally** nice when it's available!  
(Circles, triangles, symmetry, etc.)

- ▶ Often not available – most functions don't make such nice shapes.



- ▶ Up next: Fundamental Theorem of Calculus

- ▶ **Conceptually** less obvious – we'll spend about a day explaining why it works

- ▶ **Computationally** generally nicer than Riemann sums

- ▶ Doesn't work for every function

## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$


for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$


FTC(I) gives us the derivative of a very specific function (subject to some fine print).

It shows a close relationship between integrals and derivatives.


AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$



AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$




AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$






AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$



Notation: the function  $A$  depends on the variable  $x$ .

We need to know how the function  $f$  behaves on the whole interval  $(0, x)$  to find  $A(x)$ . That's why we use  $f(t)$ , not  $f(x)$ .


AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$



Notation: the function  $A$  depends on the variable  $x$ .

We need to know how the function  $f$  behaves on the whole interval  $(0, x)$  to find  $A(x)$ . That's why we use  $f(t)$ , not  $f(x)$ .


AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$



Notation: the function  $A$  depends on the variable  $x$ .

We need to know how the function  $f$  behaves on the whole interval  $(0, x)$  to find  $A(x)$ . That's why we use  $f(t)$ , not  $f(x)$ .


AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$



Notation: the function  $A$  depends on the variable  $x$ .

We need to know how the function  $f$  behaves on the whole interval  $(0, x)$  to find  $A(x)$ . That's why we use  $f(t)$ , not  $f(x)$ .

AREA FUNCTION:  $A(x) = \int_a^x f(t)dt$  FOR  $a \leq x \leq b$



Notation: the function  $A$  depends on the variable  $x$ .

We need to know how the function  $f$  behaves on the whole interval  $(0, x)$  to find  $A(x)$ . That's why we use  $f(t)$ , not  $f(x)$ .


# AREA FUNCTION NOTATION

It might look strange at first to see two different variables. Let's consider the alternatives:

$$A(x) = \int_0^x f(t) \, dt$$

$$B(x) = \int_0^x f(x) \, dt$$

$$C(x) = \int_0^x f(x) \, dx$$



# AREA FUNCTION NOTATION


It might look strange at first to see two different variables. Let's consider the alternatives:

$$A(x) = \int_0^x f(t) \, dt$$

$$B(x) = \int_0^x f(x) \, dt$$

$$C(x) = \int_0^x f(x) \, dx$$

$$A(1) = \int_0^1 f(t) \, dt$$



# AREA FUNCTION NOTATION


It might look strange at first to see two different variables. Let's consider the alternatives:

$$A(x) = \int_0^x f(t) \, dt$$

$$B(x) = \int_0^x f(x) \, dt$$

$$C(x) = \int_0^x f(x) \, dx$$

$$A(2) = \int_0^2 f(t) \, dt$$





# AREA FUNCTION NOTATION


It might look strange at first to see two different variables. Let's consider the alternatives:

$$A(x) = \int_0^x f(t) dt$$

$$B(x) = \int_0^x f(x) dt$$

$$C(x) = \int_0^x f(x) dx$$

$$A(3) = \int_0^3 f(t) dt$$



# AREA FUNCTION NOTATION

It might look strange at first to see two different variables. Let's consider the alternatives:



$$A(x) = \int_0^x f(t) \, dt$$

$$B(x) = \int_0^x f(x) \, dt$$

$$C(x) = \int_0^x f(x) \, dx$$

$$A(3) = \int_0^3 f(t) \, dt$$

$$B(1) = \int_0^1 f(1) \, dt$$



# AREA FUNCTION NOTATION

It might look strange at first to see two different variables. Let's consider the alternatives:



$$A(x) = \int_0^x f(t) \, dt$$

$$B(x) = \int_0^x f(x) \, dt$$

$$C(x) = \int_0^x f(x) \, dx$$

$$A(3) = \int_0^3 f(t) \, dt$$

$$B(2) = \int_0^2 f(2) \, dt$$



# AREA FUNCTION NOTATION

It might look strange at first to see two different variables. Let's consider the alternatives:



$$A(x) = \int_0^x f(t) \, dt$$

$$B(x) = \int_0^x f(x) \, dt$$

$$C(x) = \int_0^x f(x) \, dx$$

$$A(3) = \int_0^3 f(t) \, dt$$

$$B(3) = \int_0^3 f(3) \, dt$$



# AREA FUNCTION NOTATION

It might look strange at first to see two different variables. Let's consider the alternatives:

$$A(x) = \int_0^x f(t) dt$$



$$B(x) = \int_0^x f(x) dt$$

$$C(x) = \int_0^x f(x) dx$$

$$A(3) = \int_0^3 f(t) dt$$

$$B(3) = \int_0^3 f(3) dt$$

$$C(1) = \int_0^1 f(1) \underbrace{d1}_{??}$$



## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let


$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

Question: Why is it true?


# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$



$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h}$$




# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$




$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h}$$

# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$




$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h}$$

# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$




$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h}$$

# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$




$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h}$$

# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$




$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h}$$

# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$



$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h} = \lim_{h \rightarrow 0} \frac{hf(x)}{h}$$

# DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t)dt$



$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h} = \lim_{h \rightarrow 0} \frac{hf(x)}{h} = f(x)$$

When  $h$  is very small, the purple area looks like a rectangle with base  $h$  and height  $f(x)$ , so  $A(x+h) - A(x) \approx hf(x)$  and  $\frac{A(x+h) - A(x)}{h} \approx f(x)$ . As  $h$  tends to zero, the error in this approximation approaches 0.

## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

Suppose  $A(x) = \int_2^x \sin t \, dt$ . What is  $A'(x)$ ?



## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

Suppose  $A(x) = \int_2^x \sin t \, dt$ . What is  $A'(x)$ ?

$$A'(x) = \sin x$$

## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

Suppose  $A(x) = \int_2^x \sin t \, dt$ . What is  $A'(x)$ ?

$$A'(x) = \sin x$$

Suppose  $B(x) = \int_x^2 \sin t \, dt$ . What is  $B'(x)$ ?

## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

Suppose  $A(x) = \int_2^x \sin t \, dt$ . What is  $A'(x)$ ?

$$A'(x) = \sin x$$

Suppose  $B(x) = \int_x^2 \sin t \, dt$ . What is  $B'(x)$ ?

$$B'(x) = \frac{d}{dx} \left\{ - \int_2^x f(t) \, dt \right\} = - \sin x$$

## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) \, dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

Suppose  $C(x) = \int_2^{e^x} \sin t \, dt$ . What is  $C'(x)$ ?



## Fundamental Theorem of Calculus, Part 1

Let  $a < b$  and let  $f(x)$  be a function which is defined and continuous on  $[a, b]$ . Let

$$A(x) = \int_a^x f(t) dt$$

for any  $x$  in  $[a, b]$ . Then the function  $A(x)$  is differentiable and

$$A'(x) = f(x) .$$

Suppose  $C(x) = \int_2^{e^x} \sin t dt$ . What is  $C'(x)$ ?


$C'(x) = e^x \sin(e^x)$ : if we set  $a = 2$ , then

$$\begin{aligned} C(x) &= A(e^x) \\ \implies C'(x) &= A'(e^x) \cdot \frac{d}{dx}\{e^x\} = \sin(e^x) \cdot e^x \end{aligned}$$




It's possible to have two different functions with the same derivative.

$$A(x) = \int_0^x 2t \, dt$$




$$B(x) = \int_1^x 2t \, dt$$




It's possible to have two different functions with the same derivative.

$$A(x) = \int_0^x 2t \, dt$$




$$A'(x) = 2x$$

$$B(x) = \int_1^x 2t \, dt$$




It's possible to have two different functions with the same derivative.

$$A(x) = \int_0^x 2t \, dt$$



$$A'(x) = 2x$$

$$B(x) = \int_1^x 2t \, dt$$




$$B'(x) = 2x$$




It's possible to have two different functions with the same derivative.

$$A(x) = \int_0^x 2t \, dt = x^2$$



$$A'(x) = 2x$$


$$B(x) = \int_1^x 2t \, dt$$



$$B'(x) = 2x$$


It's possible to have two different functions with the same derivative.

$$A(x) = \int_0^x 2t \, dt = x^2$$



$$A'(x) = 2x$$


$$B(x) = \int_1^x 2t \, dt = x^2 - 1$$



$$B'(x) = 2x$$


It's possible to have two different functions with the same derivative.

$$A(x) = \int_0^x 2t \, dt = x^2$$



$$A'(x) = 2x$$


$$B(x) = \int_1^x 2t \, dt = x^2 - 1$$



$$B'(x) = 2x$$

When two functions have the same derivative, they differ only by a constant.

In this example:  $B(x) = A(x) - 1$



If two continuous functions have the same derivative, then one is a constant plus the other.

Two clues for finding  $A(x) = \int_a^x f(t) \, dt$ :

- ▶ If  $A(x) = \int_a^x f(t) \, dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_a^x e^t dt$ . What functions could  $A(x)$  be?

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_a^x e^t dt$ . What functions could  $A(x)$  be?

- ▶  $A'(x) = e^x$ .
- ▶ Guess a function with derivative  $e^x$ :  $F(x) = e^x$ .
- ▶ Then  $A(x) = e^x + C$  for some constant  $C$ .

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_a^x \cos t dt$ . What functions could  $A(x)$  be?

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )



Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_a^x \cos t dt$ . What functions could  $A(x)$  be?

- ▶  $A'(x) = \cos x$ .
- ▶ Guess a function with derivative  $\cos x$ :  $F(x) = \sin x$ .
- ▶ Then  $A(x) = \sin x + C$  for some constant  $C$ .

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_{-2}^x 5t^4 dt$ . What functions could  $A(x)$  be?

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_{-2}^x 5t^4 dt$ . What functions could  $A(x)$  be?

- ▶  $A'(x) = 5x^4$ .
- ▶ Guess a function with derivative  $5x^4$ :  $F(x) = x^5$ .
- ▶ Then  $A(x) = x^5 + C$  for some constant  $C$ .
- ▶ We ALSO know  $A(-2) = \int_{-2}^{-2} 5t^4 dt = 0$ , so we can find  $C$ :


$$0 = A(-2) = (-2)^5 + C \implies C = 32$$

- ▶ So,  $A(x) = x^5 + 32$


---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$




$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$




$$A(-1) = \int_{-2}^{-1} 5t^4 dt = (-1)^5 + 32 = 31$$

$$A(x) = \int_{-2}^x 5t^4 \, dt = x^5 + 32$$




$$A(0) = \int_{-2}^0 5t^4 \, dt = (0)^5 + 32 = 32$$

$$A(x) = \int_{-2}^x 5t^4 \, dt = x^5 + 32$$



$$A(1) = \int_{-2}^1 5t^4 \, dt = (1)^5 + 32 = 33$$


$$A(x) = \int_{-2}^x 5t^4 \, dt = x^5 + 32$$



$$A(2) = \int_{-2}^2 5t^4 \, dt = (2)^5 + 32 = 64$$



$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$



$$A(3) = \int_{-2}^3 5t^4 dt = (3)^5 + 32 = 275$$

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_a^x f(t) dt$ . What functions could  $A(x)$  be?

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_a^x f(t) dt$ . What functions could  $A(x)$  be?

- ▶  $A'(x) = f(x)$ .
- ▶ Guess a function with derivative  $f(x)$ :  $F(x)$ .
- ▶ Then  $A(x) = F(x) + C$  for some constant  $C$ .
- ▶ Also  $A(a) = 0$ , so  $0 = F(a) + C$ , so  $C = -F(a)$
- ▶ So,  $A(x) = F(x) - F(a)$

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .

$A(x) = \int_a^x f(t) dt$ . What functions could  $A(x)$  be?

- ▶  $A'(x) = f(x)$ .
- ▶ Guess a function with derivative  $f(x)$ :  $F(x)$ .
- ▶ Then  $A(x) = F(x) + C$  for some constant  $C$ .
- ▶ Also  $A(a) = 0$ , so  $0 = F(a) + C$ , so  $C = -F(a)$
- ▶ So,  $A(x) = F(x) - F(a)$

---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

Two clues for finding  $A(x) = \int_a^x f(t) dt$ :

- ▶ If  $A(x) = \int_a^x f(t) dt$ , then<sup>1</sup>  $A'(x) = f(x)$
- ▶ If  $F'(x) = A'(x)$ , then  $A(x) = F(x) + C$  for some constant  $C$ .


$A(b) = \int_a^b f(t) dt$ . What functions could  $A(b)$  be?

- ▶  $A'(x) = f(x)$ .
- ▶ Guess a function with derivative  $f(x)$ :  $F(x)$ .
- ▶ Then  $A(x) = F(x) + C$  for some constant  $C$ .
- ▶ Also  $A(a) = 0$ , so  $0 = F(a) + C$ , so  $C = -F(a)$
- ▶ So,  $A(b) = F(b) - F(a)$


---

<sup>1</sup>(as long as  $f(t)$  is continuous on  $[a, x]$ )

$$\int_a^b f(t) \, dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$




$$\int_a^b f(t) \, dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$



$$\int_{-2}^1 f(t) \, dt = F(1) - F(-2)$$


$$\int_a^b f(t) \, dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$



$$\int_{-2}^2 f(t) \, dt = F(2) - F(-2)$$



$$\int_a^b f(t) \, dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$



$$\int_{-3}^3 f(t) \, dt = F(3) - F(-3)$$

## Fundamental Theorem of Calculus, Part 2

Let  $F(x)$  be differentiable, defined, and continuous on the interval  $[a, b]$  with  $F'(x) = f(x)$  for all  $a < x < b$ . Then

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

## Fundamental Theorem of Calculus, Part 2

Let  $F(x)$  be differentiable, defined, and continuous on the interval  $[a, b]$  with  $F'(x) = f(x)$  for all  $a < x < b$ . Then

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Examples:

$$\frac{d}{dx} \{5x^7\} = 35x^6, \text{ so}$$

$$\int_0^3 35x^6 \, dx =$$

$$\frac{d}{dx} \{\tan x\} = \sec^2 x, \text{ so}$$

$$\int_0^{\pi/4} \sec^2 x \, dx =$$

## Fundamental Theorem of Calculus, Part 2

Let  $F(x)$  be differentiable, defined, and continuous on the interval  $[a, b]$  with  $F'(x) = f(x)$  for all  $a < x < b$ . Then

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Examples:

$\frac{d}{dx} \{5x^7\} = 35x^6$ , so

$$\int_0^3 35x^6 \, dx = 5x^7 \Big|_{x=3} - 5x^7 \Big|_{x=0} = 5(3^7) - 5(0^7) = 5 \cdot 3^7$$

$\frac{d}{dx} \{\tan x\} = \sec^2 x$ , so

$$\int_0^{\pi/4} \sec^2 x \, dx =$$

## Fundamental Theorem of Calculus, Part 2

Let  $F(x)$  be differentiable, defined, and continuous on the interval  $[a, b]$  with  $F'(x) = f(x)$  for all  $a < x < b$ . Then

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Examples:


$\frac{d}{dx} \{5x^7\} = 35x^6$ , so

$$\int_0^3 35x^6 \, dx = 5x^7 \Big|_{x=3} - 5x^7 \Big|_{x=0} = 5(3^7) - 5(0^7) = 5 \cdot 3^7$$

$\frac{d}{dx} \{\tan x\} = \sec^2 x$ , so


$$\int_0^{\pi/4} \sec^2 x \, dx = \tan x \Big|_{x=\pi/4} - \tan x \Big|_{x=0} = \tan(\pi/4) - \tan 0 = 1$$

$$\int_0^3 35x^6 \, dx = F(b) - F(a) \quad \text{where} \quad F(x) = 5x^7$$



$$\int_0^3 35x^6 \, dx = 5(3)^7 - 5(0)^7$$

$$\int_0^{\pi/4} \sec^2 x \, dx = F(b) - F(a) \quad \text{where} \quad F(x) = \tan x$$



$$\int_0^{\pi/4} \sec^2 x \, dx = \tan\left(\frac{\pi}{4}\right) - \tan 0 = 1$$

## RELEVANT VOCABULARY

### Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .



# RELEVANT VOCABULARY

## Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .

Examples:

The derivative of  $x^2$  is  $2x$ , so:  
 $x^2$  is an **antiderivative** of  $2x$ .

# RELEVANT VOCABULARY

## Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .

Examples:

The derivative of  $x^2$  is  $2x$ , so:  
 $x^2$  is an **antiderivative** of  $2x$ .

When  $x > 0$ , the derivative of  $\log x$  is  $\frac{1}{x}$ , so:

# RELEVANT VOCABULARY

## Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .

Examples:

The derivative of  $x^2$  is  $2x$ , so:  
 $x^2$  is an **antiderivative** of  $2x$ .

When  $x > 0$ , the derivative of  $\log x$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log x$ .

# RELEVANT VOCABULARY

## Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .

Examples:

The derivative of  $x^2$  is  $2x$ , so:  
 $x^2$  is an **antiderivative** of  $2x$ .

When  $x > 0$ , the derivative of  $\log x$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log x$ .

For all  $x$ , the derivative of  $\log |x|$  is  $\frac{1}{x}$ , so:

# RELEVANT VOCABULARY

## Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .

Examples:

The derivative of  $x^2$  is  $2x$ , so:  
 $x^2$  is an **antiderivative** of  $2x$ .

When  $x > 0$ , the derivative of  $\log x$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log x$ .

For all  $x$ , the derivative of  $\log |x|$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log |x|$ .

# RELEVANT VOCABULARY

## Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .

Examples:

The derivative of  $x^2$  is  $2x$ , so:  
 $x^2$  is an **antiderivative** of  $2x$ .

When  $x > 0$ , the derivative of  $\log x$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log x$ .

For all  $x$ , the derivative of  $\log |x|$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log |x|$ .

An antiderivative of  $\sin x$  is



# RELEVANT VOCABULARY

## Definition

If  $F(x)$  is a function whose derivative is  $f(x)$ , we call  $F(x)$  an **antiderivative** of  $f(x)$ .

Examples:

The derivative of  $x^2$  is  $2x$ , so:  
 $x^2$  is an **antiderivative** of  $2x$ .

When  $x > 0$ , the derivative of  $\log x$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log x$ .

For all  $x$ , the derivative of  $\log |x|$  is  $\frac{1}{x}$ , so:  
 $\frac{1}{x}$  is an **antiderivative** of  $\log |x|$ .

An antiderivative of  $\sin x$  is  $-\cos x$ , because  $\frac{d}{dx} \{-\cos x\} = \sin x$ .



# CONVENIENT NOTATION

## Definition

$$f(x) \Big|_a^b = f(b) - f(a)$$

The function  $f(x)$  evaluated from  $a$  to  $b$



# CONVENIENT NOTATION

## Definition

$$f(x) \Big|_a^b = f(b) - f(a)$$

The function  $f(x)$  evaluated from  $a$  to  $b$

Examples:

$$(5x + x^2) \Big|_1^2 =$$

# CONVENIENT NOTATION

## Definition

$$f(x) \Big|_a^b = f(b) - f(a)$$

The function  $f(x)$  evaluated from  $a$  to  $b$

Examples:

$$(5x + x^2) \Big|_1^2 = (10 + 4) - (5 + 1)$$

# CONVENIENT NOTATION

## Definition

$$f(x) \Big|_a^b = f(b) - f(a)$$

The function  $f(x)$  evaluated from  $a$  to  $b$

Examples:

$$(5x + x^2) \Big|_1^2 = (10 + 4) - (5 + 1)$$

$$\frac{x^2}{x+2} \Big|_5^{-1} =$$

# CONVENIENT NOTATION

## Definition

$$f(x) \Big|_a^b = f(b) - f(a)$$

The function  $f(x)$  evaluated from  $a$  to  $b$

Examples:

$$(5x + x^2) \Big|_1^2 = (10 + 4) - (5 + 1)$$

$$\frac{x^2}{x+2} \Big|_5^{-1} = \frac{1}{1} - \frac{25}{7}$$

# CONVENIENT NOTATION

## Definition

$$f(x) \Big|_a^b = f(b) - f(a)$$

The function  $f(x)$  evaluated from  $a$  to  $b$

## FTC Part 2, Abridged

$$\int_a^b f(x) \, dx = F(x) \Big|_a^b$$

where  $F(x)$  is an antiderivative of  $f(x)$

## Definition

The **indefinite integral** of a function  $f(x)$ :

$$\int f(x) \, dx$$

means the *most general* antiderivative of  $f(x)$ .

Examples:

$$\int 2x \, dx =$$

## Definition

The **indefinite integral** of a function  $f(x)$ :

$$\int f(x) \, dx$$

means the *most general* antiderivative of  $f(x)$ .

Examples:

$$\int 2x \, dx = x^2 + C,$$

$C$  “arbitrary constant.”

## Definition

The **indefinite integral** of a function  $f(x)$ :

$$\int f(x) \, dx$$

means the *most general* antiderivative of  $f(x)$ .

Examples:

$$\int 2x \, dx = x^2 + C,$$

$C$  “arbitrary constant.”

$$\int \frac{1}{x} \, dx =$$



## Definition

The **indefinite integral** of a function  $f(x)$ :

$$\int f(x) \, dx$$

means the *most general* antiderivative of  $f(x)$ .

Examples:

$$\int 2x \, dx = x^2 + C, \quad C \text{ "arbitrary constant."}$$

$$\int \frac{1}{x} \, dx = \log |x| + C$$

Remember: two functions with the same derivative differ by a constant, so we include the “+C” for indefinite integrals.

# DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	
Limits (or bounds) of integration, $\int_a^b f(x) dx$	
Area under a curve	
Antiderivative	
Number	
Function	

# DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	<b>indefinite</b>
Limits (or bounds) of integration, $\int_a^b f(x) dx$	
Area under a curve	
Antiderivative	
Number	
Function	

# DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	indefinite
Limits (or bounds) of integration, $\int_a^b f(x) dx$	definite
Area under a curve	
Antiderivative	
Number	
Function	

# DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	indefinite
Limits (or bounds) of integration, $\int_a^b f(x) dx$	definite
Area under a curve	definite
Antiderivative	
Number	
Function	

# DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	indefinite
Limits (or bounds) of integration, $\int_a^b f(x) dx$	definite
Area under a curve	definite
Antiderivative	indefinite
Number	
Function	

# DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	indefinite
Limits (or bounds) of integration, $\int_a^b f(x) dx$	definite
Area under a curve	definite
Antiderivative	indefinite
Number	definite
Function	

# DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	indefinite
Limits (or bounds) of integration, $\int_a^b f(x) dx$	definite
Area under a curve	definite
Antiderivative	indefinite
Number	definite
Function	indefinite



# ANTIDIFFERENTIATION BY INSPECTION

1.  $\int e^x \, dx$

# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

# ANTIDIFFERENTIATION BY INSPECTION

1.  $\int e^x \, dx = e^x + C$

2.  $\int \cos x \, dx$

# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x \, dx = e^x + C$$

$$2. \int \cos x \, dx = \sin x + C$$

# ANTIDIFFERENTIATION BY INSPECTION

1.  $\int e^x \, dx = e^x + C$

2.  $\int \cos x \, dx = \sin x + C$

3.  $\int -\sin x \, dx$

# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x \, dx = e^x + C$$

$$2. \int \cos x \, dx = \sin x + C$$

$$3. \int -\sin x \, dx = \cos x + C$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

$$2. \int \cos x dx = \sin x + C$$

$$3. \int -\sin x dx = \cos x + C$$

$$4. \int \frac{1}{x} dx$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x \, dx = e^x + C$$

$$2. \int \cos x \, dx = \sin x + C$$

$$3. \int -\sin x \, dx = \cos x + C$$

$$4. \int \frac{1}{x} \, dx = \log |x| + C$$





# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x \, dx = e^x + C$$

$$2. \int \cos x \, dx = \sin x + C$$

$$3. \int -\sin x \, dx = \cos x + C$$

$$4. \int \frac{1}{x} \, dx = \log |x| + C$$

$$5. \int 1 \, dx$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x \, dx = e^x + C$$

$$2. \int \cos x \, dx = \sin x + C$$

$$3. \int -\sin x \, dx = \cos x + C$$

$$4. \int \frac{1}{x} \, dx = \log |x| + C$$

$$5. \int 1 \, dx = x + C$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

$$2. \int \cos x dx = \sin x + C$$

$$3. \int -\sin x dx = \cos x + C$$

$$4. \int \frac{1}{x} dx = \log |x| + C$$

$$5. \int 1 dx = x + C$$

$$6. \int 2x dx$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

$$2. \int \cos x dx = \sin x + C$$

$$3. \int -\sin x dx = \cos x + C$$

$$4. \int \frac{1}{x} dx = \log |x| + C$$

$$5. \int 1 dx = x + C$$

$$6. \int 2x dx = x^2 + C$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

$$2. \int \cos x dx = \sin x + C$$

$$3. \int -\sin x dx = \cos x + C$$

$$4. \int \frac{1}{x} dx = \log |x| + C$$

$$5. \int 1 dx = x + C$$

$$6. \int 2x dx = x^2 + C$$

$$7. \int nx^{n-1} dx \quad (n \neq 0, \text{ constant})$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

$$2. \int \cos x dx = \sin x + C$$

$$3. \int -\sin x dx = \cos x + C$$

$$4. \int \frac{1}{x} dx = \log |x| + C$$

$$5. \int 1 dx = x + C$$

$$6. \int 2x dx = x^2 + C$$

$$7. \int nx^{n-1} dx = x^n + C \quad (n \neq 0, \text{ constant})$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

$$2. \int \cos x dx = \sin x + C$$

$$3. \int -\sin x dx = \cos x + C$$

$$4. \int \frac{1}{x} dx = \log |x| + C$$

$$5. \int 1 dx = x + C$$

$$6. \int 2x dx = x^2 + C$$

$$7. \int nx^{n-1} dx = x^n + C \quad (n \neq 0, \text{ constant})$$

$$8. \int x^n dx \quad (n \neq -1, \text{ constant})$$



# ANTIDIFFERENTIATION BY INSPECTION

$$1. \int e^x dx = e^x + C$$

$$2. \int \cos x dx = \sin x + C$$

$$3. \int -\sin x dx = \cos x + C$$

$$4. \int \frac{1}{x} dx = \log |x| + C$$

$$5. \int 1 dx = x + C$$

$$6. \int 2x dx = x^2 + C$$

$$7. \int nx^{n-1} dx = x^n + C \quad (n \neq 0, \text{ constant})$$

$$8. \int x^n dx = \frac{1}{n+1}x^{n+1} + C \quad (n \neq -1, \text{ constant})$$





## Power Rule for Antidifferentiation

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

if  $n \neq -1$  is a constant

Example:

$$\int (5x^2 - 15x + 3) dx =$$



## Power Rule for Antidifferentiation

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C$$

if  $n \neq -1$  is a constant

Example:

$$\int (5x^2 - 15x + 3) dx = \frac{5}{3}x^3 - \frac{15}{2}x^2 + 3x + C$$



# ANTIDERIVATIVES TO RECOGNIZE

- ▶  $\int x^n dx = \frac{1}{n+1}x^{n+1} + C$
- ▶  $\int a dx = ax + C$
- ▶  $\int e^x dx = e^x + C$
- ▶  $\int \frac{1}{x} dx = \log |x| + C$
- ▶  $\int \sin x dx = -\cos x + C$
- ▶  $\int \cos x dx = \sin x + C$
- ▶  $\int \sec^2 x dx = \tan x + C$
- ▶  $\int \sec x \tan x dx = \sec x + C$
- ▶  $\int \csc x \cot x dx = -\csc x + C$
- ▶  $\int \csc^2 x dx = -\cot x + C$
- ▶  $\int \frac{1}{1+x^2} dx = \arctan x + C$
- ▶  $\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + C$

## Included Work



'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021, modified),