


TABLE OF CONTENTS



WARMUP

Evaluate $\int_3^7 \frac{1}{\sqrt{x^2 + 2x + 1}} dx$.

Evaluate $\int \frac{1}{\sqrt{x^2 + 1}} dx$.

CHECK OUR WORK

Let's verify that $\int \frac{1}{\sqrt{x^2 + 1}} = \log \left| \sqrt{x^2 + 1} + x \right| + C.$

Seems improbable, right?

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into $\sqrt{\cos^2 \theta} = |\cos \theta|$

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into $\sqrt{\cos^2 \theta} = |\cos \theta|$
 - ▶ $x = \tan \theta$, $1 + \tan^2 \theta = \sec^2 \theta$ changes $\sqrt{1 + x^2}$ into

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into $\sqrt{\cos^2 \theta} = |\cos \theta|$
 - ▶ $x = \tan \theta$, $1 + \tan^2 \theta = \sec^2 \theta$ changes $\sqrt{1 + x^2}$ into $\sqrt{\sec^2 \theta} = |\sec \theta|$

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into $\sqrt{\cos^2 \theta} = |\cos \theta|$
 - ▶ $x = \tan \theta$, $1 + \tan^2 \theta = \sec^2 \theta$ changes $\sqrt{1 + x^2}$ into $\sqrt{\sec^2 \theta} = |\sec \theta|$
 - ▶ $x = \sec \theta$, $\sec^2 \theta - 1 = \tan^2 \theta$ changes $\sqrt{x^2 - 1}$ into

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into $\sqrt{\cos^2 \theta} = |\cos \theta|$
 - ▶ $x = \tan \theta$, $1 + \tan^2 \theta = \sec^2 \theta$ changes $\sqrt{1 + x^2}$ into $\sqrt{\sec^2 \theta} = |\sec \theta|$
 - ▶ $x = \sec \theta$, $\sec^2 \theta - 1 = \tan^2 \theta$ changes $\sqrt{x^2 - 1}$ into $\sqrt{\tan^2 \theta} = |\tan \theta|$

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into $\sqrt{\cos^2 \theta} = |\cos \theta|$
 - ▶ $x = \tan \theta$, $1 + \tan^2 \theta = \sec^2 \theta$ changes $\sqrt{1 + x^2}$ into $\sqrt{\sec^2 \theta} = |\sec \theta|$
 - ▶ $x = \sec \theta$, $\sec^2 \theta - 1 = \tan^2 \theta$ changes $\sqrt{x^2 - 1}$ into $\sqrt{\tan^2 \theta} = |\tan \theta|$
- ▶ After integrating, convert back to the original variable (possibly using a triangle—more details later)

FOCUS ON THE ALGEBRA

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

Choose a trigonometric substitution that will allow the square root to cancel out of the following expressions:

▶ $\sqrt{x^2 - 1}$

▶ $\sqrt{x^2 + 1}$

▶ $\sqrt{1 - x^2}$



FOCUS ON THE ALGEBRA

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

Choose a trigonometric substitution that will allow the square root to cancel out of the following expressions:

► $\sqrt{x^2 + 7}$

► $\sqrt{3 - 2x^2}$



CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Consider the substitution $x = \sin \theta$, $dx = \cos \theta \, d\theta$ for the integral:

$$\int_0^1 \sqrt{1-x^2} \, dx$$

When $x = 0$ (lower limit of integration), what is θ ?

When $x = 1$ (upper limit of integration), what is θ ?



CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

More generally, suppose a is a positive constant and we use the substitution $x = a \sin \theta$ for the term $\sqrt{a^2 - x^2}$.

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

More generally, suppose a is a positive constant and we use the substitution $x = a \sin \theta$ for the term $\sqrt{a^2 - x^2}$.

▶ $\theta = \arcsin\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

More generally, suppose a is a positive constant and we use the substitution $x = a \sin \theta$ for the term $\sqrt{a^2 - x^2}$.


- ▶ $\theta = \arcsin\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶ $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|$

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

More generally, suppose a is a positive constant and we use the substitution $x = a \sin \theta$ for the term $\sqrt{a^2 - x^2}$.

- ▶ $\theta = \arcsin\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶ $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|$
- ▶ On the interval $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$, $\cos \theta \geq 0$, so $|\cos \theta| = \cos \theta$




CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

More generally, suppose a is a positive constant and we use the substitution $x = a \sin \theta$ for the term $\sqrt{a^2 - x^2}$.

- ▶ $\theta = \arcsin\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶ $\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2 \cos^2 \theta} = a |\cos \theta|$
- ▶ On the interval $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$, $\cos \theta \geq 0$, so $|\cos \theta| = \cos \theta$



- ▶ So, in general, when we use the substitution $x = \sin \theta$ with trigonometric substitution, we can expect $|\cos \theta| = \cos \theta$.

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Now, consider the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, where a is a positive constant.

CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Now, consider the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, where a is a positive constant.

► $\theta = \arctan\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Now, consider the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, where a is a positive constant.


- ▶ $\theta = \arctan\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶ $\sqrt{a^2 + x^2} = \sqrt{a^2 + a^2 \tan^2 \theta} = \sqrt{a^2 \sec^2 \theta} = \frac{a}{|\cos \theta|}$

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Now, consider the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, where a is a positive constant.

- ▶ $\theta = \arctan\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶ $\sqrt{a^2 + x^2} = \sqrt{a^2 + a^2 \tan^2 \theta} = \sqrt{a^2 \sec^2 \theta} = \frac{a}{|\cos \theta|}$
- ▶ On the interval $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$, $\cos \theta \geq 0$, so $|\cos \theta| = \cos \theta$ and $|\sec \theta| = \sec \theta$.




CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Now, consider the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, where a is a positive constant.

- ▶ $\theta = \arctan\left(\frac{x}{a}\right)$, so $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$
- ▶ $\sqrt{a^2 + x^2} = \sqrt{a^2 + a^2 \tan^2 \theta} = \sqrt{a^2 \sec^2 \theta} = \frac{a}{|\cos \theta|}$
- ▶ On the interval $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$, $\cos \theta \geq 0$, so $|\cos \theta| = \cos \theta$ and $|\sec \theta| = \sec \theta$.



- ▶ So, in general, when we use the substitution $x = \tan \theta$ with trigonometric substitution, we can expect $|\sec \theta| = \sec \theta$.

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.

CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.

► $\sec \theta = \frac{x}{a}$, so $\cos \theta = \frac{a}{x}$, so $\theta = \arccos\left(\frac{a}{x}\right)$. Then $0 \leq \theta \leq \pi$

CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.

- ▶ $\sec \theta = \frac{x}{a}$, so $\cos \theta = \frac{a}{x}$, so $\theta = \arccos\left(\frac{a}{x}\right)$. Then $0 \leq \theta \leq \pi$
- ▶ $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$

CLOSER LOOK AT ABSOLUTE VALUES

[▶ SKIP CLOSER LOOK](#)

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.

- ▶ $\sec \theta = \frac{x}{a}$, so $\cos \theta = \frac{a}{x}$, so $\theta = \arccos\left(\frac{a}{x}\right)$. Then $0 \leq \theta \leq \pi$
- ▶ $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.

- ▶ $\sec \theta = \frac{x}{a}$, so $\cos \theta = \frac{a}{x}$, so $\theta = \arccos\left(\frac{a}{x}\right)$. Then $0 \leq \theta \leq \pi$
- ▶ $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:
 - ▶ For $\sqrt{x^2 - a^2}$ to be defined, we need $x^2 \geq a^2$. I.e. $x \geq a$ or $x \leq -a$.

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.



- ▶ $\sec \theta = \frac{x}{a}$, so $\cos \theta = \frac{a}{x}$, so $\theta = \arccos\left(\frac{a}{x}\right)$. Then $0 \leq \theta \leq \pi$
- ▶ $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:
 - ▶ For $\sqrt{x^2 - a^2}$ to be defined, we need $x^2 \geq a^2$. I.e. $x \geq a$ or $x \leq -a$.
 - ▶ When $x \geq a$, we have $0 \leq \theta < \frac{\pi}{2}$, $\tan \theta \geq 0$, so $|\tan \theta| = \tan \theta$.

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.

- ▶ $\sec \theta = \frac{x}{a}$, so $\cos \theta = \frac{a}{x}$, so $\theta = \arccos\left(\frac{a}{x}\right)$. Then $0 \leq \theta \leq \pi$
- ▶ $\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a|\tan \theta|$
- ▶ Now this case gets slightly more complicated than the others:
 - ▶ For $\sqrt{x^2 - a^2}$ to be defined, we need $x^2 \geq a^2$. I.e. $x \geq a$ or $x \leq -a$.
 - ▶ When $x \geq a$, we have $0 \leq \theta < \frac{\pi}{2}$, $\tan \theta \geq 0$, so $|\tan \theta| = \tan \theta$.
 - ▶ When $x \leq -a$, we have $\frac{\pi}{2} < \theta \leq \pi$, $\tan \theta < 0$, so $|\tan \theta| = -\tan \theta$.



ABSOLUTE VALUES

From now on, we will assume:

- ▶ With the substitution $x = a \sin \theta$ for $\sqrt{a^2 - x^2}$, $|\cos \theta| = \cos \theta$
- ▶ With the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, $|\sec \theta| = \sec \theta$

Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate $\int_0^1 (1 + x^2)^{-3/2} dx$

Evaluate $\int_0^1 (1+x^2)^{-3/2} dx$

Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate $\int \sqrt{1 - 4x^2} \, dx$

Evaluate $\int \sqrt{1 - 4x^2} \, dx$

CHECK OUR WORK

In the last example, we computed

$$\int \sqrt{1 - 4x^2} \, dx = \frac{1}{4} (\arcsin(2x) + 2x\sqrt{1 - 4x^2}) + C.$$

To check, we differentiate.

Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate $\int \frac{1}{\sqrt{x^2 - 1}} dx$

Evaluate $\int \frac{1}{\sqrt{x^2 - 1}} dx$

CHECK OUR WORK

Let's check our result, $\int \frac{1}{\sqrt{x^2 - 1}} dx = \log \left| x + \sqrt{x^2 - 1} \right| + C.$

COMPLETING THE SQUARE

Choose a trigonometric substitution to simplify $\sqrt{3 - x^2 + 2x}$.

Identities have two “parts” that turn into one part:

- ▶ $1 - \sin^2 \theta = \cos^2 \theta$
- ▶ $1 + \tan^2 \theta = \sec^2 \theta$
- ▶ $\sec^2 \theta - 1 = \tan^2 \theta$

But our quadratic expression has *three* parts.

COMPLETING THE SQUARE

Choose a trigonometric substitution to simplify $\sqrt{3 - x^2 + 2x}$.

Identities have two “parts” that turn into one part:

- ▶ $1 - \sin^2 \theta = \cos^2 \theta$
- ▶ $1 + \tan^2 \theta = \sec^2 \theta$
- ▶ $\sec^2 \theta - 1 = \tan^2 \theta$

But our quadratic expression has *three* parts.

Fact: $3 - x^2 + 2x = 4 - (x - 1)^2$

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

1. Find b :

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

1. Find b : $-2bx = 2x$, so $b = -1$

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

1. Find b : $-2bx = 2x$, so $b = -1$

2. Solve for c :

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

1. Find b : $-2bx = 2x$, so $b = -1$

2. Solve for c : $3 = c - b^2 = c - 1$, so $c = 4$

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

1. Find b : $-2bx = 2x$, so $b = -1$
2. Solve for c : $3 = c - b^2 = c - 1$, so $c = 4$
3. All together:

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$

$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

1. Find b : $-2bx = 2x$, so $b = -1$

2. Solve for c : $3 = c - b^2 = c - 1$, so $c = 4$

3. All together: $3 - x^2 + 2x = 4 - (x - 1)^2$

Evaluate $\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} dx$.

Identities have two “parts” that turn into one part:

- ▶ $1 - \sin^2 \theta = \cos^2 \theta$
- ▶ $1 + \tan^2 \theta = \sec^2 \theta$
- ▶ $\sec^2 \theta - 1 = \tan^2 \theta$

One of those parts is a constant, and one is squared.

Evaluate $\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} dx = \int \frac{(x - 3)^2}{\sqrt{9 - (x - 3)^2}} dx.$

CHECK OUR WORK

Let's verify that

$$\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} = \frac{9}{2} \left(\arcsin \left(\frac{x-3}{3} \right) - \frac{x-3}{3} \cdot \frac{\sqrt{6x-x^2}}{3} \right) + C :$$

Included Work

 'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021, modified),
14–16

 'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021), 35, 37, 40,
52