


TABLE OF CONTENTS




SEQUENCES AND SERIES

A **sequence** is a list of numbers
A **series** is the sum of such a list.

Date	Time	Location	Observations


3/159

Size of Tiles: $\frac{1}{2}$




Covered Area: $\frac{1}{2}$

Size of Tiles: $\frac{1}{2}, \frac{1}{2^2}$




Covered Area: $\frac{1}{2} + \frac{1}{2^2}$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$




Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3}$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$




Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4}$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$, $\frac{1}{2^5}$




Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5}$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$, $\frac{1}{2^5}$, $\frac{1}{2^6}$




Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6}$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$, $\frac{1}{2^5}$, $\frac{1}{2^6}$, $\frac{1}{2^7}$




Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7}$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$, $\frac{1}{2^5}$, $\frac{1}{2^6}$, $\frac{1}{2^7}$, $\frac{1}{2^8}$



Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8}$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$, $\frac{1}{2^5}$, $\frac{1}{2^6}$, $\frac{1}{2^7}$, $\frac{1}{2^8}$, $\frac{1}{2^9}$




Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9}$

SEQUENCES AND SERIES

Size of Tiles:

$$\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \frac{1}{2^4}, \frac{1}{2^5}, \frac{1}{2^6}, \frac{1}{2^7}, \frac{1}{2^8}, \frac{1}{2^9}, \dots$$

Sequence




Covered Area:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$$

Size of Tiles: $\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \frac{1}{2^4}, \frac{1}{2^5}, \frac{1}{2^6}, \frac{1}{2^7}, \frac{1}{2^8}, \frac{1}{2^9}, \dots$


List of numbers,
approaching



Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$

Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$, $\frac{1}{2^5}$, $\frac{1}{2^6}$, $\frac{1}{2^7}$, $\frac{1}{2^8}$, $\frac{1}{2^9}$, $\dots$

**List of numbers,
approaching zero.**



Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$

Size of Tiles: $\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \frac{1}{2^4}, \frac{1}{2^5}, \frac{1}{2^6}, \frac{1}{2^7}, \frac{1}{2^8}, \frac{1}{2^9}, \dots$

**List of numbers,
approaching zero.**

Series

Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$

3.2 Series

[illegible]

3.2 Series

[illegible]

3.2 Series

[illegible]

3.2 Series

[illegible]

3.2 Series




Size of Tiles: $\frac{1}{2}$, $\frac{1}{2^2}$, $\frac{1}{2^3}$, $\frac{1}{2^4}$, $\frac{1}{2^5}$, $\frac{1}{2^6}$, $\frac{1}{2^7}$, $\frac{1}{2^8}$, $\frac{1}{2^9}$, $\dots$

**List of numbers,
approaching zero.**

Sum of numbers,
approaching one.

Covered Area: $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \frac{1}{2^8} + \frac{1}{2^9} + \cdots$



Recall:

$$\sum_{n=1}^5 \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2}$$

Recall:

$$\sum_{n=1}^5 \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2}$$

We informally interpret:

$$\sum_{n=1}^{\infty} \frac{1}{n^2} =$$

Recall:

$$\sum_{n=1}^5 \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2}$$

We informally interpret:

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \frac{1}{7^2} + \frac{1}{8^2} + \frac{1}{9^2} + \frac{1}{10^2} + \cdots$$

(a more rigorous definition will be discussed soon)

Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (C \cdot a_n) =$$

$$\text{A. } \sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$$

$$\text{B. } \sum_{n=1}^{\infty} C + \sum_{n=1}^{\infty} a_n$$

$$\text{C. } \sum_{n=1}^{\infty} a_n$$

$$\text{D. } a_n \sum_{n=1}^{\infty} C$$

E. none of the above

Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (C \cdot a_n) =$$

$$\text{A. } \sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$$

$$\text{B. } \sum_{n=1}^{\infty} C + \sum_{n=1}^{\infty} a_n$$

C. $C \sum_{n=1}^{\infty} a_n$

$$\text{D. } a_n \sum_{n=1}^{\infty} C$$

E. none of the above

Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n + b_n) =$$

$$\text{A. } \sum_{n=1}^{\infty} a_n \cdot \sum_{n=1}^{\infty} b_n$$

$$\text{B. } \sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n$$

C. $a_n + \sum_{n=1}^{\infty} b_n$

$$\text{D. } a_n \sum_{n=1}^{\infty} b_n$$

E. none of the above

Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n + b_n) =$$

$$\text{A. } \sum_{n=1}^{\infty} a_n \cdot \sum_{n=1}^{\infty} b_n$$

B. $\sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n$

C. $a_n + \sum_{n=1}^{\infty} b_n$

$$\text{D. } a_n \sum_{n=1}^{\infty} b_n$$

E. none of the above

Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n)^C =$$

$$\text{A. } \sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$$

B. $\left(\sum_{n=1}^{\infty} a_n\right)^C$

$$\text{C. } C^n \sum_{n=1}^{\infty} a_n$$

$$\text{D. } \sum_{n=1}^{\infty} C(a_n)^{C-1}$$

E. none of the above

Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n)^C =$$

$$\text{A. } \sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$$

B. $\left(\sum_{n=1}^{\infty} a_n\right)^C$

$$\text{C. } C^n \sum_{n=1}^{\infty} a_n$$

$$\text{D. } \sum_{n=1}^{\infty} C(a_n)^{C-1}$$

E. none of the above

What does it really mean to add up infinitely many things?

$$1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + \cdots$$

What does it really mean to add up infinitely many things?

$$\underbrace{1-1}_0 + \underbrace{1-1}_0 + \underbrace{1-1}_0 + \underbrace{1-1}_0 + \underbrace{1-1}_0 + \underbrace{1-1}_0 + \cdots$$

What does it really mean to add up infinitely many things?

$$\underbrace{1}_{1} - \underbrace{1+1}_{0} + \underbrace{1-1}_{0} + \underbrace{1-1}_{0} - \underbrace{1+1}_{0} + \underbrace{1-1}_{0} - \underbrace{1+1}_{0} + \cdots$$

What does it really mean to add up infinitely many things?



Diagram illustrating the construction of a sequence of 1s and -1s. The sequence is $1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + \dots$. Below the sequence, there are three '1's. Arrows show the construction: a vertical purple arrow from the first '1' to the first '1' in the sequence; a blue arrow from the second '1' to the second '-1' and a blue arrow from the third '1' to the third '1'; a purple arrow from the fourth '1' to the fourth '-1' and a purple arrow from the fifth '1' to the fifth '1'.

What does it really mean to add up infinitely many things?

We need an unambiguous definition.

HOW CAN WE ADD UP INFINITELY MANY THINGS?



SEQUENCE OF PARTIAL SUMS



HOW CAN WE ADD UP INFINITELY MANY THINGS?

SEQUENCE OF PARTIAL SUMS

$$S_1 = 0.2000$$




HOW CAN WE ADD UP INFINITELY MANY THINGS?

SEQUENCE OF PARTIAL SUMS

$$S_1 = 0.2000$$

$$S_2 = 0.2400$$





HOW CAN WE ADD UP INFINITELY MANY THINGS?

SEQUENCE OF PARTIAL SUMS

$$S_1 = 0.2000$$

$$S_2 = 0.2400$$

$$S_3 = 0.2480$$




SEQUENCE OF PARTIAL SUMS

$$S_4 = 0.2496$$



$$\frac{1}{55}$$



HOW CAN WE ADD UP INFINITELY MANY THINGS?

SEQUENCE OF PARTIAL SUMS


$$S_1 = 0.2000$$

$$S_2 = 0.2400$$

$$S_3 = 0.2480$$

$$S_4 = 0.2496$$

$$S_5 = 0.2499$$



Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04 \qquad S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04 \qquad S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04 \qquad S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008 \qquad S_3 = 0.248$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04 \qquad S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008 \qquad S_3 = 0.248$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04 \qquad S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008 \qquad S_3 = 0.248$$

$$a_4 = \frac{1}{5^4} = 0.0016 \qquad S_4 = 0.2496$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2 \qquad S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04 \qquad S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008 \qquad S_3 = 0.248$$

$$a_4 = \frac{1}{5^4} = 0.0016 \qquad S_4 = 0.2496$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2$$

$$S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$S_3 = 0.248$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$S_4 = 0.2496$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

$$S_5 = 0.24992$$

We define $\sum_{n=1}^{\infty} a_n = \lim_{N \rightarrow \infty} \sum_{n=1}^N a_n = \lim_{N \rightarrow \infty} S_N.$

$a_1 = \frac{1}{5} = 0.2$	$S_1 = 0.2$	$a_5 = \frac{1}{5^5} = 0.00032$	$S_5 = 0.24992$
$a_2 = \frac{1}{5^2} = 0.04$	$S_2 = 0.24$	$a_6 = \frac{1}{5^6} = 0.000064$	$S_6 = 0.249984$
$a_3 = \frac{1}{5^3} = 0.008$	$S_3 = 0.248$	$a_7 = \frac{1}{5^7} = 0.0000128$	$S_7 = 0.2499968$
$a_4 = \frac{1}{5^4} = 0.0016$	$S_4 = 0.2496$	$a_8 = \frac{1}{5^8} = 0.00000256$	$S_8 = 0.24999936$

From the sequence of partial sums, we guess

$$\sum_{n=1}^{\infty} \quad = \quad \lim_{N \rightarrow \infty} S_N \quad =$$


We define $\sum_{n=1}^{\infty} a_n = \lim_{N \rightarrow \infty} \sum_{n=1}^N a_n = \lim_{N \rightarrow \infty} S_N.$

$a_1 = \frac{1}{5} = 0.2$	$S_1 = 0.2$	$a_5 = \frac{1}{5^5} = 0.00032$	$S_5 = 0.24992$
$a_2 = \frac{1}{5^2} = 0.04$	$S_2 = 0.24$	$a_6 = \frac{1}{5^6} = 0.000064$	$S_6 = 0.249984$
$a_3 = \frac{1}{5^3} = 0.008$	$S_3 = 0.248$	$a_7 = \frac{1}{5^7} = 0.0000128$	$S_7 = 0.2499968$
$a_4 = \frac{1}{5^4} = 0.0016$	$S_4 = 0.2496$	$a_8 = \frac{1}{5^8} = 0.00000256$	$S_8 = 0.24999936$

From the sequence of partial sums, we guess



$$\sum_{n=1}^{\infty} \frac{1}{5^n} = \lim_{N \rightarrow \infty} S_N = \frac{1}{4}$$

NOTATION: $S_N = \sum_{n=1}^N a_n$

 a_1  a_2  a_3  a_4  a_5  a_6 

NOTATION: $S_N = \sum_{n=1}^N a_n$


$$S_1 = a_1$$



NOTATION: $S_N = \sum_{n=1}^N a_n$

$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$




NOTATION: $S_N = \sum_{n=1}^N a_n$

$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$




NOTATION: $S_N = \sum_{n=1}^N a_n$

$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

$$S_4 = a_1 + \cdots + a_4$$



NOTATION: $S_N = \sum_{n=1}^N a_n$


$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

$$S_4 = a_1 + \cdots + a_4$$

$$S_5 = a_1 + \cdots + a_5$$


 a_6


NOTATION: $S_N = \sum_{n=1}^N a_n$

$$S_1 = a_1$$


$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

$$S_4 = a_1 + \cdots + a_4$$

$$S_5 = a_1 + \cdots + a_5$$

$$S_6 = a_1 + \cdots + a_6$$



NOTATION PRACTICE

Suppose $\sum_{n=1}^{\infty} a_n$ has partial sums $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$.

► Evaluate $\sum_{n=1}^{100} a_n$.

► Evaluate $\sum_{n=1}^{\infty} a_n$.


NOTATION PRACTICE

Suppose $\sum_{n=1}^{\infty} a_n$ has partial sums $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$.

► Evaluate $\sum_{n=1}^{100} a_n$. $\sum_{n=1}^{100} a_n = S_{100} = \frac{100}{101}$


► Evaluate $\sum_{n=1}^{\infty} a_n$. $\sum_{n=1}^{\infty} a_n = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \frac{N}{N+1} = 1$

NOTATION PRACTICE



Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

NOTATION PRACTICE




Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.

How many cookies did each person bring?

NOTATION PRACTICE




Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.

How many cookies did each person bring?

Andrew brought 10, and Joel brought $19 - 10 = 9$.

NOTATION PRACTICE




Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.

How many cookies did each person bring?

Andrew brought 10, and Joel brought $19 - 10 = 9$.

NOTATION PRACTICE



Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.

How many cookies did each person bring?

Andrew brought 10, and Joel brought $19 - 10 = 9$.

NOTATION PRACTICE

Suppose $\sum_{n=1}^{\infty} a_n$ has partial sums $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$.

- Find a_1 .
- Give an explicit expression for a_n , when $n > 1$.

NOTATION PRACTICE

Suppose $\sum_{n=1}^{\infty} a_n$ has partial sums $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$.

- Find a_1 . $a_1 = \sum_{n=1}^1 a_n = S_1 = \frac{1}{2}$
- Give an explicit expression for a_n , when $n > 1$.

$$\begin{aligned} a_n &= \left(\sum_{k=1}^n a_k \right) - \left(\sum_{k=1}^{n-1} a_k \right) = S_n - S_{n-1} \\ &= \frac{n}{n+1} - \frac{n-1}{n} = \frac{1}{n(n+1)} \end{aligned}$$

oooooooooooo●oooooooooooooooooooo

$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$


 a_1

 a_2


 a_3

 a_4

 a_5


 a_6

 a_7

 a_8


$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$


$$S_1 = 1/(1+1)$$



$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$

$$S_1 = 1/(1+1)$$

$$S_2 = 2/(2+1)$$




$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$

$$S_1 = 1/(1+1)$$

$$S_2 = 2/(2+1)$$

$$S_3 = 3/(3+1)$$




$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$

$$S_1 = 1/(1+1)$$

$$S_2 = 2/(2+1)$$

$$S_3 = 3/(3+1)$$

$$S_4 = 4/(4+1)$$



Definition

The N^{th} partial sum of the series $\sum_{n=1}^{\infty} a_n$ is the sum of its first N terms

$$S_N = \sum_{n=1}^N a_n.$$

The partial sums form a sequence $\{S_N\}_{N=1}^{\infty}$. If this sequence of partial sums converges $S_N \rightarrow S$ as $N \rightarrow \infty$ then we say that the series $\sum_{n=1}^{\infty} a_n$ converges to S and we write

$$\sum_{n=1}^{\infty} a_n = S$$

If the sequence of partial sums diverges, we say that the series diverges.

Geometric Series

Let a and r be two fixed real numbers with $a \neq 0$. The series

$$a + ar + ar^2 + ar^3 + \cdots$$

is called the **geometric series** with first term a and ratio r .

We call r the *ratio* because it is the quotient of consecutive terms:

$$\frac{ar^{n+1}}{ar^n} = r$$

Geometric Series

Let a and r be two fixed real numbers with $a \neq 0$. The series

$$a + ar + ar^2 + ar^3 + \dots$$

is called the **geometric series** with first term a and ratio r .

We call r the *ratio* because it is the quotient of consecutive terms:

$$\frac{ar^{n+1}}{ar^n} = r$$

Another useful way of identifying geometric series is to determine whether all pairs of consecutive terms have the same ratio.

- ▶ Geometric: $1 + \frac{1}{5} + \frac{1}{5^2} + \frac{1}{5^3} + \frac{1}{5^4} + \dots$
- ▶ Geometric: $\sum_{n=0}^{\infty} \frac{1}{2^n}$
- ▶ Not geometric: $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \dots$

Consider the partial sum S_N of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \cdots + ar^N$$

Consider the partial sum S_N of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \cdots + ar^N$$

$$rS_N =$$

Consider the partial sum S_N of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \cdots + ar^N$$

$$rS_N = ar + ar^2 + ar^3 + \cdots + ar^N + ar^{N+1}$$

Consider the partial sum S_N of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \cdots + ar^N$$

$$rS_N = ar + ar^2 + ar^3 + \cdots + ar^N + ar^{N+1}$$

$$rS_N - S_N =$$

Consider the partial sum S_N of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \cdots + ar^N$$

$$rS_N = ar + ar^2 + ar^3 + \cdots + ar^N + ar^{N+1}$$

$$rS_N - S_N = -a + ar^{N+1}$$

Consider the partial sum S_N of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \cdots + ar^N$$

$$rS_N = \quad ar + ar^2 + ar^3 + \cdots + ar^N + ar^{N+1}$$

$$rS_N - S_N = \quad -a \quad \quad \quad + \quad \quad \quad ar^{N+1}$$

$$S_N(r - 1) = ar^{N+1} - a$$

If $r \neq 1$, then

$$S_N = \frac{ar^{N+1} - a}{r - 1} = a \frac{r^{N+1} - 1}{r - 1}$$

Geometric Series and Partial Sums

Let a and r be constants with $a \neq 0$, and let N be a natural number.

► If $r \neq 1$, then $a + ar + ar^2 + ar^3 + \cdots + ar^N = a \frac{r^{N+1} - 1}{r - 1}.$

► If $r = 1$, then $a + ar + ar^2 + ar^3 + \cdots + ar^N =$

► If $|r| < 1$, then $\sum_{n=0}^{\infty} ar^n =$

► If $r = 1$, then $\sum_{n=0}^{\infty} ar^n$

► If $r = -1$, then $\sum_{n=0}^{\infty} ar^n$

► If $|r| > 1$, then $\sum_{n=0}^{\infty} ar^n$

Geometric Series and Partial Sums

Let a and r be constants with $a \neq 0$, and let N be a natural number.

► If $r \neq 1$, then $a + ar + ar^2 + ar^3 + \cdots + ar^N = a \frac{r^{N+1} - 1}{r - 1}$.

► If $r = 1$, then $a + ar + ar^2 + ar^3 + \cdots + ar^N = (N + 1)a$.


► If $|r| < 1$, then $\sum_{n=0}^{\infty} ar^n = \lim_{N \rightarrow \infty} a \frac{r^{N+1} - 1}{r - 1} = a \frac{1}{1 - r}$

► If $r = 1$, then $\sum_{n=0}^{\infty} ar^n$ **diverges**

► If $r = -1$, then $\sum_{n=0}^{\infty} ar^n$ **diverges**


► If $|r| > 1$, then $\sum_{n=0}^{\infty} ar^n$ **diverges**

$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$



$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$


$$S_0 = a$$



$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$

$$S_0 = a$$

$$S_1 = 2a$$




$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$

$$S_0 = a$$

$$S_1 = 2a$$

$$S_2 = 3a$$





$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$

$$S_0 = a$$

$$S_1 = 2a$$

$$S_2 = 3a$$

$$S_3 = 4a$$



$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$


$$S_0 = a$$

$$S_1 = 2a$$

$$S_2 = 3a$$

$$S_3 = 4a$$

$$S_4 = 5a$$



$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$

$$S_0 = a$$


$$S_1 = 2a$$

$$S_2 = 3a$$


$$S_3 = 4a$$

$$S_4 = 5a$$

$$S_5 = 6a$$



$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$



$$S_0 = a$$


$$S_1 = 2a$$

$$S_2 = 3a$$



$$S_3 = 4a$$

$$S_4 = 5a$$

$$S_5 = 6a$$




$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$



$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$



$$S_0 = a$$



$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$

$$S_0 = a$$

$$S_1 = 0$$





$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$

$$S_0 = a$$

$$S_1 = 0$$

$$S_2 = a$$





$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$

$$S_0 = a$$

$$S_1 = 0$$

$$S_2 = a$$

$$S_3 = 0$$



$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$


$$S_0 = a$$

$$S_1 = 0$$

$$S_2 = a$$

$$S_3 = 0$$

$$S_4 = a$$



$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$

$$S_0 = a$$


$$S_1 = 0$$

$$S_2 = a$$


$$S_3 = 0$$

$$S_4 = a$$

$$S_5 = 0$$



$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$



$$S_0 = a$$


$$S_1 = 0$$

$$S_2 = a$$



$$S_3 = 0$$

$$S_4 = a$$

$$S_5 = 0$$





$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$



$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$

$$S_0 = a$$





○○○○○○○○○○○○○○○○○○○○●○○○○○○○○○


$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$

$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$




$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$



$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$




$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$

$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$



$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$


$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$


$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$



$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$




$$S_0 = a$$

$$S_1 = a \frac{r^2-1}{r-1}$$



$$S_2 = a \frac{r^3-1}{r-1}$$

$$S_3 = a \frac{r^4-1}{r-1}$$

$$S_4 = a \frac{r^5-1}{r-1}$$





$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$




$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$

$$S_0 = a$$




$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$




$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$




$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$




$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$



$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$




$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$



$$S_0 = a$$


$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$

$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$



$$S_0 = a$$

$$S_1 = a \frac{r^2 - 1}{r - 1}$$


$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$

$$S_5 = a \frac{r^6 - 1}{r - 1}$$

$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$



$$S_0 = a$$


$$S_1 = a \frac{r^2 - 1}{r - 1}$$

$$S_2 = a \frac{r^3 - 1}{r - 1}$$

$$S_3 = a \frac{r^4 - 1}{r - 1}$$

$$S_4 = a \frac{r^5 - 1}{r - 1}$$

$$S_5 = a \frac{r^6 - 1}{r - 1}$$



GEOMETRIC SERIES

New bitcoins are produced when a particular type of computational problem is solved. Every time 210,000 solutions are found, the number of bitcoins each solution can produce is cut in half.

- ▶ Each of the first 210,000 solutions can produce 50 bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2}$ bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2^2}$ bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2^3}$ bitcoins.

Assume that this continues forever, and that bitcoins are infinitely divisible.¹ How many bitcoins can possibly be produced?

¹Actually the smallest allowed division of a bitcoin is 10^{-8} .

GEOMETRIC SERIES

New bitcoins are produced when a particular type of computational problem is solved. Every time 210,000 solutions are found, the number of bitcoins each solution can produce is cut in half.

- ▶ Each of the first 210,000 solutions can produce 50 bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2}$ bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2^2}$ bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2^3}$ bitcoins.

Assume that this continues forever, and that bitcoins are infinitely divisible.¹ How many bitcoins can possibly be produced?

We start by writing the total number of bitcoin produced as a series. Since we want to know an upper bound, we'll assume that infinite solutions can be found and used to make bitcoin.

$$210\,000(50) + 210\,000\left(\frac{50}{2}\right) + 210\,000\left(\frac{50}{2^2}\right) + \cdots = \sum_{n=0}^{\infty} (210\,000) \left(\frac{50}{2^n}\right)$$

¹Actually the smallest allowed division of a bitcoin is 10^{-8} .

GEOMETRIC SERIES

$$\sum_{n=0}^{\infty} (210\,000) \left(\frac{50}{2^n} \right) =$$

GEOMETRIC SERIES

$$\begin{aligned}
 \sum_{n=0}^{\infty} (210\,000) \left(\frac{50}{2^n} \right) &= \sum_{n=0}^{\infty} (210\,000 \cdot 50) \left(\frac{1}{2} \right)^n \\
 &= (210\,000 \cdot 50) \frac{1}{1 - \frac{1}{2}} \\
 &= (210\,000 \cdot 50)(2) \\
 &= 21\,000\,000
 \end{aligned}$$

So there will never be more than 21,000,000 bitcoins produced this way.

$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n} \right) = 21\,000\,000$$



10 500 000



5 250 000




2 625 000



1 312 500



656 250



$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n} \right) = 21\,000\,000$$

$$S_0 = 10\,500\,000$$



5 250 000




2 625 000



1 312 500



656 250




10 500 000

$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n} \right) = 21\,000\,000$$

$$S_0 = 10\,500\,000$$

$$S_1 = 15\,750\,000$$



$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n} \right) = 21\,000\,000$$


$$S_0 = 10\,500\,000$$

$$S_1 = 15\,750\,000$$

$$S_2 = 18\,375\,000$$



 1 312 500 656 250



$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n} \right) = 21\,000\,000$$


$$S_0 = 10\,500\,000$$

$$S_1 = 15\,750\,000$$

$$S_2 = 18\,375\,000$$

$$S_3 = 19\,687\,500$$


656 250



$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n} \right) = 21\,000\,000$$


$$S_0 = 10\,500\,000$$

$$S_1 = 15\,750\,000$$

$$S_2 = 18\,375\,000$$

$$S_3 = 19\,687\,500$$

$$S_4 = 20\,343\,750$$



Arithmetic of Series

Let S , T , and C be real numbers. Let the two series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ converge to S and T respectively. Then

$$\sum_{n=1}^{\infty} [a_n + b_n] = S + T$$

$$\sum_{n=1}^{\infty} [a_n - b_n] = S - T$$

$$\sum_{n=1}^{\infty} [Ca_n] = CS$$

Geometric Series and Partial Sums

Let a and r be fixed numbers, and let N be a positive integer. Then

$$\sum_{n=0}^N ar^n = \begin{cases} a \cdot \frac{1-r^{N+1}}{1-r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate $\sum_{n=0}^{\infty} \left(\frac{2}{3^n} + \frac{4}{5^n} \right)$

$$\sum_{n=0}^{\infty} \left(\frac{2}{3^n} + \frac{4}{5^n} \right) =$$

$$\begin{aligned}\sum_{n=0}^{\infty} \left(\frac{2}{3^n} + \frac{4}{5^n} \right) &= \sum_{n=0}^{\infty} \frac{2}{3^n} + \sum_{n=0}^{\infty} \frac{4}{5^n} \\ &= \sum_{n=0}^{\infty} 2 \left(\frac{1}{3} \right)^n + \sum_{n=0}^{\infty} 4 \left(\frac{1}{5} \right)^n \\ &= \frac{2}{1 - \frac{1}{3}} + \frac{4}{1 - \frac{1}{5}} \\ &= \frac{2}{2/3} + \frac{4}{4/5} \\ &= 3 + 5 = 8\end{aligned}$$

Geometric Series and Partial Sums

Let a and r be fixed numbers, and let N be a positive integer. Then

$$\sum_{n=0}^N ar^n = \begin{cases} a \cdot \frac{1-r^{N+1}}{1-r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate $\sum_{n=6}^{\infty} \left(\frac{3^{n-1}}{5^{2n}} \right)$

$$\sum_{n=6}^{\infty} \left(\frac{3^{n-1}}{5^{2n}} \right) =$$

$$\sum_{n=6}^{\infty} \left(\frac{3^{n-1}}{5^{2n}} \right) = \sum_{n=6}^{\infty} \frac{1}{3} \left(\frac{3^n}{5^{2n}} \right) = \sum_{n=6}^{\infty} \frac{1}{3} \left(\frac{3^n}{25^n} \right) = \sum_{n=6}^{\infty} \frac{1}{3} \left(\frac{3}{25} \right)^n$$

Set $k = n - 6$. Then $n = k + 6$ and when $n = 6$, $k = 0$.

$$\begin{aligned} &= \sum_{k=0}^{\infty} \frac{1}{3} \left(\frac{3}{25} \right)^{k+6} = \sum_{k=1}^{\infty} \frac{1}{3} \left(\frac{3}{25} \right)^6 \left(\frac{3}{25} \right)^k \\ &= \frac{1}{3} \left(\frac{3}{25} \right)^6 \cdot \frac{1}{1 - 3/25} = \frac{3^5}{25^6} \cdot \frac{25}{22} = \frac{3^5}{25^5 \cdot 22} \end{aligned}$$

Geometric Series and Partial Sums

Let a and r be fixed numbers, and let N be a positive integer. Then

$$\sum_{n=0}^N ar^n = \begin{cases} a \cdot \frac{1-r^{N+1}}{1-r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate $\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right)$

$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) =$$


$$\begin{aligned}\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) &= \sum_{n=0}^{\infty} \left(\frac{4^n}{3^n} \right) \\ &= \sum_{n=1}^{\infty} \left(\frac{4}{3} \right)^n\end{aligned}$$

Since $\left| \frac{4}{3} \right| > 1$, the series diverges.

$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) \text{ diverges}$$





1

 $\frac{4}{3}$  $\frac{16}{9}$  $\frac{64}{27}$  $\frac{256}{81}$  $\frac{1024}{243}$ 

$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) \text{ diverges}$$



$$S_0 = 1.0000$$



$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) \text{ diverges}$$

$$S_0 = 1.0000$$

$$S_1 = 2.3333$$





$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) \text{ diverges}$$


$$S_0 = 1.0000$$

$$S_1 = 2.3333$$

$$S_2 = 4.1111$$



$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) \text{ diverges}$$




$$S_0 = 1.0000$$

$$S_1 = 2.3333$$

$$S_2 = 4.1111$$

$$S_3 = 6.4815$$



$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) \text{ diverges}$$



$$\frac{1024}{243}$$


$$S_0 = 1.0000$$

$$S_1 = 2.3333$$

$$S_2 = 4.1111$$

$$S_3 = 6.4815$$

$$S_4 = 9.6420$$



$$\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n} \right) \text{ diverges}$$

$$S_0 = 1.0000$$


$$S_1 = 2.3333$$

$$S_2 = 4.1111$$

$$S_3 = 6.4815$$

$$S_4 = 9.6420$$

$$S_5 = 13.856$$



TELESCOPING SUMS

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right).$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right).$

TELESCOPING SUMS

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right).$

$$\begin{array}{rclcl}
 a_1 : & \frac{1}{1} & - & \frac{1}{2} & \\
 a_2 : & \frac{1}{2} & - & \frac{1}{3} & \\
 a_3 : & \frac{1}{3} & - & \frac{1}{4} & \\
 a_4 : & \frac{1}{4} & - & \frac{1}{5} & \\
 & \vdots & & & \\
 a_{N-1} : & \frac{1}{N-1} & - & \frac{1}{N} & \\
 a_N : & \frac{1}{N} & - & \frac{1}{N+1} &
 \end{array}$$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right).$

$$S_1 = \frac{1}{1} - \frac{1}{2}$$

$$S_2 =$$

$$S_3 =$$

$$S_4 =$$

$$\vdots$$

$$S_N =$$

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{ccc} a_1 : & \frac{1}{1} & - & \frac{1}{2} \\ a_2 : & \frac{1}{2} & - & \frac{1}{3} \\ a_3 : & \frac{1}{3} & - & \frac{1}{4} \\ a_4 : & \frac{1}{4} & - & \frac{1}{5} \\ \vdots & & & \\ a_{N-1} : & \frac{1}{N-1} & - & \frac{1}{N} \\ a_N : & \frac{1}{N} & - & \frac{1}{N+1} \end{array}$$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rcl} S_1 = & \frac{1}{1} & - \quad \frac{1}{2} \\ S_2 = & \frac{1}{1} & - \quad \frac{1}{3} \\ S_3 = & & \\ S_4 = & & \\ \vdots & & \\ S_N = & & \end{array}$$

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} a_1 : & \frac{1}{1} & - & \frac{1}{2} \\ a_2 : & \frac{1}{2} & - & \frac{1}{3} \\ a_3 : & \frac{1}{3} & - & \frac{1}{4} \\ a_4 : & \frac{1}{4} & - & \frac{1}{5} \\ \vdots & & & \\ a_{N-1} : & \frac{1}{N-1} & - & \frac{1}{N} \\ a_N : & \frac{1}{N} & - & \frac{1}{N+1} \end{array}$$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rcl} S_1 & = & \frac{1}{1} \quad - \quad \frac{1}{2} \\ S_2 & = & \frac{1}{1} \quad - \quad \frac{1}{3} \\ S_3 & = & \frac{1}{1} \quad - \quad \frac{1}{4} \\ S_4 & = & \\ \vdots & & \\ S_N & = & \end{array}$$

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} a_1 : & \frac{1}{1} & - & \frac{1}{2} \\ a_2 : & \frac{1}{2} & - & \frac{1}{3} \\ a_3 : & \frac{1}{3} & - & \frac{1}{4} \\ a_4 : & \frac{1}{4} & - & \frac{1}{5} \\ & \vdots & & \\ a_{N-1} : & \frac{1}{N-1} & - & \frac{1}{N} \\ a_N : & \frac{1}{N} & - & \frac{1}{N+1} \end{array}$$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} S_1 & = & \frac{1}{1} & - & \frac{1}{2} \\ S_2 & = & \frac{1}{1} & - & \frac{1}{3} \\ S_3 & = & \frac{1}{1} & - & \frac{1}{4} \\ S_4 & = & \frac{1}{1} & - & \frac{1}{5} \\ & \vdots & & & \\ S_N & = & & & \end{array}$$

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{ccccc} a_1 : & \frac{1}{1} & - & \frac{1}{2} \\ a_2 : & \frac{1}{2} & - & \frac{1}{3} \\ a_3 : & \frac{1}{3} & - & \frac{1}{4} \\ a_4 : & \frac{1}{4} & - & \frac{1}{5} \\ \vdots & & & \\ a_{N-1} : & \frac{1}{N-1} & - & \frac{1}{N} \\ a_N : & \frac{1}{N} & - & \frac{1}{N+1} \end{array}$$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} S_1 & = & \frac{1}{1} & - & \frac{1}{2} \\ S_2 & = & \frac{1}{1} & - & \frac{1}{3} \\ S_3 & = & \frac{1}{1} & - & \frac{1}{4} \\ S_4 & = & \frac{1}{1} & - & \frac{1}{5} \\ & \vdots & & & \\ S_N & = & & & \end{array}$$

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} a_1 : & \frac{1}{1} & - & \frac{1}{2} \\ a_2 : & \frac{1}{2} & - & \frac{1}{3} \\ a_3 : & \frac{1}{3} & - & \frac{1}{4} \\ a_4 : & \frac{1}{4} & - & \frac{1}{5} \\ \vdots & & & \\ a_{N-1} : & \frac{1}{N-1} & - & \frac{1}{N} \\ a_N : & \frac{1}{N} & - & \frac{1}{N+1} \end{array}$$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} S_1 & = & \frac{1}{1} & - & \frac{1}{2} \\ S_2 & = & \frac{1}{1} & - & \frac{1}{3} \\ S_3 & = & \frac{1}{1} & - & \frac{1}{4} \\ S_4 & = & \frac{1}{1} & - & \frac{1}{5} \\ & \vdots & & & \\ S_N & = & \frac{1}{1} & - & \frac{1}{N+1} = \frac{N}{N+1} \end{array}$$

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} a_1 : & \frac{1}{1} & - & \frac{1}{2} \\ a_2 : & \frac{1}{2} & - & \frac{1}{3} \\ a_3 : & \frac{1}{3} & - & \frac{1}{4} \\ a_4 : & \frac{1}{4} & - & \frac{1}{5} \\ \vdots & & & \\ a_{N-1} : & \frac{1}{N-1} & - & \frac{1}{N} \\ a_N : & \frac{1}{N} & - & \frac{1}{N+1} \end{array}$$

$$\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1} \right) = S_{800} = \frac{800}{801}$$

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right)$.

$$\begin{array}{rclcl} S_1 & = & \frac{1}{1} & - & \frac{1}{2} \\ S_2 & = & \frac{1}{1} & - & \frac{1}{3} \\ S_3 & = & \frac{1}{1} & - & \frac{1}{4} \\ S_4 & = & \frac{1}{1} & - & \frac{1}{5} \\ & \vdots & & & \\ S_N & = & \frac{1}{1} & - & \frac{1}{N+1} = \frac{N}{N+1} \end{array}$$

$$\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right) = \lim_{N \rightarrow \infty} S_N = 1$$

Evaluate $\sum_{n=1}^{1000} \log \left(\frac{n+1}{n} \right) .$

Evaluate $\sum_{n=1}^{\infty} \log \left(\frac{n+1}{n} \right) .$

Evaluate $\sum_{n=1}^{1000} \log \left(\frac{n+1}{n} \right).$

Evaluate $\sum_{n=1}^{\infty} \log \left(\frac{n+1}{n} \right).$

$$a_1 : \log(2) - \log(1)$$

$$S_1 = \log(2)$$

$$a_2 : \log(3) - \log(2)$$

$$S_2 = \log(3)$$

$$a_3 : \log(4) - \log(3)$$

$$S_3 = \log(4)$$

$$\vdots$$

$$a_{n-1} : \log(n) - \log(n-1)$$

$$a_n : \log(n+1) - \log(n) \quad S_n = \log(n+1)$$

So, $\sum_{n=1}^{1000} \log \left(\frac{n+1}{n} \right) = S_{1000} = \log(1001)$ and

$$\sum_{n=1}^{\infty} \log \left(\frac{n+1}{n} \right) = \lim_{n \rightarrow \infty} \log(n+1) = \infty$$

$$\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right) \text{ diverges}$$



$$\log 2$$



$$\log \frac{3}{2}$$



$$\log \frac{4}{3}$$



$$\log \frac{5}{4}$$




$$\log \frac{6}{5}$$



$$\log \frac{7}{6}$$



$$\log \frac{8}{7}$$



$$\sum_{n=1}^{\infty} \log \left(\frac{n+1}{n} \right) \text{ diverges}$$

$$S_1 = 0.6931$$



$$\log \frac{3}{2}$$



$$\log \frac{4}{3}$$



$$\log \frac{5}{4}$$




$$\log \frac{6}{5}$$



$$\log \frac{7}{6}$$



$$\log \frac{8}{7}$$




$$\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right) \text{ diverges}$$

$$S_1 = 0.6931$$

$$S_2 = 1.0986$$

$$\begin{array}{ccccc} \text{[Weight Icon]} & \text{[Weight Icon]} & \text{[Weight Icon]} & \text{[Weight Icon]} & \text{[Weight Icon]} \\ \log \frac{4}{3} & \log \frac{5}{4} & \log \frac{6}{5} & \log \frac{7}{6} & \log \frac{8}{7} \end{array}$$



$$\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right) \text{ diverges}$$


$$S_1 = 0.6931$$

$$S_2 = 1.0986$$

$$S_3 = 1.3863$$



$$\log \frac{5}{4} \quad \log \frac{6}{5} \quad \log \frac{7}{6} \quad \log \frac{8}{7}$$



$$\sum_{n=1}^{\infty} \log \left(\frac{n+1}{n} \right) \text{ diverges}$$


$$S_1 = 0.6931$$

$$S_2 = 1.0986$$

$$S_3 = 1.3863$$

$$S_4 = 1.6094$$

$$\begin{array}{ccc} \text{[weight icon]} & \text{[weight icon]} & \text{[weight icon]} \\ \log \frac{6}{5} & \log \frac{7}{6} & \log \frac{8}{7} \end{array}$$



$$\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right) \text{ diverges}$$

$$S_1 = 0.6931$$


$$S_2 = 1.0986$$

$$S_3 = 1.3863$$

$$S_4 = 1.6094$$

$$S_5 = 1.7917$$

$$\begin{array}{cc} \text{[weight icon]} & \text{[weight icon]} \\ \log \frac{7}{6} & \log \frac{8}{7} \end{array}$$



$$\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right) \text{ diverges}$$

$$S_1 = 0.6931$$

$$S_2 = 1.0986$$


$$S_3 = 1.3863$$

$$S_4 = 1.6094$$

$$S_5 = 1.7917$$

$$S_6 = 1.9459$$

$$\log \frac{8}{7}$$



$$\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right) \text{ diverges}$$

$$S_1 = 0.6931$$

$$S_2 = 1.0986$$


$$S_3 = 1.3863$$

$$S_4 = 1.6094$$

$$S_5 = 1.7917$$

$$S_6 = 1.9459$$

$$S_7 = 2.0794$$



Included Work

📍 'Balloon' by [Simon Farkas](#) is licensed under [CC-BY](#) (accessed November 2022, edited), 92–99, 107–114

 'Waage/Libra' by [B. Lachner](#) is in the public domain (accessed April 2021, edited), 33–38, 52–58, 68–72, 84–114, 119–124, 135–141, 152–159

📦 'Weight' by [Kris Brauer](#) is licensed under [CC-BY](#) (accessed May 2021), 33–38, 52–58, 68–72, 84–114, 119–124, 135–141, 152–159

👤 'boy' by [Xinh Studio](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 61–65

🍪 'cookies' by [Azam Ishaq](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 61–65

🍪 'cookies' by [Vectors Point](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 61–65

🖱️ 'office desk' by [Abdul Baasith](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 61–65

👤 'Man' by [Xinh Studio](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 61–65

📖 'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021, modified), 142–149

📖 'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021), 116, 126, 129, 132