

## RAINFALL

Station BOWDEN HALL No. 24-459523  
 County GLOUCESTERSHIRE (Old No. \_\_\_\_\_)  
 Authority \_\_\_\_\_ Grid Ref. 50(32) 871151  
 Altitude 232 FT 714 ft  
 River Basin SEVERN  
 File \_\_\_\_\_ TF \_\_\_\_\_ / MWR / m / a Tributary Twyver

YEAR	1840	1841	1842	1843	1844	1845	1846	1847	1848	1849	Means
January ...				1.40	0.80	1.20	2.10	1.30	0.88	0.65	
February ...				2.50	1.40	0.37	0.70	0.60	2.70	1.62	
March ...				0.70	1.75	0.75	1.70	1.50	2.60	0.35	
April ...				1.80	0.60	0.90	3.50	2.20	2.80	3.00	
May ...				2.40	0.60	2.00	1.40	1.50	1.10	2.50	
June ...				2.50	0.80	2.60	1.20	2.60	4.20	1.33	
July ...				3.25	1.40	2.90	2.40	0.20	1.70	3.08	
August ...				4.20	3.10	2.60	3.30	1.00	3.10	1.14	
September ...				1.10	2.75	2.30	3.20	1.80	4.70	2.90	
October ...				4.70	2.30	1.60	4.50	4.00	3.80	2.10	
November ...				1.90	3.00	2.00	1.20	2.00	1.15	1.15	
December ...				0.60	0.20	1.60	0.50	2.10	2.00	1.70	
TOTAL ...				27.05	18.70	20.82	25.70	20.80	30.73	21.52	
Percentage ...											

## AVERAGE ANNUAL RAINFALL

Actual

Estimated from data for:-

1881-1915 in. \_\_\_\_\_

1916-1950 in. \_\_\_\_\_

\_\_\_\_\_ in. \_\_\_\_\_

Type and capacity of gauge \_\_\_\_\_

\*NOTES on unorthodox equipment or exposure, inspections, changes of equipment, site or altitude:-

8.12.76 9/L DATA EXTRACTED FROM GLOUCESTERSHIRE RAINFALL - AS HELPS "  
 PROCEEDINGS OF THE COTTESWOLD NATURALISTS FIELD CLUB  
 Vol XII PART II 1896-1897

Earliest year with complete data (D : W : M) 1843

\*NOTES on years with incomplete or unsatisfactory data:-

\*Each note to be dated. Continue notes overleaf if necessary.

## RAINFALL

Station BOWDEN HALLNo. 2A-459523County GLOUCESTERSHIRE

(Old No. \_\_\_\_\_)

Authority \_\_\_\_\_

Grid Ref. 50(32) 871151Altitude 232 FT 711 ft ftRiver Basin SEVERN

File \_\_\_\_\_ TP \_\_\_\_\_ / MWR / m / a

Tributary TWYVER

YEAR	1850	1851	1852	1853	1854	1855	1856	1857	1858	1859	Means
January ...	0.65	2.71	4.62	2.55	1.50		3.18	1.73			
February...	1.06	0.20	0.68	0.70	0.68		1.78	1.08			
March ...	0.72	3.00	0.15	0.62	0.40		0.88	1.82			
April ...	1.80	1.12	0.50	1.38	0.30		2.52	2.04			
May ...	3.82	1.36	1.36	2.60	2.10		2.86	1.96			
June ...	0.68	2.20	4.20	3.70	1.45		1.30	2.72			
July ...	4.60	2.30	3.30	2.58	3.90		0.82	2.30			
August ...	2.38	1.80	4.58	2.63	1.20		3.52	2.64			
September	1.60	0.45	3.68	1.91	1.10		2.70	2.04			
October ...	1.20	2.42	2.98	2.90	2.55		2.44	3.31			
November	1.80	0.40	5.52	2.10	0.84		0.46	1.20			
December	2.06	1.33	2.45	0.20	0.86		2.06	0.41			
TOTAL, ...	22.37	19.11	34.02	23.17	16.88		24.48	23.35			
Percentage											

## AVERAGE ANNUAL RAINFALL

Actual

Estimated from data for:-

1881-1915

in.

1916-1950

in.

in.

Type and capacity of gauge \_\_\_\_\_

\*NOTES on unorthodox equipment or exposure, inspections, changes of equipment, site or altitude:-

8.12.76 ORL DATA EXTRACTED FROM GLOUCESTERSHIRE RAINFALL - P.S. HELPS "

PROCEEDINGS OF THE COTTESWOLD NATURALISTS FIELD CLUB  
VOL XII PART II 1856-1857Earliest year with complete data (D : W : M) 1843

\*NOTES on years with incomplete or unsatisfactory data:-

\*Each note to be dated. Continue notes overleaf if necessary.