



WFP
World Food Programme

SAVING LIVES
CHANGING LIVES

WFP
wfp.org
vam
food security analysis

Global Seasonal Outlook

November 2025



1 HORN OF AFRICA — drought

The Deyr/short rains season has been delayed, with rainfall significantly below average, limiting planting and crop development. Dry conditions are expected to persist until mid-December, and any improvement may come too late to allow for meaningful vegetation recovery. Rainfall accumulation along the Shabelle–Juba basin remain low, raising concerns for irrigation and downstream water supplies

2 SOUTHERN AFRICA — flood — cyclone

La Niña conditions are expected to bring above-average rainfall and heightened flood risks, while northern areas contend with more variable weather patterns similar to those observed in the Horn of Africa. A normal to above-normal tropical cyclone season is also forecast

3 MENA — drought

Early-season rainfall remains below average across Syria, Iraq, Jordan, and northern Algeria, slowing moisture recovery. Ongoing water shortages and reduced upstream inflows continue to constrain irrigation and heighten seasonal water-stress risks in Middle East. Improved rainfall conditions are expected early next year and may support crop development during spring.

4 CENTRAL ASIA — drought

A dry start of the 2025/26 winter season is delaying crop planting and raising concerns. Forecasts indicate near-normal rainfall between December and February likely associated with a weakening La Niña

5 INDONESIA & PACIFIC — cyclone

The wet season began under La Niña with above-average rainfall, but forecasts point to a shift toward drier conditions ahead. Near to slightly above-normal tropical cyclone activity is also forecast for the region.

6 LATAM — flood — drought

Above-average rains marked the end of the Postrera rainy season, easing deficits from the Primera season but also causing floods. Record Hurricane Melissa brought severe flooding and damage across the Caribbeans. In South America, La Niña is driving wetter conditions in the north and drier outlooks in the south.

HORN OF AFRICA

Seasonal Rainfall Calendar

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
		HOA Long Rains, Ethiopia (Belg)							HOA Short Rains		


Seasonal Rainfall Deficits Deepen Across the Region

Across Horn of Africa, the Deyr/Short Rains season have been delayed and are significantly below average. October rainfall totals were generally less than 60 percent of normal across Somalia, southeastern Ethiopia, and eastern Kenya. As a result, the cropping season has been delayed, early crop establishment has been weak, pasture development has been poor, and the recharge of water pans and shallow wells has been limited.


Forecasts through the end of November indicate that deficits will persist and may even worsen. The medium-term forecast indicates that below-average rainfall will persist until mid-December, with recovery expected to begin in the latter part of the month. Unfortunately, the forecasted improvement will come too late to allow for meaningful vegetation recovery

These rainfall shortfalls reflect the combined influence of La Niña like event and a strong negative Indian Ocean Dipole, both of which are constraining moisture inflows into the region, while above-average temperatures are further increasing evapotranspiration.

Rainfall across the Shabelle–Juba basin in late October was about one standard deviation below the long-term average. Should this deficit continue, it may result in declining river levels and reduced irrigation capacity along riverine farming areas.



Rainfall **October 2025**, as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Observed and Rainfall forecast for the 1-month **until 30 Nov 2025** as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall

SOUTHERN AFRICA

Seasonal Rainfall Calendar

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Southern Africa												

Season Begins: Good Rains, Cyclone and Flood Risks Ahead


The start of the Southern Africa's 2025/26 rainy season has brought below-average rainfall in October, but good rains in November that are expected to continue through the month for most of the region. Northern parts of the region, however, is experiencing below average rainfall conditions, similar to Horn of Africa trends

La Nina conditions are present and typically bring above normal rainfall to Southern Africa. While this is usually associated with better crop production and hydropower generation, it also brings a heightened risk of flooding.


According to seasonal forecasts, the season is expected to bring normal to above-normal rainfall for most countries, though pockets of below-normal rains may occur in Angola, Namibia, Mozambique, DRC and Tanzania.

The SWIOCOF cyclone forecast for the season indicates normal to above average cyclonic activity across the Southwest Indian Ocean, suggesting the formation of 10-14 tropical systems, 5-8 reaching cyclone intensity. Madagascar and Mozambique are at highest risk, but Malawi, Tanzania and Zimbabwe have also been affected by cyclonic activity in the past.

Flood risk is expected along major river basins in the region, namely the Zambezi, Limpopo and Congo, as well as due to cyclonic activity in Madagascar and Mozambique. In addition, the risk of localized flash floods and associated landslides cannot be ruled out. As such, southern and central Mozambique, Madagascar, southeastern DRC, Republic of Congo and Malawi are at increased flood risk.



Observed and Rainfall forecast for the 1-month **until 30 Nov 2025** as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for the 3-month periods **Dec2025-Feb 2026** as the likelihood of exceeding the long-term median. Blue (orange) shades for likely wetter (drier) than usual conditions.

SPECIAL FOCUS: TROPICAL CYCLONES


Strong Late-Season Tropical Cyclones in the Caribbeans and Western Pacific

The final weeks of the 2025 tropical cyclones season in the Caribbeans and western Pacific were marked by an unusual concentration of high impact storms. Historically, storms of such intensity are rare during this period, making recent events particularly noteworthy

In the Caribbean, Hurricane Melissa delivered a devastating end to the Atlantic season. Making landfall in Jamaica as a Category 5 storm, Melissa caused widespread destruction across Jamaica, Cuba, Haiti and the Dominican Republic. Its slow movement prolonged exposure to extreme winds and rainfall, amplifying flooding, storm surge, and infrastructure damage.

In the western Pacific, Philippines faced a similarly intense late season Typhoons (Kalmaegi/Tino and Fung/Uwan) that struck the country in rapid succession bringing powerful winds, storm surge and widespread flood.

The timing and intensity of these storms are consistent with the influence of La Niña, which tends to favour late-season hurricane development in the Atlantic and shift cyclone formation westward toward Southeast Asia. While not the sole driver, La Niña likely contributed to the elevated activity observed during a period when cyclone frequency typically declines.




Warning for SW Indian Ocean cyclone season

The two plots illustrate wind speeds of historical tropical storms in the Caribbean (top) and Southern Africa (bottom) according to ENSO phase. In both regions, storm wind speeds are higher during La Niña years (blue area) compared to Neutral (white area) or El Niño phases (red area). This pattern highlights the amplifying effect of La Niña on cyclone intensity. For the upcoming Southwest Indian Ocean cyclone season, this serves as a warning: eastern coastal areas of Southern Africa and Madagascar may face heightened risks from stronger storms.

Data Source: IBTrACS (NOAA)

LATIN AMERICA & THE CARIBBEAN




End of Rainfall Season in Central America, Dry Start in South America

Central America and the Caribbean: The Postrera season is nearing its end, with above-average rainfall observed across much of Central America and parts of Caribbean through November, helping to ease earlier deficits from the Primera season. However excessive rainfall in the late October and early November triggered floods and landslides in several areas including Guatemala, El Salvador and Honduras.


Hurricane Melissa, the final storm of the 2025 Atlantic season, brought severe flooding and damage to Haiti, Jamaica and Cuba, marking a destructive close of the hurricane season.

As the region transitions into dry season in December, soil moisture remains adequate for late season crops, but localized floods damage and high temperatures may hinder recovery in some farming areas. Overall, the Postrera harvest is expected to conclude under generally favorable conditions, though Haiti, Jamaica and eastern Cuba remain affected by storm impacts and excess rainfall.


South America: The onset of the rainy season began in November with contrasting rainfall patterns typical of la Niña conditions. Forecasts for December 2025 to February 2026 indicate above-average rainfall across the Amazon basin and northern South America, which could sustain favorable crop growth but also heighten flood risk. In contrast southern South America are likely to experience below-average rainfall, signaling a drier start to the cropping season.



Rainfall observed and forecasted in 1 month ending **30 Nov 2025** as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall **Apr to Oct 2025**, as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for **Dec 25 - Feb 26** as the likelihood of exceeding the long- term median. Blue (orange) shades for likely wetter (drier) than usual condition

MENA REGION

Seasonal Rainfall Calendar


	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Mena Region												

High Likelihood of Continued Dry Conditions until early 2026


Across the MENA region October–November rainfall has been generally below average, limiting soil moisture recharge and slowing the start of the winter season in major rainfed wheat and barley producing areas. Dry conditions dominate across Syria, Iraq, Jordan, Turkey and northern Algeria.

Precipitation levels in the Tigris–Euphrates basin were well below normal last season. As a result, most water reservoirs in the region are at very low levels, increasing the vulnerability of agricultural activities in Iraq and Syria to any additional seasonal rainfall deficits

Seasonal forecasts suggest continued below-average rainfall through December and January, with recovery anticipated from February onward. This improvement could provide support for crop development during spring 2026.



Observed and forecast rainfall 1-month ending **30 Nov 2025** , as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for the 3-month periods **Dec 2025 - Feb 2026** as the likelihood of exceeding the long-term median. Blue (orange) shades for likely wetter (drier) than usual conditions

CENTRAL ASIA


Seasonal Rainfall Calendar

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Central Asia												

Dry Start Raises Concerns, but Improvement Likely Ahead


The 2025/2026 winter season has begun with below-average rainfall across much of the region, particularly in Afghanistan, Tajikistan, and Kyrgyzstan, where early-season dryness is delaying winter wheat planting in some rainfed areas. Reduced snow accumulation from the 2024/2025 winter season may continue to limit water availability for irrigation.

Looking ahead, forecasts suggest near-normal rainfall between December and February, likely associated with a weakening La Niña.




Rainfall for **Oct 2025**, as a proportion of the long- term average.

Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Observed and Rainfall forecast for the 1-month **until 30 Nov 2025** as the likelihood of exceeding the long-term median. Blue (orange) shades for likely wetter (drier) than usual conditions



Rainfall forecast for the 3-month periods **Dec 2025 - Feb 2026**, as the likelihood of exceeding the long-term median.Blue (orange) shades for likely wetter (drier) than usual conditions

Indonesia and Pacific

Seasonal Rainfall Calendar


Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Wet Start, Uncertain Outlook Ahead


The 2025/2026 wet season began under the La Niña conditions, bringing above-average rainfall across much of Indonesia and Papua New Guinea. Early rains have been beneficial for soil moisture and the start of wet season planting, progressing under mostly favorable conditions than last year.

However, seasonal forecasts for December-February indicate a shift toward near-normal to below-average rainfall across much of the region, suggesting a drier mid-season period that could negatively affect crops if it becomes prolonged. Overall agricultural prospects remain favorable in the short term supported by sufficient water availability.


Tropical cyclone activity is expected to remain near normal, with slightly above-normal activity possible, consistent with the ongoing La Niña influence.



Rainfall for **October 2025**, as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall.



Observed and Rainfall forecast for the 1-month **until 30 Nov 2025** as the likelihood of exceeding the long-term median. Blue (orange) shades for likely wetter (drier) than usual conditions



Rainfall forecast for the 3-month periods **Dec 2025 - Feb 2026**, as the likelihood of exceeding the long-term median.Blue (orange) shades for likely wetter (drier) than usual conditions

All satellite data (rainfall, vegetation index and land surface temperature) both current and historical are stored and processed at the WFP-HQ Humanitarian Data Cube AWS cloud system.

Rainfall: Primary dekadal data (CHIRPS) from Univ California St Barbara Climate Hazards Centre (https://www.chc.ucsb.edu/data/chirps)
Various accumulations are computed on a dekadal basis and anomalies derived relative to a 25 year-long mean (1994-2018).

Land Surface Temperature: Primary data from MODIS-AQUA (NASA), cloud cleared and gap-filled, both daytime and nighttime images. Thermal amplitude derived from difference between the two. Long term averages 2002-2018 used to derive anomalies

Vegetation Index: Primary data from MODIS-AQUA and TERRA (NASA), cloud cleared and gap-filled. Long term averages 2002-2018 used to derive anomalies

Seasonal Forecast: ECMWF seasonal forecasts **(SEAS5) |** The forecast anomalies are presented as the probability of exceeding the median, indicating the likelihood that seasonal precipitation will be above the median of the 24-year climatological distribution. These anomalies are derived from a 51-member forecast ensemble and are calculated relative to a 24-year model climatology (based on a 25-member ensemble reforecast) spanning 1993–2016

Water level in reservoirs: G-REALM data from USDA Foreign Agricultural Service.

Flood Forecast: GLOFAS-Copernicus and GOOGLE Flood Hub

Prepared by: Irene Ferrari, Diego Oliveira De Souza, Virginia Leape, Sara Miller, Giancarlo Pini

WFP World Food Programme

For further information please contact:

hq.geospatial@wfp.org