



WFP
World Food Programme

SAVING LIVES
CHANGING LIVES

WFP wfp.org | vam food security analysis

Global Seasonal Outlook

August 2025



KEY HIGHLIGHTS

ENSO OUTLOOK

ENSO-neutral conditions persist, but a transition to La Niña is likely during the northern hemisphere fall and early winter.

1 WESTERN AFRICA

drought | flood

Rainfall across the Sahel remains uneven. While the Sahelian belt has received average to above-average rainfall, persistent dryness in southern areas is causing vegetation stress in some regions. Flood risks remain elevated, with fluvial flooding likely in the coming weeks. Forecasts through the end of the month and into the next three months suggest continued below-average rainfall across the Gulf of Guinea and Atlantic coastal countries, while the Sahelian zone is expected to maintain average to above-average precipitation.

2 EASTERN AFRICA

flood | drought

Early-season dryness is improving. Recent heavy rainfall has led to flooding, and the risk of further events remains high in the coming weeks. In the Sudd wetlands, flood extent is projected to reach or surpass those observed in 2024. The seasonal forecast through November indicates above-average rainfall across much of the region supporting vegetation growth but heightens the risk of floods. Early forecasts indicate a potential for below-average rainfall in Kenya and Somalia during the upcoming Deyr/Short Rains season.

3 L. AMERICA & THE CARIBBEAN

drought | cyclone

Dry conditions persist across the Dry Corridor and parts of the Caribbean, disrupting the Primera season and impacting subsistence farming. However, forecasts through November suggest a return to typical rainfall, easing concerns for the Postrera season. Favorable rainfall has supported crop development in Colombia and Venezuela. The 2025 hurricane season has already brought storms near Haiti and the Dominican Republic, with forecasts indicating continued above-average cyclone activity.

4 SOUTH & SOUTHEAST ASIA

flood

The 2025 monsoon season has generally brought favorable rainfall, supporting crop development across key agricultural zones with just few localized concerns. Heavy rains have caused flooding and further floods are forecast in the coming weeks. The seasonal forecast through November indicates a continued likelihood of above-average rainfall, which may further support crop growth but also increases the risk of flooding and landslides.

WESTERN AFRICA

Seasonal Rainfall Calendar

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Sahel					▬	▬	▬	▬	▬	▬		
Gulf of Guinea			▬	▬	▬	▬	▬	▬	▬	▬	▬	

Flood risks in the Sahelian belt while Gulf of Guinea remains unusually dry


As the rainy season progresses across the Sahelian region, conditions remain mixed. Persistent dryness continues in southwestern Mali, Sierra Leone, Liberia, Guinea, and parts of Côte d'Ivoire, Ghana, southern Nigeria, southern Cameroon and northern Central Africa Republic. Signs of vegetation stress are emerging in southern Chad, central Nigeria, parts of Ghana and Côte d'Ivoire.

In contrast, the Sahelian belt has received average to above-average rainfall and these conditions have generally supported healthy vegetation growth in both croplands and pastures. Very high levels of accumulated rainfall were recorded over the past month in northern Burkina Faso, southwestern Niger, and central Chad. Localized flooding occurred in central Mali, along the Niger and Benue rivers in Nigeria, in southern Chad and along the Senegal river.


The forecast through the end of the month continues to indicate below-average rainfall across much of the Gulf of Guinea while the eastern Sahelian region show likely above-average precipitation.

Flood risk remains elevated in the coming weeks. Latest forecasts indicate a likelihood of fluvial flooding in SE Mali, NE and NW Nigeria, northern Senegal, Guinea Bissau and central-southern Chad.


Looking ahead, the three-month seasonal forecast indicates an increased likelihood of below-average rainfall across much of the Gulf of Guinea region, extending into southern Mali, Burkina Faso, Senegal, and Cameroon. In contrast, the Sahelian zone is expected to receive average rainfall, with some areas likely to experience above-average precipitation.



Rainfall **May 2025** to **20 Aug 2025**, as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for the 10 days **31 Aug 2025** as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for the 3-month **Sep - Nov 2025** as the likelihood of exceeding the long-term median. Blue (orange) shades for likely wetter (drier) than usual conditions

Seasonal Rainfall Calendar

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			■	■	■	■	■	■	■		

Eastern Africa
Sudans and Ethiopia (Meher)

Vegetation has improved but flood risks remain high

Early-season dryness in parts of Sudan, South Sudan, and Ethiopia has begun to ease, allowing vegetation to recover across most areas. However, areas in south-eastern Sudan*, northern Upper Nile in South Sudan and northern Ethiopia continue to experience limited improvements.

Recent rainfall has triggered flooding in several regions, particularly in eastern Sudan, Jonglei and Unity State in South Sudan.


Flood risk remains elevated in the coming weeks. Latest forecasts indicate a likelihood of fluvial flooding in eastern and south-western Sudan, northern Ethiopia and western Yemen.

Overall current flooding extent in South Sudan's Sudd wetlands is at record levels for this period. According to the latest forecast, it will likely match or exceed the levels observed in 2024. The flood extent is expected to peak between late September and October.


Looking ahead, the seasonal forecast through November indicates a high probability of above-average rainfall across much of the region—especially in South Sudan, Sudan, and central-northern Ethiopia. This is expected to support favorable vegetation growth but also heightens the risk of both riverine and flash floods due to intense rainfall events.

It is worth to note that early forecasts for the coming Deyr/Short Rains season in the Horn of Africa suggest a likelihood of below-average rainfall across much of Kenya and Somalia. This trend warrants close monitoring in the coming months, with updated forecasts expected to provide further insights.

* Insecurity is still affecting farming activities in most of the country




Rainfall **April 2025** to **20 August 2025**, as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for the 3-month periods **Sep-Nov 2025** as the likelihood of exceeding the long-term median. Blue (orange) shades for likely wetter (drier) than usual conditions

LATIN AMERICA & THE CARIBBEAN

Seasonal Rainfall Calendar




Stress continues in Primera Season while Postrera outlook is promising

Central America and the Caribbean: Over the past five months, rainfall patterns have been dominated by persistent dryness across northern areas and parts of the Caribbean, while scattered episodes of rains brought temporary relief in localized areas. These conditions have shaped the Primera season, where repeated dry spells and erratic rains constrained planting and crop growth, especially in Guatemala, El Salvador, western Honduras, Northern Nicaragua. Subsistence farming in the Dry Corridor are likely to be the most affected. Cuba and Haiti continue to face rainfall deficits, with signs of vegetation stress emerging in southwestern Haiti and eastern Cuba.


Looking ahead, the outlook through November indicates a return to more typical rainfall across most of the region, reducing concerns for the upcoming Postrera season. However, uncertainty remains around the potential for delayed planting. Haiti remains exposed to below-average rains increasing the risk of further crop stress.

The 2025 hurricane season has already seen Hurricane Erin brushing past Haiti, the Dominican Republic, and nearby islands, and two Pacific storms in northern Central America without making landfall. Forecasts still showing above-average cyclone activity through the remainder of the season, with continued risks for coastal and island territories.

Northern South America: Above-average rainfall has been the dominant feature, particularly in Colombia and Venezuela. While this has generally supported crop development, it also triggered localized flooding. Looking ahead, near-average conditions are expected, though occasional dry spells may occur in southern Venezuela. The prospects for the ongoing cropping season remain generally favorable.




Observed rainfall from **April to 20 Aug 2025** as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for **Sep-Nov 2025** as the likelihood of exceeding the long- term median. Blue (orange) shades for likely wetter (drier) than usual condition

SOUTH & SOUTHEAST ASIA




Favorable Monsoon Conditions Support Agriculture but Flood Risks Persist

The 2025 monsoon season has so far brought generally favorable rainfall across South and South-east Asia, supporting crop development in key agricultural zones. However, below-average rainfall in southern Myanmar, northern Laos, and eastern Cambodia raises some localized concerns.


Heavy rainfall events have affected several parts of the region, including Bangladesh, northeastern Pakistan, southern and central Myanmar, and southern Cambodia, some of them resulting in localized flooding.

Fluvial floods are forecast over the next couple of weeks in eastern Pakistan and central Myanmar.

Looking ahead, the seasonal forecast through November indicates a continued likelihood of above-average rainfall, particularly across the Philippines, DPRK, most of India, and the eastern parts of the region. While this may further benefit crop growth, it also raises the risk of flooding and landslides.



Rainfall **April 2025** to **20 August 2025**, as a proportion of the long-term average. Blue and purple (orange and brown) shades correspond to above (below) average rainfall



Rainfall forecast for the 3-month **Sep-Nov 2025** as the likelihood of exceeding the long-term median. Blue (orange) shades for likely wetter (drier) than usual conditions

All satellite data (rainfall, vegetation index and land surface temperature) both current and historical are stored and processed at the WFP-HQ Humanitarian Data Cube AWS cloud system.

Rainfall: Primary dekadal data (CHIRPS) from Univ California St Barbara Climate Hazards Centre (https://www.chc.ucsb.edu/data/chirps)
Various accumulations are computed on a dekadal basis and anomalies derived relative to a 25 year-long mean (1994-2018). Rainfall charts from WFP Seasonal Explorer (https://dataviz.vam.wfp.org/seasonal_explorer/rainfall_vegetation/visualizations)

Land Surface Temperature: Primary data from MODIS-AQUA (NASA), cloud cleared and gap-filled, both daytime and nighttime images. Thermal amplitude derived from difference between the two. Long term averages 2002-2018 used to derive anomalies

Vegetation Index: Primary data from MODIS-AQUA and TERRA (NASA), cloud cleared and gap-filled. Long term averages 2002-2018 used to derive anomalies

Seasonal Forecast: ECMWF seasonal forecasts (SEAS5) | The forecast anomalies are presented as the probability of exceeding the median, indicating the likelihood that seasonal precipitation will be above the median of the 24-year climatological distribution. These anomalies are derived from a 51-member forecast ensemble and are calculated relative to a 24-year model climatology (based on a 25-member ensemble reforecast) spanning 1993–2016

Water level in reservoirs: G-REALM data from USDA Foreign Agricultural Service.

Flood Forecast: GLOFAS-Copernicus and GOOGLE Flood Hub

Prepared by: Irene Ferrari, Diego Oliveira De Souza, Sirio Modugno, Giancarlo Pini

WFP World Food Programme

For further information please contact:

hq.geospatial@wfp.org