MICHIGAN MEDICINE
UNIVERSITY OF MICHIGAN

Criteria restricted antimicrobials in patients on adult services

Use of certain antimicrobial agents is restricted at Michigan Medicine. Agents are classified as Tier I or Tier II agents depending on whether Antimicrobial Stewardship Team (AST) approval is required prior to dispensing.

TIER II RESTRICTED ANTIMICROBIALS

Use of the following agents (i.e., Tier II agents) do not require approval prior to dispensing but is restricted to the criteria below. Even for the indications listed below, approved use is often limited to specific situations, such as drug allergy or certain risk factors. Other use requires approval from AST (pager #30780) or ID. Please consult appropriate treatment guidelines

Note: The below indications generally refer to appropriate EMPIRIC use. When cultures are available, antibiotic therapy should be escalated/de-escalated as appropriate based on organism and susceptibility. Restricted agents should only be utilized if narrower-spectrum agents are resistant or otherwise inappropriate. When cultures are not available, please refer to individual treatment guidelines for appropriate definitive therapy strategies.

UMHHC Policy 07-01-015 ("Use of Infectious Diseases Restricted Antimicrobials")

List of Tier II restricted Antimicrobials

Vancomycin	Clindamycin	Ciprofloxacin	Levofloxacin
Moxifloxacin	Ceftriaxone		

Empiric and definitive therapy: Tier II antimicrobials should be rarely used as first-line therapy unless patients have allergies, resistance, or pathogen-specific risk factors. Please refer to stewardship webpage for specific disease-state recommendations:

University of Michigan Health System (umich.edu)