
MICHIGAN MEDICINE
UNIVERSITY OF MICHIGAN

Antibiotic Treatment Guidelines for Group A Streptococcal Pharyngitis in Adult Patients

This guideline is designed to provide guidance for pharyngitis caused by group A Streptococcus (GAS) in most adults. Patients with a recent history of oropharyngeal or other deep neck space surgery or infection are excluded. Patients who have evidence of suppurative complications on exam (such as peritonsillar abscess) OR patients with suspected acute rheumatic fever will need additional management that is outside the scope of this guideline.

In adults, strep is the cause of pharyngitis < 15% of the time, so viral etiologies should be strongly considered.



Feature	Symptom	Score
Viral Symptoms	Absence of Cough	1
Cervical Adenopathy	Swollen tender anterior cervical nodes	1
Fever	>100.4°F (38°C)	1
Tonsillar Appearance	Tonsillar exudate or swelling	1
Age	3 y - 14 y 15 y - 44 y > 45 y	1 0 -1

†Testing for GAS in patients with concomitant symptoms of viral illness is not recommended as a positive result is of questionable diagnostic value. Rates of asymptomatic carriage of GAS are reported as high as 25% (AAP Red Book). In these cases risk of transmission and suppurative/nonsuppurative complications is low.

§High-risk individuals include: those with sore throat and household exposure to GAS (e.g.,living or sleeping in the same indoor shared space as a person diagnosed with GAS infection), a history of a previous rheumatic fever diagnosis, or symptoms or signs suggestive of complicated local or systemic GAS infection (e.g., peritonsillar or retropharyngeal abscess, scarlet fever and/or toxic shock syndrome).

Recommended Therapy	Duration/Comments
1st line: **Penicillin V** *Preferred*: **5 days:** Oral suspension: 800 mg PO QID Oral tablet/capsule: 750 mg PO QID *Alternative for risk factors (see comments):* **10 days**: 500 mg PO BID OR **Amoxicillin*** *Preferred:* **5 days:** 1000 mg BID *Alternative for risk factors (see comments):* **10 days:** 1000 mg once daily OR **Penicillin G benzathine** 1,200,000 units (1.2 million units) as a **single dose** Low-risk penicillin allergy:[1] **Cephalexin*** 500 mg PO BID for **10 days** High-risk penicillin allergy or low-risk allergy and inability to take cephalexin:[1] **Clindamycin** 300 mg PO TID for **10 days** OR **Azithromycin** 500 mg PO once daily for **5 days**	**Duration of amoxicillin and penicillin** For most adults, we recommend shared decision making with a preference for **5 days** versus **10 days**. **5 days** is recommended for the following reasons: • Symptoms are typically resolved by 5 days regardless of treatment • Longer courses do not affect likelihood of suppurative complications • Risk of acute rheumatic fever (ARF) is very low in the US for most populations (see more details below) • Transmission risk significantly decreases after just one dose of antibiotics • Longer courses are associated with increased adverse drug events and antimicrobial resistance **10 days** (5 days for azithromycin) is recommended for the following scenarios: • If using alternative agents due to allergy (cephalexin or clindamycin) or • If there is concern for current or past ARF or • Concern for poststreptococcal reactive arthritis or glomerulonephritis or • Strong desire for bacterial eradication, such as: ○ local outbreak of ARF, poststreptococcal glomerulonephritis, or severe or invasive GAS disease ○ an outbreak of GAS pharyngitis in a closed or semi-closed community ○ family history of ARF ○ multiple episodes of documented symptomatic GAS pharyngitis occurring within a family for many weeks despite appropriate therapy ○ when tonsillectomy is being considered solely because of frequent GAS isolations • Indigenous or Polynesian ancestry including Hawaii and American Samoa, indigenous peoples of New Zealand and Australia or other Pacific Islands • Recent travel to **areas with high-risk of ARF which include parts of**: ○ **Africa** ○ **Asia** ○ **Pacific** ▪ **Includes Australian and New Zealand indigenous populations** The **United States**, Canada, and most of Europe are considered **low-risk populations.** The incidence in the United States is unknown but is believed to be < 0.5%. **Comments** • GAS is uniformly susceptible to all beta-lactams (penicillins and cephalosporins), whereas macrolide and clindamycin resistance rates of close to 30% have been reported among invasive GAS isolates in the US. These agents should only be used in the case of true beta-lactam allergies. • In case of penicillin allergy, review β-lactam allergy evaluation and empiric guidance and remove allergy label if possible. If not, consider referral to Allergy clinic to provide guidance on future episodes.

* Renal adjustment may be necessary. See Adult Antimicrobial Dosing Guidelines.

[1] Low-risk allergies include: pruritus without rash, remote (> 10 years) unknown reaction, mild to severe rash with no other symptoms; see β-lactam allergy evaluation and empiric guidance for further information.

[2] High-risk allergies include: anaphylaxis, respiratory symptoms (chest tightness, bronchospasm, wheezing, cough), angioedema (swelling, throat tightness), cardiovascular symptoms (hypotension, dizzy/lightheadedness, syncope/passing out, arrhythmia); see β-lactam allergy evaluation and empiric guidance for further information.

[3] Non-IgE mediated allergies are contraindications to further beta-lactam use unless approved by Allergy: organ damage (kidney, liver), drug-induced immune-mediated anemia/thrombocytopenia/leukopenia, rash with mucosal lesions (Stevens Johnson Syndrome/Toxic Epidermal Necrosis), rash with pustules (acute generalized exanthematous pustulosis), rash with eosinophils and organ injury (DRESS – drug rash eosinophilia and systemic symptoms), rash with joint pain, fever, and myalgia (Serum Sickness); see β-lactam allergy evaluation and empiric guidance for further information.

Authors:
Elizabeth Lloyd (Pediatric Infectious Diseases / Antimicrobial Stewardship)
Lindsay Petty (Infectious Diseases / Antimicrobial Stewardship)
Nicholas Dillman (Pharmacy / Antimicrobial Stewardship)
Margeaux Naughton (General Pediatrics)
Aleksandr Belakovskiy (Family Medicine)
Natalie Schellpfeffer (Pediatric Emergency Medicine)

Consultants:
Alison Tribble (Pediatric Infectious Diseases / Antimicrobial Stewardship)
Daniel Riggsbee (Pediatric Infectious Disease / Pharmacy)
Karen Davidge (Pediatric Infectious Disease / Pharmacy)

Last revised: 01/2026	**Antimicrobial Stewardship Subcommittee:** 01/2026	**P&T:** 02/2026

References

	Citation
1	Shulman ST, Bisno AL, Clegg HW, et al. Clinical practice guideline for the diagnosis and management of group A streptococcal pharyngitis: 2012 update by the Infectious Diseases Society of America. Clin Infect Dis. 2012 Nov 15;55(10):e86-102. doi: 10.1093/cid/cis629. Epub 2012 Sep 9.
2	El Feghaly RE, Jaggi P, Katz SE, Poole NM. "Give Me Five": The Case for 5 Days of Antibiotics as the Default Duration for Acute Respiratory Tract Infections. J Pediatric Infect Dis Soc. 2024 Jun 28;13(6):328-333. doi: 10.1093/jpids/piae034.
3	Bradley-Hewitt T, Longenecker CT, Nkomo V, et al. Trends and presentation patterns of acute rheumatic fever hospitalisations in the United States. Cardiol Young. 2019 Nov;29(11):1387-1390. doi: 10.1017/S1047951119002270. Epub 2019 Oct 1.
4	Beaudoin A, Edison L, Introcaso CE, et al. Acute Rheumatic Fever and Rheumatic Heart Disease Among Children — American Samoa, 2011–2012. MMWR Morb Mortal Wkly Rep 2015;64: 555-558.
5	Stockmann C, Ampofo K, Hersh AL, et al. Evolving epidemiologic characteristics of invasive group a streptococcal disease in Utah, 2002-2010. Clin Infect Dis. 2012 Aug;55(4):479-87. doi: 10.1093/cid/cis422. Epub 2012 Apr 24.
6	Stahlgren GK, Tyrstrup M, Edlund C, et al. Penicillin V four times daily for five days versus three times daily for 10 days in patients with pharyngotonsillitis caused by group A streptococci: randomised controlled, open label, non-inferiority study. BMJ. 2019 Oct 4:367:l5337. doi: 10.1136/bmj.l5337.
7	Moore M, Stuart B, Hobbs FdR, et al. Symptom response to antibiotic prescribing strategies in acute sore throat in adults: the DESCARTE prospective cohort study in UK general practice. Br J Gen Pract. 2017 Sep;67(662):e634-e642. doi: 10.3399/bjgp17X692321. Epub 2017 Aug 14.
8	Schwartz RH, Kim D, Martin M, Pichichero ME. A Reappraisal of the Minimum Duration of Antibiotic Treatment Before Approval of Return to School for Children With Streptococcal Pharyngitis. Pediatr Infect Dis J. 2015 Dec;34(12):1302-4. doi: 10.1097/INF.0000000000000883.
9	Dharmapalan D, Bielicki J, Sharland M. Harmonization of Amoxicillin Dose, Duration, and Formulation for Acute Childhood Respiratory Infections. Antibiotics (Basel). 2023 Jun 30;12(7):1138. doi: 10.3390/antibiotics12071138.

Revision History

Date	Revision

The recommendations in this guide are meant to serve as treatment guidelines for use at Michigan Medicine facilities. If you are an individual experiencing a medical emergency, call 911 immediately. These guidelines should not replace a provider's professional medical advice based on clinical judgment or be used in lieu of an Infectious Diseases consultation when necessary. As a result of ongoing research, practice guidelines may from time-to-time change. The authors of these guidelines have made all attempts to ensure the accuracy based on current information, however, due to ongoing research, users of these guidelines are strongly encouraged to confirm the information contained within them through an independent source.