

Question 1

Tries remaining: 2

Marked out of 1.00

$I(q)$ -plots (reciprocal space plots) are usually preferred to $I(r)$ -plots (real space plots) in neutron scattering. Below you will find three different explanations to why that is. Which ones are correct?

Select one or more:

- ☐ a. The magnitude of q is a fundamental physical quantity independent on the instrumental setup (for example, the size of the detector and the wavelength of the neutrons). Therefore, $I(q)$ -plots will be more general than $I(r)$ -plots.
- ☐ b. Traditionally in scattering physics, we prefer notations relating to reciprocal space, as the majority of the data analysis and interpretation is done in this space. As q is a reciprocal space-variable, this is usually the way it is shown.
- ☐ c. q describes actual properties in the sample, which is the most relevant part of the experiment. While r might be interesting for instrumental design, it is irrelevant for the purposes of analyses.

CheckStart againSaveFill in correct responsesSubmit and finishClose preview[Technical information](#)[Download this question in Moodle XML format](#)[Collapse all](#)

Attempt options

How questions behave ?

Interactive with multiple tries

Marked out of

1Start again with these options

Display options

Whether correct

Not shown

Marks

Show max mark only

Decimal places in grades

2

Specific feedback

Shown

General feedback

Shown

Right answer

Not shown

Response history

Not shownUpdate display options

