

DTU Physics

Sources and Monitors

part 2.

2019 CSNS

*McStas
School*

McStas



Sources: Source model overview

➤ *Mathematical:*

- *Source_simple.comp*
- *Source_div.comp*

➤ *Pulsed sources:*

- *ESS_butterfly.comp*
- *ESS_moderator.comp*
- *Moderator.comp*
- *SNS_source.comp (*)*
- *SNS_source_analytic (*)*
- *ViewModISIS (*)*
- *ISIS_moderator.comp (*)*

➤ *Reactors :*


- *Source_Maxwell_3.comp*
- *Source_gen.comp*
- *Source_gen4.comp*
- *Source_multi_surfaces.comp (*)*
- *I/O mechanisms:*
 - ***MCPL_input/output.comp***
 - *Virtual_input/output.comp*
 - *Virtual_mcnp_ss_input/output.comp*
 - *Virtual_tripoli4_input/output.comp*
 - *Vitess_input/output.comp*

Sources: Source_Maxwell_3

```
COMPONENT source = Source_Maxwell_3(yheight=0.156, xwidth=0.126,  
    Lmin=0.1, Lmax=9.0, dist=1.5, focus_xw = 0.025, focus_yh = 0.12,  
    T1=150.42, I1=3.67E11, T2=38.74, I2=3.64E11, T3=14.84, I3=0.95E11)
```

Parameters from the PSI cold source

Initial position and direction: as for Source_simple



Sources: Source_Maxwell_3

```
COMPONENT source = Source_Maxwell_3(yheight=0.156, xwidth=0.126,  
    Lmin=0.1, Lmax=9.0, dist=1.5, focus_xw = 0.025, focus_yh = 0.12,  
    T1=150.42, I1=3.67E11, T2=38.74, I2=3.64E11, T3=14.84, I3=0.95E11)
```

Parameters from the PSI cold source

Intensity at a given wavelength drawn from a sum of (up to) 3 normalized Maxwellian distributions:

$$I(\lambda) = \sum_{i \in 1,2,3} I_i M(\lambda, T_i); \quad M(\lambda, T_i) = 2 \alpha^2 \exp\left(\frac{-\alpha}{\lambda^2}\right) / \lambda^5;$$


$$\alpha = 949.0 \text{ K } \text{\AA}^2 / T_i$$

Sources: Source_Maxwell_3

COMPONENT source

Lmin

T1=1



26,

yh = 0.12,


I3=0.95E11)

from the PSI cold source

Sources: Source_Maxwell_3

```
COMPONENT source = Source_Maxwell_3
  Lmin=0.1, Lmax=9.0, dis
  T1=150.42, I1=3.67E11,
```

Just for fun – let's see what happens if we remove the fast peak...



Input parameters

Parameters in **boldface** are required; the others are optional.

Name	Unit	Description	Default
size	m	Edge of cube shaped source (for backward compatibility)	0
yheight	m	Height of rectangular source	0
xwidth	m	Width of rectangular source	0
Lmin	AA	Lower edge of lambda distribution	
Lmax	AA	Upper edge of lambda distribution	
dist	m	Distance from source to focusing rectangle; at (0,0,dist)	
focus_xw	m	Width of focusing rectangle	
focus_yh	m	Height of focusing rectangle	
T1	K	1st temperature of thermal distribution	
T2	K	2nd temperature of thermal distribution	300
T3	K	3rd temperature of - - -	300
I1	1/(cm**2*st)	flux, 1 (in flux units, see above)	
I2	1/(cm**2*st)	flux, 2 (in flux units, see above)	0
I3	1/(cm**2*st)	flux, 3 - - -	0
target_index	1	relative index of component to focus at, e.g. next is +1 this is used to compute 'dist' automatically.	+1
lambda0	AA	Mean wavelength of neutrons.	0
dlambda	AA	Wavelength spread of neutrons.	0

2019 CSNS

McStas
School

McStas



Sources: Source_gen (Source_gen4)


```
COMPONENT source = Source_gen(yheight=0.156, xwidth=0.126,  
    Lmin=0.1, Lmax=9.0, dist=1.5, focus_xw = 0.025, focus_yh = 0.12,  
    T1=150.42, I1=3.67E11, T2=38.74, I2=3.64E11, T3=14.84, I3=0.95E11)
```

Almost the same as Source_Maxwell_3: but with optional flux-files as input.

2019 CSNS

McStas
School

McStas



MCPL_input/output

Reads/writes events directly from MCPL-format files:
“T. Kittelmann et. al., “”, J. Phys. Comp., 2017


```
COMPONENT vout = MCPL_output(filename="voutput.mcpl",
                                doubleprec=1,polarisationuse=1,verbose=1)
AT (Xout,Yout,Zout) RELATIVE PREVIOUS
```

```
COMPONENT vin = MCPL_input(filename="voutput.mcpl",
                              polarisationuse=1,verbose=1)
AT (Xin,Yin,Zin) RELATIVE PREVIOUS
```

MCPL_input/output

Can include an Implicit Translation:


MCPL_output.comp



MCPL_input/output


Can include an Implicit Translation:

MCPL_output.comp



BECOMES

MCPL_input.comp



Pulsed sources:

Simplest case:

Use a continuous source!

Model a source with given wavelength and spatial distribution

and

... an infinitely short pulse length. I.e. $t = 0$ for all neutron rays.

```
COMPONENT src = Source_simple(
    radius=0.05, lambda0=2.5, dlambda=1.5,
    focus_xw=0.1, focus_yh=0.1, dist=5 )
AT(0,0,0) RELATIVE origin
```

Pulsed sources:

Add a sample here
TOF_monitor.

Simplest case:

Use a continuous source!

Model a source with given wavelength and spatial distribution

and

... an infinitely short pulse length. I.e. $t = 0$ for all neutron rays.

```
COMPONENT src = Source_simple(
    radius=0.05, lambda0=2.5, dlambda=1.5,
    focus_xw=0.1, focus_yh=0.1, dist=5 )
AT(0,0,0) RELATIVE origin
```

Pulsed Sources:

Simplest case:

Use a continuous source!

Model a source with given wavelength and spatial distribution

... an infinitely sharp neutron rays.

Or: Use a chopper
(Wednesday)

```
COMPONENT s1
radius=0.1, dlambda=2.5, dlambda=1.5,
focus_xw=, focus_yh=0.1, dist=5 )
AT(0,0,0) RELATIVE origin
```

Pulsed Sources: Moderator

A flat pulsed source with uniform energy spectrum:


$$x \in U\left[-\frac{xwidth}{2}, \frac{xwidth}{2}\right] \quad y \in U\left[-\frac{yheight}{2}, \frac{yheight}{2}\right]$$

$$|v| = f(\lambda) ; \lambda \in U[L_{min} L_{max}]$$

Time structure is given by energy dependent probability density function:

$$f_t = \frac{1}{\tau} \exp\left(-\frac{t}{\tau}\right)$$

$$\tau = \begin{cases} t_0; & E < E_c \\ t_0 \left(\frac{1}{1 + \frac{(E - E_c)}{\gamma}} \right); & E \geq E_c \end{cases}$$





中

国

散

裂

中

子

源

Pulsed Sources: SNS_source

- Samples directly from tallies coming from e.g. MCNP target+moderator calculations.
- Originally from SNS but also used extensively at J-PARC
- Can be used (with the proper input files) to model CSNS-source.
Example (coming from you) is expected to be included in next release of McStas.


2019 CSNS

McStas
School

McStas



Pulsed Sources: SNS_source



calculations.

McStas.

2019 CSNS

McStas
School

McStas



Pulsed Sources: SNS_source_analytic

- Samples from fits of Padé-functions to tallies from SNS_source.
 - Requires a complex fitting campaign
 - + Much faster than SNS_source
 - + “Cleaner” distributions where statistics are sketchy
- Can be used (with the proper input files) to model CSNS-source.

2019 CSNS

McStas
School

McStas



Pulsed Sources: ViewModISIS


- Samples directly from tallies coming from e.g. MCNP target+moderator calculations.
- Data file supplied for each beam port at ISIS.

Pulsed Sources: ViewModISIS

ISIS T2: IMAT

● S

● D



culations.

Monitors (some)

1D

- ◆ L_monitor $\rightarrow I(\lambda)$
- ◆ TOF_monitor $\rightarrow I(t)$
- ◆ Hdiv_monitor $\rightarrow I(\text{div}_x)$
- ◆ MeanPolLambda $\rightarrow \langle \bar{P} \rangle(\lambda)$
- ◆ E_monitor $\rightarrow I(E)$

2D

- PSD_monitor $\rightarrow I(x, y)$
- PSD_monitor_4PI $\rightarrow I(\theta, \phi)$
- PolLambda_monitor $\rightarrow I(\bar{P}, \lambda)$
- Divergence_monitor $\rightarrow I(\text{div}_x, \text{div}_y)$
- DivPos_monitor $\rightarrow I(\text{div}_x, x)$

nD

- Monitor_nD \rightarrow
 $I(X)$
or
 $I(X, Y)$
or
 $Z(X, Y, Z)$
or ...

2019 CSNS

McStas
School

McStas




Monitors: Quick examples

```
COMPONENT my_L_monitor = L_monitor(xwidth=0.2, yheight=0.2,  
                                     nL=20, filename="Output.L", Lmin=2, Lmax=10)
```

In a histogram sense


Imagine a histogram, e.g. $\mathbf{I}(\lambda)$





In bin i , \mathbf{N} events each carrying a fractional intensity p_j so that

$$I = \sum_N p_j$$

The RMS variance over that set becomes our statistical error bar \mathbf{E}







From "Virtual experiments - the ultimate aim of neutron ray-tracing simulations", K. Lefmann et al., Journal of Neutron Research 16, 97-111 (2008)

Let n be the number of neutron rays reaching the detector, and let the rays have (different) weights, w_i . The simulated intensity is then given by

$$I = \sum_{i=1}^n w_i. \quad (1)$$

The estimate of the error on this number is calculated in the McStas manual [1], and the standard deviation is approximated by

$$\sigma^2(I) = \sum_{i=1}^n w_i^2. \quad (2)$$

In real experiments, $w_i = 1$, whence we reach $I = n$ and $\sigma(I) = \sqrt{I}$ as expected (for counts exceeding 10). Let the virtual time be denoted by t . The simulated counts during this time becomes

$$C = tI, \quad (3)$$

From "Virtual experiments - the ultimate aim of neutron ray-tracing simulations", K. Lefmann et al., Journal of Neutron Research 16, 97-111 (2008)

and its error bar estimate is

$$\sigma^2(C) = t^2 \sigma^2(I). \quad (4)$$

However, to simulate a realistic counting statistics, we must fulfill

$$\sigma_{VE}(C_{VE}) = \sqrt{C_{VE}}. \quad (5)$$

This is obtained by adding to (3) a Gaussian noise $E(\Sigma)$ of mean value zero and standard deviation Σ :

$$C_{VE} = tI + E(\Sigma). \quad (6)$$

The standard deviation for the VE becomes

$$\sigma_{VE}^2(C) = t^2 \sigma^2(I) + \Sigma^2. \quad (7)$$

Now, the requirement (5) allows us to determine Σ :

$$\Sigma^2 = tI - t^2 \sigma^2(I). \quad (8)$$

Since Σ^2 must remain positive, we reach an upper limit on t

$$t_{\max} = \frac{I}{\sigma^2(I)}. \quad (9)$$

Sketch of an algorithm...

1. On a given McStas histogram

2. For the non-zero bins. calculate

$$t_{\max} = \frac{I}{\sigma^2(I)}.$$

t_{\max}

The *smallest* defines the “maximal counting time” allowed by your statistics

3. Preferably a “background” should be added - use a “known experimental value” or an estimate...



Monitor_nD

The all-in-one , swiss-army-knife of monitors

***Monitor_nD** can have almost any shape, and record any requested standard quantities*



2019 CSNS

McStas
School

McStas



Monitor_nD

A general monitor for 0D/1D/2D records

Examples

```
COMPONENT MyMon = Monitor_nD( xwidth = 0.1, yheight = 0.1, zdepth = 0,  
    options = "intensity per cm2 angle,limits=[-5 5],  
    bins=10,with borders, file = mon1")
```

options = "banana, theta limits=[10,130], bins=120, y"

options = "multiple kx ky kz, auto abs log t, and list all neutrons"

Monitor_nD


... or monitor just about anything:

```
COMPONENT MyMon = Monitor_nD(xwidth = 0.1, yheight = 0.1,  
    user1=age, username1="Age of the Captain [years]",  
    options="user1, auto")
```

2019 CSNS

McStas
School

McStas



Exercise 2:

Head over to the github site and continue the exercise we started before:

https://github.com/McStasMcXtrace/Schools/tree/master/CSNS_March_2019/1_Monday_March_25th/2_Sources_and_Monitors

2019 CSNS

**McStas
School**

McStas






EUROPEAN
SPALLATION
SOURCE

McStas



DTU



中国
China

中国散裂中子源

China Spallation Neutron Source

**2019 CSNS
McStas School**