# **NEOS**

Air & water carcinogens

France

**Epidemiology** 

Care

## **NEoplasm Open-Source Environmental Risk Factors Standardization**

Pascal Deschaseaux<sup>1</sup>, MD, MBA; Sébastien de Longeaux<sup>1</sup>, MBA; Edouard Debonneuil<sup>2</sup>, PhD; Rachel Aronoff<sup>3</sup>, PhD

**Environmental** 

Behavioral &

occupational

CHALLENGE: accelerating research on cancer risk factors by structuring open data sources, and completing the OSIRIS clinical and -omics databases, in order to standardize variables related to the environment (terminology, interoperability,..)

**OBJECTIVE**: easing analyses of environmental cancer risk factor data by structuring and harmonizing open source epidemiological data sets, in a FAIR approach

## **WORK PACKAGES:**

WP1: ontology

WP2: data & metadata

WP3: data sources

## **TARGET OUTCOME:**

Standardized cancer epidemiology dataset framework & examples

**RISK FACTOR SELECTION:** 

Cancer

Other diseases

SCOPE:

Easily measurable  $\rightarrow$  air & water agents, France

Prognostic factors

**Risk factors** 

- International reference → IARC (International Agency for Research on Cancer) monographs
- Scientifically proven → Group 1 carcinogens (substances known to have carcinogenic potential for humans)

Alliance 3 AGiR! & Hackuarium



**IARC Group 1** 

IARC groups

2A. 2B. 3

Air & water

Other agents

## **EXAMPLES:**

Private sources

**Open sources** 

From a list of 37 IARC Group I air & water biological, chemical and physical agents with open source data, we selected two carcinogens:

Rest of world

- An air pollutant: PM 2.5 (fine particle matter), associated with lung cancer risk
- A water pollutant: arsenic, associated with lung, urinary bladder and skin cancer risk

## **NEOS FRAMEWORK (SELECTED FIELDS):**

Item group ● Objectives ● Item N° ● Collection status ● Item ● Item definition ● Expected value

Geographic location of measure ● Geographic granularity of measure ● Date of measure ● Temporal granularity of measure ● Data source ● Geographic and temporal relevance

Main cancer sites associated with agent ● Reference value ● Guidelines ● Monograph/backup paper ● Main sources of exposure

Current address ● For how long ● Past addresses (starting with most recent, as detailed as possible) ● For how long (years) for each past address ● Main occupation ● Usual place of main occupation ● For how long (years) ● Main mode of transportation ● How many days a month ● How many hours a week

Exposure to carcinogen (concentration in medium)



atmospheric pollution » scenario (source : InVS, Santé Publique France). Map shows data measurement coverage

## **CONCLUSIONS:**

- Open source environmental data are very heterogeneous
- Two types of data are crucial for the NEOS Framework: place of residence/occupation, total duration of exposure.
- Definition of variables must be in context and precise to avoid bias
- Data collection and analyses at the patient level require a precise address and geocoding.
- This work will be expanded to other IARC Group I environmental cancer risk factors with open sources, using the NEOS Framework

Possible limitations, particularly for rural areas, include the place where measurements are obtained, and agents' geographical coverage.