CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International Advanced Subsidiary and Advanced Level

MARK SCHEME for the May/June 2015 series

9700 BIOLOGY

9700/34

Paper 3 (Advanced Practical Skills 2), maximum raw mark 40

www.PapaCambridge.com

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme Sy. per
	Cambridge International AS/A Level – May/June 2015 970 970
Mark scher	me abbreviations:
;	separates marking points
1	alternative answers for the same point
R	reject
Α	accept (for answers correctly cued by the question, or by extra guidance)
AW	alternative wording (where responses vary more than usual)
<u>underline</u>	actual word given must be used by candidate (grammatical variants accepted)

Mark scheme abbreviations:

max indicates the maximum number of marks that can be given

or reverse argument ora

marking point (with relevant number) mp

error carried forward ecf

ignore

Page	3	Mark Scheme Syl oer
		Cambridge International AS/A Level – May/June 2015 970 970
1 (a) le	vel of risk) medium or high ;
(b) (i)	(labels under correct sequence of beakers) 0.03 + 0.003 + 0.0003 + %;
		shows transfer of 1 cm ³ of solution from previous beaker to 2 beakers;

- (a) level of risk) medium or high;
 - (b) (i) (labels under correct sequence of beakers) 0.03 + 0.003 + 0.0003 + %;

adds 9 cm³water/W to three beakers;

[3]

- (ii) 1 table with heading + percentage concentration of X;
 - 2 table with heading + number or no. of bubbles;
 - 3 records results for W or 0% and 4 concentrations;
 - 4 records lowest concentration of **X** with a higher number of bubbles than highest concentration of X;
 - repeats at least one concentration;

[5]

(iii) whole seconds recorded and shows 2 divided by this value;

correct answer calculated to correct number of significant figures;

[2]

(iv) idea of inhibits activity;

idea of preventing substrate binding to the enzyme/active site or fewer enzymesubstrate complexes formed;

[2]

(v) (counting bubbles) different sizes or too fast or bubbles group together;

(displacement of water) gas escapes from delivery tube or not all bubbles go into syringe or parallax error;

[2]

(vi) (independent variable) use the same concentration of X;

5 or more temperatures;

use thermostatically-controlled water-bath;

[3]

[Total: 18]

_		•		W 101	
Р	age	4		Mark Scheme Sy. Cambridge International AS/A Level – May/June 2015 970	er
2	(a)	(i)	orie	Mark Scheme Cambridge International AS/A Level – May/June 2015 entation axis) length of neck/cm + (y-axis) thickness of muscle wall in left ventricle/mm ale axis) 2 cm to 10 labelled each 2 cm + must have 50 at the origin + (y-axis) 2 cm to	Brick
			sca (x-a	ale axis) 2 cm to 10 labelled each 2 cm + must have 50 at the origin + (y-axis) 2cm to 5 labelled each 2 cm, + must have 20 at origin;	Se.
			plo	tting correct plotting of 5 points;	
			line 5 p	lots with ruled lines exactly point to point or line of best fit + quality smooth line les than 1 mm thick ;	ss [4]
		(ii)	cor	rect estimate from candidate graph;	[1]
		(iii)	ide	a of thicker/stronger/wall or muscle to push blood up longer neck or to push the blood further;	[1]
	(b)	(i)	1	correct selection of vessel Q or T ;	
			2	size at least 100 mm + no shading;	
			3	length of drawing is at least twice the size of the narrowest width;	
			4	draws at least three lines across wall + inner line crinkled;	
			5	proportions of vessel walls correct with one selected;	[5]
		(ii)	1	shows on Fig. 2.1 where measured S ;	
			2	shows at least 5 of measurements of the diameter + 5 measurements of the thickness of the wall ;	
			3	measures at least 3 for each in whole mm or to $\pm 0.5\text{mm}$;	
			4	answer shown as larger number to smaller number to lowest common denomina	tor ; [4]
	(c)	(i)	1	sharp continuous lines + size at least 40 mm for at least one cell;	
			2	draws only 4 xylem vessels + at least 2 touching;	
			3	for at least 2 cells, walls drawn as double lines, with middle lamella;	
			4	straight line where 2 cells meet or at least one cell with at least one angle present	nt ;
			5	correct label with label line ending in the lumen;	[5]
		(ii)	ide	a that Fig. 2.1 has thicker walls than the xylem or Fig. 2.1 has more than one laye xylem has only one ;	er [1]
		, <u>.</u>	1		F41

(iii) lumen + space / no (cell) contents or lumen + idea of less resistance;

[1]

[Total: 22]

2