

Week 6 Notes: From One Measurement to a Phase Diagram

Seminar notes

May 21, 2026

1 Purpose of Week 6

Week 5 established a proof of principle: after mapping the Kitaev chain to qubits, a gate-prepared state can reveal topology through a non-local Pauli string. The Week 5 example was deliberately narrow. It focused on the topological sweet spot

$$t = \Delta, \quad \mu = 0, \quad (1)$$

and measured the boundary string

$$X_0 Z_1 Z_2 X_3 \quad (2)$$

for a four-site chain.

Week 6 turns this single-point demonstration into a phase diagnostic. The parameter μ is swept through trivial, critical, and topological regimes. The central question is:

Can the same circuit-measurable non-local string reconstruct the known phase boundary $|\mu| = 2t$?

This is still part of Block 3, not yet the full Block 4 noise study. The role of Week 6 is to define the ideal observable curve that later noisy-circuit simulations should be compared against.

2 Qubit Hamiltonian Used in the Sweep

The Jordan-Wigner mapping gives the open-chain qubit Hamiltonian

$$H(\mu) = -\frac{\mu}{2} \sum_{j=0}^{L-1} Z_j + \frac{t - \Delta}{2} \sum_{j=0}^{L-2} X_j X_{j+1} + \frac{t + \Delta}{2} \sum_{j=0}^{L-2} Y_j Y_{j+1}. \quad (3)$$

In the Week 6 script we keep

$$t = \Delta = 1, \quad (4)$$

so the XX term vanishes and the Hamiltonian becomes

$$H(\mu) = -\frac{\mu}{2} \sum_j Z_j + \sum_{j=0}^{L-2} Y_j Y_{j+1}. \quad (5)$$

The sweep changes only the transverse-field strength μ . The known thermodynamic transition points are

$$\mu = \pm 2t. \quad (6)$$

For finite L , the transition is rounded, but the same scale should still organize the numerical data.

3 Why a Non-Local String Is Needed

A single boundary Majorana operator changes fermion parity. Since the Hamiltonian conserves fermion parity, an energy eigenstate can be chosen with definite parity. For any operator A that anticommutes with parity,

$$\{A, P\} = 0, \quad (7)$$

its expectation value in a parity eigenstate vanishes:

$$\langle A \rangle = \langle \psi | A | \psi \rangle \quad (8)$$

$$= \langle \psi | P A P | \psi \rangle \quad (9)$$

$$= -\langle \psi | A | \psi \rangle = 0. \quad (10)$$

This is why local edge measurements alone are not a stable diagnostic of the topological phase.

The parity-preserving observable used in Week 6 is

$$O_{\text{edge}} = X_0 Z_1 \cdots Z_{L-2} X_{L-1}. \quad (11)$$

For $L = 4$ this reduces to

$$O_{\text{edge}} = X_0 Z_1 Z_2 X_3, \quad (12)$$

which is the same Pauli string emphasized in the Week 5 deck, up to an overall sign convention inherited from the Majorana operator ordering. The sign is not important for the phase diagnostic, so the plotted quantity is the magnitude

$$|\langle O_{\text{edge}} \rangle|. \quad (13)$$

4 Measurement Protocol

Quantum hardware measures in the computational, or Z , basis. A Pauli string expectation value is obtained by rotating each non- Z factor into the Z basis, measuring all qubits, and multiplying the resulting eigenvalues.

For

$$O_{\text{edge}} = X_0 Z_1 \cdots Z_{L-2} X_{L-1}, \quad (14)$$

the measurement circuit is:

1. Prepare the ground-state approximation with the VQE ansatz.
2. Apply a Hadamard gate to qubit 0 and qubit $L - 1$, because

$$H X H = Z. \quad (15)$$

3. Measure all qubits in the computational basis.
4. Convert every measured bit b_j into an eigenvalue $z_j = (-1)^{b_j}$.
5. Multiply the relevant eigenvalues:

$$m_s = z_0 z_1 \cdots z_{L-2} z_{L-1}. \quad (16)$$

Each shot returns $m_s = \pm 1$. The shot estimator is

$$\widehat{\langle O \rangle} = \frac{1}{N} \sum_{s=1}^N m_s, \quad (17)$$

with standard error

$$\sigma_{\hat{O}} \approx \sqrt{\frac{1 - \langle O \rangle^2}{N}}. \quad (18)$$

The Week 6 script uses a binomial sampling model with $N = 4096$ shots to emulate this final measurement layer.

5 Validation Path Used in the Current Script

The current Week 6 implementation is in `src/block3_week6.py`. It is intentionally a validation script rather than the final hardware-style VQE sweep. For each μ value it:

1. builds the qubit Hamiltonian $H(\mu)$ using `kitaev_qubit_hamiltonian`;
2. diagonalizes the small $2^L \times 2^L$ matrix;
3. selects the lowest even-parity eigenstate;
4. evaluates $\langle O_{\text{edge}} \rangle$ exactly;
5. samples a synthetic finite-shot estimator for the same Pauli string;
6. computes the parity gap $|E_0^{(+)} - E_0^{(-)}|$ as a spectral cross-check.

This use of exact diagonalization is not meant to replace VQE. Its purpose is to produce a trusted small- L target curve and verify the observable convention before spending time on a VQE loop at every μ point.

6 Reading the Week 6 Figure

The shaded region marks the known topological interval

$$|\mu| < 2t. \quad (19)$$

The expected qualitative behavior is:

- inside the topological region, the two-edge string remains large because the boundary Majoranas are correlated through the non-local fermion;
- outside the topological region, the string drops because the chain becomes trivial and long-range edge correlation is lost;
- near $|\mu| = 2t$, finite-size rounding replaces the sharp thermodynamic transition;
- the parity gap is small in the topological region and grows in the trivial region.

The left panel therefore gives the measurement-side signature, while the right panel gives a spectrum-side check from the qubit Hamiltonian. The point of the figure is not that exact diagonalization is the final algorithm. The point is that the chosen Pauli string carries the same phase information as the more traditional parity-gap diagnostic.




Figure 1: Week 6 phase sweep for $L = 4$, $t = \Delta = 1$. The left panel shows the non-local edge-string correlator and a finite-shot estimator. The right panel shows the parity-gap cross-check.

7 Connection to VQE

The final Block 3 version should replace the exact even-parity eigenstate in the sweep with a VQE-prepared state

$$|\psi(\theta^*(\mu))\rangle = U(\theta^*(\mu))|0 \cdots 0\rangle, \quad (20)$$

where

$$\theta^*(\mu) = \arg \min_{\theta} C_{\mu}(\theta). \quad (21)$$

A parity-biased cost function can be reused from the Week 5 notebook:

$$C_{\mu}(\theta) = \langle H(\mu) \rangle_{\theta} - \lambda \langle P \rangle_{\theta}, \quad P = \prod_{j=0}^{L-1} Z_j. \quad (22)$$

After optimization, the same Pauli-string measurement protocol estimates

$$\langle \psi(\theta^*) | O_{\text{edge}} | \psi(\theta^*) \rangle. \quad (23)$$

The small- L exact curve from Week 6 should be kept as a benchmark. It can answer whether VQE error, shot noise, or sign/string-convention mistakes are responsible when the circuit curve deviates from the expected phase diagram.

8 Bridge to Block 4

Block 4 will ask whether this diagnostic survives realistic near-term hardware effects. The Week 6 curve defines the ideal reference. The natural Block 4 extensions are:

- add depolarizing noise after one- and two-qubit gates;
- add readout error before classical bitstring processing;
- optionally add thermal relaxation through T_1 and T_2 parameters;
- compare ideal, noisy, and mitigated estimates of $|\langle O_{\text{edge}} \rangle|$;

- quantify how much noise can be tolerated before the phase boundary is no longer visible.

In short, Week 6 finishes the ideal measurement story: a non-local Pauli string can be swept across μ and used as a circuit-compatible phase diagnostic. Block 4 will test the robustness of that diagnostic under noisy measurement and imperfect gates.