

Majorana Modes in the Machine

Week 5: Transitioning to Quantum Circuit Measurement

Dobromir Stoev
Edgar Harutyunyan
Guilherme Schewtschik
Jaskaran Singh
Zhengyi Liu
Zhenming Shi

Advanced Seminar — AI-Augmented Theoretical Physics

May 17, 2026

Week 5 Objectives: Moving to Block 3


- **Main Goal:** Shift from "Exact Matrix Math" to "Gate-Based Simulation."
- **Task 1:** Ground State Preparation via VQE (Variational Quantum Eigensolver).
- **Task 2:** Defining the Non-local String Order Parameter $\mathcal{O}_{string}$.
- **Task 3:** Mapping Majorana Observables to Qubit Measurement Gates.
- **Task 4:** Numerical Evidence.

Task 1: VQE Circuit Ansatz (Visual)

Hardware-Efficient Ansatz:

- Uses only R_y to keep the state in the real domain.
- Linear CNOT entanglement mimics 1D hopping.

Hardware-Efficient Ansatz Structure (L=4)



R_y Gate

For $L = 4$, the configuration space consists of $2^4 = 16$ possible states:

$$|\Psi\rangle = c_0|0000\rangle + c_1|0001\rangle + \cdots + c_{15}|1111\rangle$$

Initially, $|\psi\rangle = [1, 0, 0, \dots, 0]^T$. An $R_y(\theta)$ gate on q_0 performs:

$$R_y(\theta)|0\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)|1\rangle$$

In the 16D vector, the value "1" splits:

- Index 0 ($|0000\rangle$): becomes $\cos(\theta/2)$
- Index 8 ($|1000\rangle$): becomes $\sin(\theta/2)$

CNOT Gate

Entanglement occurs when a gate moves amplitudes between binary configurations.

CNOT(0,1) Operation

If q_0 is 1, flip q_1 . In the vector:

- 1 $|1000\rangle$ (Index 9) satisfies the condition.
- 2 It is transformed into $|1100\rangle$ (Index 13).

Physical meaning: isolated excitations are combined into **paired states**, building the topological correlations needed for Majoranas.

Hilbert Space Evolution: Mathematical Verification

Mathematical Constraint: Post-entanglement states cannot be factored into independent single-qubit probabilities. The evolution must be tracked across the full $2^4 = 16$ dimensional state vector.

State Vector Evolution Table (Tracking non-zero amplitudes):

Index (Dec)	Basis ($q_0q_1q_2q_3$)	Init State $ \psi_0\rangle$	Post $R_y(\theta_0)$ $ \psi_1\rangle$	Post CNOT _{0→1} $ \psi_2\rangle$
0	$ 0000\rangle$	1	$\cos(\frac{\theta_0}{2})$	$\cos(\frac{\theta_0}{2})$
1	$ 0001\rangle$	0	0	0
⋮	⋮	⋮	⋮	⋮
8	$ 1000\rangle$	0	$\sin(\frac{\theta_0}{2})$	0 (<i>Amplitude shifted</i>)
⋮	⋮	⋮	⋮	⋮
12	$ 1100\rangle$	0	0	$\sin(\frac{\theta_0}{2})$

Hamiltonian Simplification: The Pure YY Limit

General Qubit Mapping (Block 2):


$$H_K = -\frac{\mu}{2} \sum Z_j + \frac{t-\Delta}{2} \sum X_j X_{j+1} + \frac{t+\Delta}{2} \sum Y_j Y_{j+1}$$

Topological Sweet Spot ($t = \Delta$, $\mu = 0$):

$$H = \sum_{j=0}^{L-2} Y_j Y_{j+1}$$

- **XX Cancellation:** Setting $t = \Delta$ (hopping = pairing) mathematically eliminates all $X_j X_{j+1}$ terms.
- **Field Removal:** Setting $\mu = 0$ removes the transverse Z fields, reaching the deepest topological phase.

Numerical Verification: VQE Convergence Dynamics



Optimization Parameters:

- **Algorithm:** L-BFGS-B (Gradient-based).
- **Cost Function:**
$$C(\vec{\theta}) = \langle H \rangle - \lambda \langle P \rangle.$$
- **Penalty ($\lambda = 0.1$):** Enforces strictly even parity ($P = +1$) selection, breaking topological degeneracy.

Task 2: String Operator

Boundary Majorana Correlator to Qubit Mapping:

$$\hat{O}_{corr} \propto i\gamma_{0,2}\gamma_{3,1} = i(Y_0)(Z_0Z_1Z_2X_3) = -X_0Z_1Z_2X_3$$


- **1. Breakdown of the Landau Paradigm:** The topological phase hosts no local symmetry breaking. All local single-qubit expectations strictly vanish:

$$\forall j \in \{0, 1, 2, 3\}, \quad \langle X_j \rangle = 0, \quad \langle Y_j \rangle = 0, \quad \langle Z_j \rangle = 0$$

Non-local order parameters are mandatory to detect this hidden order.

- **2. Ground State Commutation Constraint:** A saturated non-zero invariant requires $[H, \hat{O}_{string}] = 0$. The naive boundary operator $\hat{O}_{naive} = X_0X_3$ anti-commutes with H , creating high-energy quasiparticle excitations and yielding $\langle X_0X_3 \rangle \equiv 0$.


Task 3: Measuring Topology via Basis Rotation



Hardware measures in Z -basis. Majoranas live in X -basis.

- **Observable** ($X_0 Z_1 Z_2 X_3$)
- **Hadamard (H) Gate:** Maps $X \rightarrow Z$.
- Applied to q_0 and q_3 to extract the non-local correlation.

Task 4: Numerical Evidence: The 1.0 Correlation



- **Left (Y_0):** ≈ 0 due to anti-commutation with the global parity operator ($\{Y_0, \hat{P}\} = 0$).
- **String ($X_0 Z_1 Z_2 X_3$):** Exactly 1.0000 (Theoretical limit).
- **Algebraic Symmetry:** The commutation $[H, \hat{O}_{corr}] = 0$ identifies this as a conserved topological invariant.
- **Result:** Non-local Topology confirmed in the gate-based model.

Summary & Next Steps

- **Completed:** VQE ground state preparation.
- **Discovery:** The non-local string order parameter is a robust topological signature.
- **Next:** Sweep μ to generate the full Phase Diagram using only quantum circuit results.