

# Majorana Modes in the Machine

Blocks 1 & 2: The Physics Bridge & Finite-Size Physics

Dobromir Stoev  
Edgar Harutyunyan  
Guilherme Schewtschik  
Jaskaran Singh  
Zhengyi Liu  
Zhenming Shi

Advanced Seminar — AI-Augmented Theoretical Physics

July 6, 2026

# Outline


- 1 The Kitaev Hamiltonian & Edge Modes
- 2 Bulk Spectrum & Phase Transition
- 3 Topological Invariant
- 4 Topology & Protection
- 5 Block 1 Summary
- 6 Numerical Method
- 7 Finite-Size Results
- 8 Edge Mode Localization
- 9 Majorana Hybridization Splitting
- 10 Block 2 Summary

# Starting point: the lattice Kitaev Hamiltonian

$$H_K = -\mu \sum_j c_j^\dagger c_j - t \sum_j (c_j^\dagger c_{j+1} + \text{h.c.}) + \Delta \sum_j (c_j^\dagger c_{j+1}^\dagger + \text{h.c.}) \quad (1)$$

## Physical ingredients

- Spinless (or spin-polarized) fermions
- Nearest-neighbor hopping  $t$
- $p$ -wave pairing  $\Delta \in \mathbb{R}$
- Chemical potential  $\mu$  tunes filling



## Questions we will answer

- When does topology emerge?
- How do edge zero modes appear?

# Majorana Representation & The “Sweet Spot”

Ordinary fermions can be split into two Hermitian **Majorana operators**:

$$\gamma_{A,j} = \frac{c_j + c_j^\dagger}{\sqrt{2}}, \quad \gamma_{B,j} = \frac{i(c_j^\dagger - c_j)}{\sqrt{2}} \quad (2)$$

By rewriting  $H_K$  in terms of Majoranas (see Appendix for details), we find a special limit:

**Sweet spot:  $\mu = 0, t = \Delta$**

The intra-site coupling vanishes. Only inter-site coupling  $i\gamma_{B,j}\gamma_{A,j+1}$  survives. In a chain with open boundaries, this leaves **one unpaired Majorana zero-mode at each end**.

**Why is this interesting?**

These edge Majorana modes are **robust** (protected by topology) and candidates for **fault-tolerant quantum computation**.

# Bulk dispersion: topological / critical / trivial

Applying a Fourier transform and using the BdG formalism (see Appendix), the energy spectrum is given by:

$$E_k = \pm \sqrt{n_z(k)^2 + n_y(k)^2} = \pm \sqrt{(-\mu - 2t \cos k)^2 + (-2\Delta \sin k)^2} \quad (3)$$

The topological phase transition occurs when the **bulk gap closes** ( $E_k = 0$ ). This happens at  $k = \{0, \pi\}$  when  $\mu = \pm 2t$ .

Bulk Dispersion of the 1D Kitaev Chain ( $t = 1.0, \Delta = 1.0$ )



# Winding number $\nu$

As  $k$  sweeps the BZ, the vector  $\mathbf{n}(k) = (n_z, n_y)$  traces a closed loop.

$$\theta_k = \arg(n_z + in_y), \quad \nu = \frac{1}{2\pi} \int_{\text{BZ}} \partial_k \theta_k dk \quad (4)$$

Winding Loops of the 1D Kitaev Chain in Complex Plane ( $t = 1.0, \Delta = 1.0$ )



- $\nu = 0$ : loop does *not* enclose origin  $\rightarrow$  trivial
- $|\nu| = 1$ : loop winds once around origin  $\rightarrow$  topological

# Phase diagram



Bulk gap vanishes at  $\mu = \pm 2t$ ; winding number jumps across the transition.

# Topology: Continuity and Invariance

- **Topological Equivalence:** Two shapes are equivalent if they can be transformed into each other via “continuous deformation” (stretching, bending).
- **Invariants:** Properties that remain unchanged during deformation (e.g., number of holes / Genus).
- **Physical Link:** Tuning Hamiltonian parameters  $(t, \mu)$  is essentially a “continuous deformation” of the wave function.

## Core Intuition


You cannot turn a **Sphere** (0 holes) into a **Donut** (1 hole) without “tearing” the surface (Gap Closing).



# Topological invariant: The Winding Number



with Genus 1



winding number  $\nu = 1$

## What is Protection?

As long as the circle doesn't cross the origin (Gap remains open), parameter fluctuations only "deform" the circle but cannot change the **integer** winding number.

# Block 1 — key takeaways

- ① The Kitaev Hamiltonian hosts **two phases** separated by gap closings at  $\mu = \pm 2t$ .
- ② Majorana representation reveals the **sweet-spot** picture directly ( $\mu = 0, t = \Delta$ ).
- ③ The BdG d-vector winding number  $\nu \in \{0, 1\}$  is the **topological invariant**.

**Now (Block 2):** Finite size & edge modes — from bulk invariant to real-space edge states.

## Step 1: Building the BdG matrix

The real-space BdG Hamiltonian is a  $(2L \times 2L)$  block matrix:

$$M = \begin{pmatrix} h & d \\ -d & -h \end{pmatrix}, \quad h_{ij} = -\mu \delta_{ij} - t(\delta_{i,j+1} + \delta_{i+1,j}), \quad d_{ij} = \Delta(\delta_{i,j+1} - \delta_{i+1,j})$$

```
def build_hamiltonian(self):
    h = np.zeros((L, L))
    np.fill_diagonal(h, -mu)           # onsite: chemical potential
    for j in range(L - 1):
        h[j, j+1] = h[j+1, j] = -t    # hopping

    d = np.zeros((L, L))
    for j in range(L - 1):
        d[j, j+1] = delta              # pairing (antisymmetric)
        d[j+1, j] = -delta

    return np.block([[h, d], [-d, -h]])
```


## Step 2: Diagonalization & extracting the spectrum

```
def positive_spectrum(self):  
    evals = np.linalg.eigvalsh(self.build_hamiltonian())  
    return np.sort(np.abs(evals))[:,2]    # |E| plus one copy per +/-E  
                                           pair
```

```
# Majorana splitting scan over chain lengths:  
EO_arr = np.array([  
    np.sort(np.abs(KitaevChain(L=1, ...).spectrum()))[0]  
    for l in L_arr  
)
```

Particle-hole symmetry gives eigenvalues in  $\pm E$  pairs — we take  $|E|$  and keep the  $L$  smallest values to get the quasiparticle spectrum. (*Why not just filter  $evals \geq 0$ ? See Appendix D.*)


# Finite-size spectrum (OBC): edge modes appear



- Pink curve: near-zero mode pinned inside the gap across the entire topological phase  $|\mu| < 2t$ .
- Gray curves: bulk quasiparticle bands — gapped in both phases, closing only at  $\mu = \pm 2t$ .

# Real-space BdG snapshot

Full BdG spectrum in real space ( $L = 30$ , OBC)



**Left** (trivial,  $\mu = -3t$ ): all eigenvalues uniformly gapped — no special states.

**Right** (topological,  $\mu = -t$ ): two eigenvalues pinned at  $E \approx 0$  — the Majorana pair localized at opposite ends of the chain.

# Exponential localization of the edge mode

At the symmetric point  $t = \Delta$ , the zero-mode wavefunction has an exact analytical form. Substituting the Ansatz  $\psi_j \sim \lambda^j$  into the bulk BdG equations gives:

$$\lambda = \frac{\mu}{2t}, \quad |\lambda| < 1 \text{ (topological phase)}$$

The left edge mode decays as:

$$\psi_j^L \propto \left(\frac{\mu}{2t}\right)^{j-1} = e^{-(j-1)/\xi}$$

with coherence length

$$\xi = \frac{1}{\ln(2t/|\mu|)}$$

## Key points

- Pure exponential at  $t = \Delta$  (no oscillations)
- $\xi \rightarrow \infty$  as  $|\mu| \rightarrow 2t$  (phase boundary)
- $\xi \rightarrow 0$  as  $|\mu| \rightarrow 0$  (sweet spot)

## Derivation: finite-size splitting $E_0(L)$

In a **finite** chain, the left and right edge modes overlap, lifting the degeneracy. First-order perturbation theory gives:

$$\delta E = \frac{|\langle \psi^L | H | \psi^R \rangle|}{\|\psi^L\| \|\psi^R\|}$$

The overlap is dominated by the mid-chain region:


$$\langle \psi^L | \psi^R \rangle \sim \sum_{j=1}^L \lambda^{j-1} \cdot \lambda^{L-j} = L \lambda^{L-1}$$

The exact transfer-matrix calculation ( $t = \Delta$ ) keeps the  $(1 - \lambda^2)$  prefactor (with  $\lambda = |\mu|/2t$ ):

$$E_0(L) \approx 2t (1 - \lambda^2) \left( \frac{|\mu|}{2t} \right)^L$$



# Majorana hybridization splitting vs system size



**Key result** — at the symmetric point  $t = \Delta$ :

$$E_0(L) \approx 2t \left( \frac{|\mu|}{2t} \right)^L$$

- Straight lines on the semilog scale confirm **exponential** decay.
- Slope  $= \ln(|\mu|/2t) = -1/\xi$  measures the **coherence length**  $\xi$ .
- Smaller  $|\mu| \Rightarrow$  faster decay  $\Rightarrow$  better protected zero mode.

## Block 2 — key takeaways

- 1 The BdG matrix is built directly from  $\mu$ ,  $t$ ,  $\Delta$  — exact diagonalization gives the full spectrum with no approximations.
- 2 OBC spectra show **near-zero edge modes** in the topological phase — direct numerical confirmation of bulk-boundary correspondence.
- 3 Edge modes are **exponentially localized** with coherence length  $\xi = 1/\ln(2t/|\mu|)$ .
- 4 Majorana splitting decays as  $E_0 \sim (|\mu|/2t)^L$  — exponential in  $L$ , measurable from the semilog slope.

**Next session (Block 2 cont.):** Jordan–Wigner transform  $\rightarrow$  qubit encoding  $\rightarrow$  quantum circuit for ground-state preparation.

# Appendix

# Appendix A: Full Majorana Representation

Recall the Majorana operators:

$$\gamma_{A,j} = \frac{c_j + c_j^\dagger}{\sqrt{2}}, \quad \gamma_{B,j} = \frac{i(c_j^\dagger - c_j)}{\sqrt{2}} \quad (5)$$

where  $\{\gamma_{\alpha,i}, \gamma_{\beta,j}\} = 2\delta_{\alpha\beta}\delta_{ij}$ .

Rewriting the full Kitaev Hamiltonian in the Majorana basis yields:

$$H_K = -\mu \sum_j \left( i\gamma_{A,j}\gamma_{B,j} + \frac{1}{2} \right) + i \sum_j \left[ (\Delta + t) \gamma_{B,j}\gamma_{A,j+1} + (\Delta - t) \gamma_{A,j}\gamma_{B,j+1} \right] \quad (6)$$

*Note how setting  $\mu = 0$  and  $t = \Delta$  simplifies this to just the  $i\gamma_{B,j}\gamma_{A,j+1}$  term.*

## Appendix B: Momentum Space Mapping

The tight-binding Hamiltonian in real space:

$$H = \sum_j \left[ -t(c_j^\dagger c_{j+1} + \text{h.c.}) - \mu \left( c_j^\dagger c_j - \frac{1}{2} \right) + \Delta(c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger) \right] \quad (7)$$

Applying the Fourier transform  $c_j = \frac{1}{\sqrt{N}} \sum_k e^{ikj} c_k$ , the kinetic and chemical potential terms diagonalize trivially. The pairing term requires antisymmetrization due to fermionic statistics ( $c_k c_{-k} = -c_{-k} c_k$ ):

$$\Delta \sum_k e^{ik} c_k c_{-k} = \Delta \sum_{k>0} (e^{ik} - e^{-ik}) c_k c_{-k} = 2i\Delta \sum_{k>0} \sin(k) c_k c_{-k} \quad (8)$$

This isolates the odd-parity  $p$ -wave superconducting nature of the model in momentum space.

## Appendix C: BdG Formalism and Pseudospin

Use the Nambu spinor  $\Psi_k = \begin{pmatrix} c_k \\ c_{-k}^\dagger \end{pmatrix}$  to cast  $H$  into a bilinear form:

$$H = \sum_{k>0} \Psi_k^\dagger h(k) \Psi_k, \quad h(k) = \begin{pmatrix} \xi_k & 2i\Delta \sin(k) \\ -2i\Delta \sin(k) & -\xi_k \end{pmatrix} \quad (9)$$

where the normal-state dispersion is  $\xi_k = -\mu - 2t \cos(k)$ .

Expanding  $h(k)$  in Pauli matrices:

$$h(k) = n_y(k)\sigma^y + n_z(k)\sigma^z \implies \begin{cases} n_z(k) = -\mu - 2t \cos(k) \\ n_y(k) = -2\Delta \sin(k) \end{cases} \quad (10)$$

Diagonalizing this yields the bulk dispersion  $E_k = \pm \sqrt{n_z(k)^2 + n_y(k)^2}$ .

# Appendix D: BdG extraction — sign filter and double counting

## The problem with `evals[evals $\geq$ 0]`

Particle-hole symmetry guarantees  $\pm E$  pairs *analytically*. Numerically, near-zero Majorana modes at large  $L$  can round to tiny *negative* values — a sign filter silently drops them, and the “lowest mode” jumps to the bulk gap, creating spurious spikes.

Sign-filter artefact vs stable  $|E|$  extraction for  $E_0$  ( $L = 30$ ,  $t = \Delta = 1$ , OBC)

