

# Spatial Analysis and Modeling (GIST 4302/5302)

Guofeng Cao

Department of Geosciences


Texas Tech University

# Outline of This Week


- Last week, we learned:
  - spatial point pattern analysis (PPA)
  - focus on location distribution of ‘events’
- This week, we will learn:
  - spatial autocorrelation
  - global measures of spatial autocorrelation
  - local measure of spatial autocorrelation

# Spatial Autocorrelation


- Tobler's first law of geography
- Spatial auto/cross correlation



If like values tend to cluster together, then the field exhibits high **positive spatial autocorrelation**



If there is no apparent relationship between attribute value and location then there is **zero spatial autocorrelation**



If like values tend to be located away from each other, then there is **negative spatial autocorrelation**




# Spatial Autocorrelation

- Spatial autocorrelation is everywhere
  - Spatial point pattern
    - K, F, G functions
    - Kernel functions
  - Areal/lattice (this topic)
  - Geostatistical data (next topic)

# Spatial Autocorrelation of Areal Data

## *Positive spatial autocorrelation*

- high values surrounded by nearby high values
- intermediate values surrounded by nearby intermediate values
- low values surrounded by nearby low values




Source: Ron Briggs of UT Dallas



# Negative spatial autocorrelation

- high values surrounded by nearby low values
- intermediate values surrounded by nearby intermediate values
- low values surrounded by nearby high values

competition for space



Grocery store density



Source: Ron Briggs of UT Dallas

# Spatial Weight Matrix

- **Core** concept in statistical analysis of areal data
- Two steps involved:
  - define which relationships between observations are to be given a nonzero weight, i.e., define spatial neighbors
  - assign weights to the neighbors
- Making the neighbors and weights is not easy as it seems to be
  - Which states are near Texas?






# Spatial Neighbors


- **Contiguity-based neighbors**
  - Zone  $i$  and  $j$  are neighbors if zone  $i$  is contiguity or adjacent to zone  $j$
  - But what constitutes contiguity?
- **Distance-based neighbors**
  - Zone  $i$  and  $j$  are neighbors if the distance between them are less than the threshold distance
  - But what distance do we use?

# Contiguity-based Spatial Neighbors


- Sharing a border or boundary
  - Rook: sharing a border
  - Queen: sharing a border or a point




rook



queen




Hexagons



Irregular

Which use?

# Example




**Fig. 9.3.** (a) Queen-style census tract contiguities, Syracuse; (b) Rook-style contiguity differences shown as thicker lines

Source: Bivand and Pebesma and Gomez-Rubio


# Higher-Order Contiguity

1<sup>st</sup>  
order


Nearest  
neighbor



rook






hexagon



queen

2<sup>nd</sup>  
order

Next  
nearest  
neighbor



# Distance-based Neighbors

- How to measure distance between polygons?
- Distance metrics
  - 2D Cartesian distance (projected data)
  - 3D spherical distance/great-circle distance (lat/long data)
    - Haversine formula

*Haversine*  $a = \sin^2(\Delta\phi/2) + \cos(\phi_1) \cdot \cos(\phi_2) \cdot \sin^2(\Delta\lambda/2)$


*formula:*  $c = 2 \cdot \text{atan2}(\sqrt{a}, \sqrt{1-a})$

$d = R \cdot c$

*where  $\phi$  is latitude,  $\lambda$  is longitude,  $R$  is earth's radius (mean radius = 6,371km)*

# Distance-based Neighbors

- $k$ -nearest neighbors




**Fig. 9.5.** (a)  $k = 1$  neighbours; (b)  $k = 2$  neighbours; (c)  $k = 4$  neighbours

Source: Bivand and Pebesma and Gomez-Rubio


# Distance-based Neighbors

- thresh-hold distance (buffer)



**Fig. 9.6.** (a) Neighbours within 1,158 m; (b) neighbours within 1,545 m; (c) neighbours within 2,317 m

# Neighbor/Connectivity Histogram




Source: Bivand and Pebesma and Gomez-Rubio




# Recap: Box-plot

- Help indicate the degree of dispersion and skewness and identify outliers
  - Non-parametric
  - 25%, 50%, 75% percentiles
  - end of the hinge could mean differently depending on implementation
  - Points outside the range are usually taken as outliers



# Spatial Weight Matrix


- Spatial weights can be seen as a list of weights indexed by a list of neighbors
- If zone  $j$  is not a neighbor of zone  $i$ , weights  $W_{ij}$  will set to zero
  - The weight matrix can be illustrated as an image
  - Sparse matrix



# A Simple Example for Rook case

- Matrix contains a:
  - 1 if share a border
  - 0 if do not share a border

4 areal units




Common border

4x4 matrix

	A	B	C	D
A	0	1	1	0
B	1	0	0	1
C	1	0	0	1
D	0	1	1	0

- 1 Washington
- 2 Oregon
- 3 California
- 4 Arizona
- 5 Nevada
- 6 Idaho
- 7 Montana
- 8 Wyoming
- 9 Utah
- 10 New Mexico
- 11 Texas
- 12 Oklahoma
- 13 Colorado
- 14 Kansas
- 15 Nebraska
- 16 South Dakota
- 17 North Dakota
- 18 Minnesota
- 19 Iowa
- 20 Missouri
- 21 Arkansas
- 22 Louisiana
- 23 Mississippi
- 24 Tennessee
- 25 Kentucky
- 26 Illinois
- 27 Wisconsin
- 28 Michigan
- 29 Indiana
- 30 Ohio
- 31 West Virginia
- 32 Florida
- 33 Alabama
- 34 Georgia
- 35 South Carolina
- 36 North Carolina
- 37 Virginia
- 38 Maryland
- 39 Delaware
- 40 District of Columbia
- 41 New Jersey
- 42 Pennsylvania
- 43 New York
- 44 Connecticut
- 45 Rhode Island
- 46 Massachusetts
- 47 New Hampshire
- 48 Vermont
- 49 Maine



**Sparse Contiguity Matrix for US States -- obtained from Anselin's web site (see powerpoint for link)**

<b>Name</b>	<b>Fips</b>	<b>Ncount</b>	<b>N1</b>	<b>N2</b>	<b>N3</b>	<b>N4</b>	<b>N5</b>	<b>N6</b>	<b>N7</b>	<b>N8</b>
Alabama	1	4	28	13	12	47				
Arizona	4	5	35	8	49	6	32			
Arkansas	5	6	22	28	48	47	40	29		
California	6	3	4	32	41					
Colorado	8	7	35	4	20	40	31	49	56	
Connecticut	9	3	44	36	25					
Delaware	10	3	24	42	34					
District of Columbia	11	2	51	24						
Florida	12	2	13	1						
Georgia	13	5	12	45	37	1	47			
Idaho	16	6	32	41	56	49	30	53		
Illinois	17	5	29	21	18	55	19			
Indiana	18	4	26	21	17	39				
Iowa	19	6	29	31	17	55	27	46		
Kansas	20	4	40	29	31	8				
Kentucky	21	7	47	29	18	39	54	51	17	
Louisiana	22	3	28	48	5					
Maine	23	1	33							
Maryland	24	5	51	10	54	42	11			
Massachusetts	25	5	44	9	36	50	33			
Michigan	26	3	18	39	55					
Minnesota	27	4	19	55	46	38				
Mississippi	28	4	22	5	1	47				
Missouri	29	8	5	40	17	21	47	20	19	31
Montana	30	4	16	56	38	46				
Nebraska	31	6	29	20	8	19	56	46		
Nevada	32	5	6	4	49	16	41			
New Hampshire	33	3	25	23	50					
New Jersey	34	3	10	36	42					
New Mexico	35	5	48	40	8	4	49			
New York	36	5	34	9	42	50	25			
North Carolina	37	4	45	13	47	51				
North Dakota	38	3	46	27	30					
Ohio	39	5	26	21	54	42	18			
Oklahoma	40	6	5	35	48	29	20	8		
Oregon	41	4	6	32	16	53				
Pennsylvania	42	6	24	54	10	39	36	34		
Rhode Island	44	2	25	9						
South Carolina	45	2	13	37						
South Dakota	46	6	56	27	19	31	38	30		
Tennessee	47	8	5	28	1	37	13	51	21	29
Texas	48	4	22	5	35	40				
Utah	49	6	4	8	35	56	32	16		
Vermont	50	3	36	25	33					
Virginia	51	6	47	37	24	54	11	21		
Washington	53	2	41	16						
West Virginia	54	5	51	21	24	39	42			
Wisconsin	55	4	26	17	19	27				
Wyoming	56	6	49	16	31	8	46	30		

# Style of Spatial Weight Matrix

- Row
  - a weight of unity for each neighbor relationship
- Row standardization
  - Symmetry not guaranteed
  - can be interpreted as allowing the calculation of average values across neighbors
- General spatial weights based on distances

# Row vs. Row standardization

A	B	C
D	E	F

Divide each  
number by the  
**row sum**

Total number of neighbors  
--some have more than others


	A	B	C	D	E	F	Row Sum
A	0	1	0	1	0	0	2
B	1	0	1	0	1	0	3
C	0	1	0	0	0	1	2
D	1	0	0	0	1	0	2
E	0	1	0	1	0	1	3
F	0	0	1	0	1	0	2

Row standardized  
--usually use this

	A	B	C	D	E	F	Row Sum
A	0.0	0.5	0.0	0.5	0.0	0.0	1
B	0.3	0.0	0.3	0.0	0.3	0.0	1
C	0.0	0.5	0.0	0.0	0.0	0.5	1
D	0.5	0.0	0.0	0.0	0.5	0.0	1
E	0.0	0.3	0.0	0.3	0.0	0.3	1
F	0.0	0.0	0.5	0.0	0.5	0.0	1

# General Spatial Weights Based on Distance

- Decay functions of distance
  - Most common choice is the inverse (reciprocal) of the distance between locations  $i$  and  $j$  ( $w_{ij} = 1/d_{ij}$ )
  - Other functions also used
    - inverse of squared distance ( $w_{ij} = 1/d_{ij}^2$ ), or
    - negative exponential ( $w_{ij} = e^{-d}$  or  $w_{ij} = e^{-d^2}$ )





# Measure of Spatial Autocorrelation

# Global Measures and Local Measures


- Global Measures
  - A single value which applies to the entire data set
    - The same pattern or process occurs over the entire geographic area
    - An average for the entire area
- Local Measures
  - A value calculated for each observation unit
    - Different patterns or processes may occur in different parts of the region
    - A unique number for each location
- Global measures usually can be decomposed into a combination of local measures

# Global Measures and Local Measures


- Global Measures
  - Join Count
  - Moran's I
- Local Measures
  - Local Moran's I

# Join (or Joint or Joins) Count Statistic


Positive autocorrelation



No autocorrelation



Negative autocorrelation



Rook's case

$$J_{BB} = 27$$

$$J_{WW} = 27$$

$$J_{BW} = 6$$

Queen's case

$$J_{BB} = 47$$

$$J_{WW} = 47$$

$$J_{BW} = 16$$

$$J_{BB} = 6$$

$$J_{WW} = 19$$

$$J_{BW} = 35$$

$$J_{BB} = 14$$

$$J_{WW} = 40$$

$$J_{BW} = 56$$

$$J_{BB} = 0$$

$$J_{WW} = 0$$

$$J_{BW} = 60$$

$$J_{BB} = 25$$

$$J_{WW} = 25$$

$$J_{BW} = 60$$

- 60 for Rook Case
- 110 for Queen Case

# Join Count: Test Statistic

Test Statistic given by:  $Z = \frac{\text{Observed} - \text{Expected}}{\text{SD of Expected}}$

**Expected** = random pattern generated by tossing a coin in each cell.

Expected given by: Standard Deviation of Expected (standard error) given by:

$$E(J_{BB}) = kp_B^2$$

$$E(s_{BB}) = \sqrt{kp_B^2 + 2mp_B^3 - (k + 2m)p_B^4}$$

$$E(J_{WW}) = kp_W^2$$

$$E(s_{WW}) = \sqrt{kp_W^2 + 2mp_W^3 - (k + 2m)p_W^4}$$

$$E(J_{BW}) = 2kp_Bp_W$$

$$E(s_{BW}) = \sqrt{2(k + m)p_Bp_W - 4(k + 2m)p_B^2p_W^2}$$

Where: k is the total number of joins (neighbors)

$p_B$  is the expected proportion Black, if random

$p_W$  is the expected proportion White

m is calculated from k according to:

$$m = \frac{1}{2} \sum_{i=1}^n k_i(k_i - 1)$$

# Gore/Bush Presidential Election 2000



	<b>Actual</b>
<b>Jbb</b>	<b>60</b>
<b>Jgg</b>	<b>21</b>
<b>Jbg</b>	<b>28</b>
<b>Total</b>	<b>109</b>

# Join Count Statistic for Gore/Bush 2000 by State

<b>candidates</b>	probability
Bush	0.49885
Gore	0.50115

	Actual	Expected	Stan Dev	Z-score
Jbb	60	27.125	8.667	3.7930
Jgg	21	27.375	8.704	-0.7325
Jbg	28	54.500	5.220	-5.0763
Total	109	109.000		

- The expected number of joins is calculated based on the proportion of votes each received in the election (for Bush =  $109 * .499 * .499 = 27.125$ )
- There are far more Bush/Bush joins (actual = 60) than would be expected (27)
  - Positive autocorrelation
- There are far fewer Bush/Gore joins (actual = 28) than would be expected (54)
  - Positive autocorrelation
- No strong clustering evidence for Gore (actual = 21 slightly less than 27.375)

# Moran's I

- The most common measure of Spatial Autocorrelation
- Use for points or polygons
  - Join Count statistic only for polygons
- Use for a continuous variable (any value)
  - Join Count statistic only for binary variable (1,0)



Patrick Alfred Pierce Moran (1917-1988)



# Formula for Moran's I

$$I = \frac{N \sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i - \bar{x})(x_j - \bar{x})}{\left( \sum_{i=1}^n \sum_{j=1}^n w_{ij} \right) \sum_{i=1}^n (x_i - \bar{x})^2}$$

- Where:

- $N$  is the number of observations (points or polygons)
- $\bar{x}$  is the mean of the variable
- $x_i$  is the variable value at a particular location
- $x_j$  is the variable value at another location
- $w_{ij}$  is a weight indexing location of  $i$  relative to  $j$

# Moran's I



- Expectation of Moran's I under no spatial autocorrelation

$$E(I) = -1 / (N-1)$$

- Variance of Moran's is complex and exact equation is given at textbook d&G&L
- $[-1, 1]$

# Moran's $I$ and Correlation Coefficient

- **Correlation Coefficient [-1, 1]**
  - Relationship between two different variables
- **Moran's  $I$  [-1, 1]**
  - Spatial autocorrelation and often involves one (spatially indexed) variable only
  - Correlation between observations of a spatial variable at location  $X$  and “spatial lag” of  $X$  formed by averaging all the observation at neighbors of  $X$



# Correlation Coefficient

Note the similarity of the numerator (top) to the measures of spatial association discussed earlier if we view  $Y_i$  as being the  $X_i$  for the neighboring polygon

**(see next slide)**

$$\frac{\sum_{i=1}^n (y_i - \bar{y})(x_i - \bar{x})/n}{\sqrt{\frac{\sum_{i=1}^n (y_i - \bar{y})^2}{n}} \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}}}$$

$$\frac{N \sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i - \bar{x})(x_j - \bar{x})}{(\sum_{i=1}^n \sum_{j=1}^n w_{ij}) \sum_{i=1}^n (x_i - \bar{x})^2}$$

Spatial  
auto-correlation

=

$$\frac{\sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i - \bar{x})(x_j - \bar{x}) / \sum_{i=1}^n \sum_{j=1}^n w_{ij}}{\sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}} \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}}}$$

# Correlation Coefficient

$$\frac{\sum_{i=1}^n 1(y_i - \bar{y})(x_i - \bar{x})/n}{\sqrt{\frac{\sum_{i=1}^n (y_i - \bar{y})^2}{n}} \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}}}$$

Spatial weights

Yi is the Xi for the neighboring polygon →


$$\frac{N \sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i - \bar{x})(x_j - \bar{x})}{(\sum_{i=1}^n \sum_{j=1}^n w_{ij}) \sum_{i=1}^n (x_i - \bar{x})^2} =$$

$$\frac{\sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i - \bar{x})(x_j - \bar{x}) / \sum_{i=1}^n \sum_{j=1}^n w_{ij}}{\sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}} \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}}}$$

Moran's I


# Moran Scatter Plots

We can draw a scatter diagram between these two variables (in standardized form):  $X$  and  $\text{lag-}X$  (or  $W\_X$ )




The slope of this *regression line* is  
Moran's I

# Moran Scatter Plots



# Moran Scatterplot: Example





# Moran's I for rate-based data

- Moran's I is often calculated for rates, such as crime rates (e.g. number of crimes per 1,000 population) or infant mortality rates (e.g. number of deaths per 1,000 births)
- An adjustment should be made, especially if the denominator in the rate (population or number of births) varies greatly (as it usually does)
- Adjustment is known as the *EB adjustment*:
  - see Assuncao-Reis *Empirical Bayes Standardization Statistics in Medicine*, 1999
- *GeoDA* software includes an option for this adjustment

# Statistical Significance Tests for Moran's I


- Based on the normal frequency distribution with

$$Z = \frac{I - E(I)}{S_{error}(I)}$$

Where: I is the calculated value for Moran's I  
from the sample  
E(I) is the expected value if random  
S is the standard error

- Statistical significance test
  - Monte Carlo test, as we did for spatial pattern analysis
  - Permutation test
    - Non-parametric
    - Data-driven, no assumption of the data
    - Implemented in GeoDa

# Test Statistic for Normal Frequency Distribution



*Null Hypothesis*: no spatial autocorrelation

\*Moran's  $I = 0$

*Alternative Hypothesis*: spatial autocorrelation exists

\*Moran's  $I > 0$

Reject *Null Hypothesis* if Z test statistic  $> 1.96$  (or  $< -1.96$ )

---less than a 5% chance that, in the population, there is no spatial autocorrelation

---95% confident that spatial auto correlation exists

*Null Hypothesis:* no spatial autocorrelation

\*Moran's  $I = 0$

*Alternative Hypothesis:* spatial autocorrelation exists

\*Moran's  $I > 0$

Reject *Null Hypothesis* if  $Z$  test statistic  $> 1.96$  (or  $< -1.96$ )

---less than a 5% chance that, in the population, there is no spatial autocorrelation

---95% confident that spatial auto correlation exists

# Local Measures of Spatial Autocorrelation

# Local Indicators of Spatial Association (LISA)

- Local versions of *Moran's I*
- Moran's I is most commonly used, and the local version is often called Anselin's LISA, or just LISA


**See:**

Luc Anselin 1995 *Local Indicators of Spatial Association-LISA* Geographical Analysis 27: 93-115

# Local Indicators of Spatial Association (LISA)

- The statistic is calculated for each areal unit in the data
- For each polygon, the index is calculated based on neighboring polygons with which it shares a border
- A measure is available for each polygon, these can be mapped to indicate how spatial autocorrelation varies over the study region
- Each index has an associated test statistic, we can also map which of the polygons has a statistically significant relationship with its neighbors, and show type of relationship

# Example:





# Calculating Anselin's LISA

- The local Moran statistic for areal unit  $i$  is:

$$I_i = z_i \sum_j w_{ij} z_j$$

where  $z_i$  is the original variable  $x_i$  in  
“standardized form”

$$z_i = \frac{x_i - \bar{x}}{SD_x}$$

or it can be in “deviation form”


$$x_i - \bar{x}$$

and  $w_{ij}$  is the spatial weight

The summation  $\sum_j$  is across each row  $i$  of the spatial weights matrix.

An example follows

Contiguity Matrix		1	2	3	4	5	6	7	Sum	Neighbors	Illiteracy
		Anhui	Zhejiang	Jiangxi	Jiangsu	Henan	Hubei	Shanghai			
Anhui	1	0	1	1	1	1	1	0	5	6 5 4 3 2	14.49
Zhejiang	2	1	0	1	1	0	0	1	4	7 4 3 1	9.36
Jiangxi	3	1	1	0	0	0	1	0	3	6 2 1	6.49
Jiangsu	4	1	1	0	0	0	0	1	3	7 2 1	8.05
Henan	5	1	0	0	0	0	1	0	2	6 1	7.36
Hubei	6	1	0	1	0	1	0	0	3	1 3 5	7.69
Shanghai	7	0	1	0	1	0	0	0	2	2 4	3.97



Source: Ron Briggs of UT Dallas

# Contiguity Matrix and Row Standardized Spatial Weights Matrix

**Contiguity Matrix**

		1	2	3	4	5	6	7	
	Code	Anhui	Zhejiang	Jiangxi	Jiangsu	Henan	Hubei	Shanghai	Sum
Anhui	1	0	1	1	1	1	1	0	5
Zhejiang	2	1	0	1	1	0	0	1	4
Jiangxi	3	1	1	0	0	0	1	0	3
Jiangsu	4	1	1	0	0	0	0	①	③
Henan	5	1	0	0	0	0	1	0	2
Hubei	6	1	0	1	0	1	0	0	3
Shanghai	7	0	1	0	1	0	0	0	2

**Row Standardized Spatial Weights Matrix**

		Code	Anhui	Zhejiang	Jiangxi	Jiangsu	Henan	Hubei	Shanghai	Sum
Anhui	1		0.00	0.20	0.20	0.20	0.20	0.20	0.00	1
Zhejiang	2		0.25	0.00	0.25	0.25	0.00	0.00	0.25	1
Jiangxi	3		0.33	0.33	0.00	0.00	0.00	0.33	0.00	1
Jiangsu	4		0.33	0.33	0.00	0.00	0.00	0.00	①	1
Henan	5		0.50	0.00	0.00	0.00	0.00	0.50	0.00	1
Hubei	6		0.33	0.00	0.33	0.00	0.33	0.00	0.00	1
Shanghai	7		0.00	0.50	0.00	0.50	0.00	0.00	0.00	1

Source: Ron Briggs of UT Dallas

# Calculating standardized (z) scores

Deviations from Mean and z scores.

	X	X-Xmean	X-Mean <sup>2</sup>	z ← $Z_i = \frac{x_i - \bar{x}}{SD_x}$
Anhui	14.49	6.29	39.55	2.101
Zhejiang	9.36	1.16	1.34	0.387
Jiangxi	6.49	(1.71)	2.93	(0.572)
Jiangsu	8.05	(0.15)	0.02	(0.051)
Henan	7.36	(0.84)	0.71	(0.281)
Hubei	7.69	(0.51)	0.26	(0.171)
Shanghai	3.97	(4.23)	17.90	(1.414)

**Mean and Standard Deviation**

Sum	57.41	0.00	62.71
Mean	57.41 / 7 =		8.20
Variance	62.71 / 7 =		8.96
SD	√ 8.96 =		2.99

# Calculating LISA

## Row Standardized Spatial Weights Matrix

	Code	Anhui	Zhejiang	Jiangxi	Jiangsu	Henan	Hubei	Shanghai
Anhui	1	0.00	0.20	0.20	0.20	0.20	0.20	0.00
Zhejiang	2	0.25	0.00	0.25	0.25	0.00	0.00	0.25
Jiangxi	3	0.33	0.33	0.00	0.00	0.00	0.33	0.00
Jiangsu	4	0.33	0.33	0.00	0.00	0.00	0.00	0.33
Henan	5	0.50	0.00	0.00	0.00	0.00	0.50	0.00
Hubei	6	0.33	0.00	0.33	0.00	0.33	0.00	0.00
Shanghai	7	0.00	0.50	0.00	0.50	0.00	0.00	0.00

$w_{ij}$

## Z-Scores for row Province and its potential neighbors

		Anhui	Zhejiang	Jiangxi	Jiangsu	Henan	Hubei	Shanghai
	Zi							
Anhui	2.101	<b>2.101</b>	0.387	(0.572)	(0.051)	(0.281)	(0.171)	(1.414)
Zhejiang	0.387	2.101	<b>0.387</b>	(0.572)	(0.051)	(0.281)	(0.171)	(1.414)
Jiangxi	(0.572)	2.101	0.387	<b>(0.572)</b>	(0.051)	(0.281)	(0.171)	(1.414)
Jiangsu	(0.051)	2.101	0.387	(0.572)	<b>(0.051)</b>	(0.281)	(0.171)	(1.414)
Henan	(0.281)	2.101	0.387	(0.572)	(0.051)	<b>(0.281)</b>	(0.171)	(1.414)
Hubei	(0.171)	2.101	0.387	(0.572)	(0.051)	(0.281)	<b>(0.171)</b>	(1.414)
Shanghai	(1.414)	2.101	0.387	(0.572)	(0.051)	(0.281)	(0.171)	<b>(1.414)</b>

$z_j$

$$I_i = z_i \sum_j w_{ij} z_j$$

## Spatial Weight Matrix multiplied by Z-Score Matrix (cell by cell multiplication)

		Anhui	Zhejiang	Jiangxi	Jiangsu	Henan	Hubei	Shanghai	SumWijZj
	Zi								
Anhui	<b>2.101</b>	-	0.077	(0.114)	(0.010)	(0.056)	(0.034)	-	<b>(0.137)</b>
Zhejiang	<b>0.387</b>	0.525	-	(0.143)	(0.013)	-	-	(0.353)	<b>0.016</b>
Jiangxi	<b>(0.572)</b>	0.700	0.129	-	-	-	(0.057)	-	<b>0.772</b>
Jiangsu	<b>(0.051)</b>	0.700	0.129	-	-	-	-	(0.471)	<b>0.358</b>
Henan	<b>(0.281)</b>	1.050	-	-	-	-	(0.085)	-	<b>0.965</b>
Hubei	<b>(0.171)</b>	0.700	-	(0.191)	-	(0.094)	-	-	<b>0.416</b>
Shanghai	<b>(1.414)</b>	-	0.194	-	(0.025)	-	-	-	<b>0.168</b>

$w_{ij}z_j$


LISA	Lisa from GeoDA
<b>-0.289</b>	-0.248
<b>0.006</b>	0.005
<b>-0.442</b>	-0.379
<b>-0.018</b>	-0.016
<b>-0.271</b>	-0.233
<b>-0.071</b>	-0.061
<b>-0.238</b>	-0.204

Source: Ron Briggs of UT Dallas


# Bivariate LISA

- Moran's I is the correlation between X and Lag-X--the same variable but in nearby areas
  - Univariate Moran's I
- Bivariate Moran's I is a correlation between X and a different variable in nearby areas.

## Moran Scatter Plot for GDI vs AL





## Moran Significance Map for GDI vs. AL




# Bivariate LISA

## and the Correlation Coefficient

- Correlation Coefficient is the relationship between two different variables in the same area
- Bivariate LISA is a correlation between two different variables in an area and in nearby areas.



# Bivariate Moran Scatter Plot





# Summary

- Spatial autocorrelation of areal data
- Spatial weight matrix
- Measures of spatial autocorrelation
- Global Measure
  - Moran's I/General G and  $G^*$
- Local
  - LISA: Moran's I/General G and  $G^*$
  - Bivariate LISA
    - Significance test

- End of this topic