


WORKING WITH FUNCTIONS: HW

Multiple-Choice

- 1. If the function f is defined by f(x) = 3x + 2, and if f(a) = 17, what is the value of a?
 - (A) 5
 - (B) 9
 - (C) 10
 - (D) 11
- 2. A function f is defined such that f(1) = 2, f(2) = 5, and f(n) = f(n-1) f(n-2) for all integer values of f greater than 2. What is the value of f(4)?
 - (A) -8
 - (B) -2
 - (C) 2
 - (D) 8




- 3. The graph of y = f(x) is shown above. If $-4 \le x \le 10$, for how many values of x does f(x) = 2?
 - (A) None
 - (B) One
 - (C) Two
 - (D) Three
- 4. If function *f* is defined by f(x) = 5x + 3, then which expression represents 2f(x) 3?
 - (A) 10x 3
 - (B) 10x + 3
 - (C) 10x
 - (D) 3
- 5. If the function *k* is defined by $k(h) = (h + 1)^2$, then k(x 2) =
 - (A) $x^2 x$
 - (B) $x^2 2x$

- (C) $x^2 2x + 1$
- (D) $x^2 + 2x 1$

х	1	2	3	4	5	
f(x)	f(x) 3		5	6	7	
х	3	4	5	6	7	
g(x)	4	6	8	10	12	


- 6. The accompanying tables define functions f and g. What is g(f(3))?
 - (A) 4
 - (B) 6
 - (C) 8
 - (D) 10
- 7. In 2014, the United States Postal Service charged \$0.48 to mail a first-class letter weighing up to 1 oz. and \$0.21 for each additional ounce. Based on these rates, which function would determine the cost, in dollars, c(z), of mailing a first-class letter weighing z ounces where z is an integer greater than 1?
 - (A) c(z) = 0.48z + 0.21
 - (B) c(z) = 0.21z + 0.48
 - (C) c(z) = 0.48(z 1) + 0.21
 - (D) c(z) = 0.21(z 1) + 0.48




- 8. Based on the graphs of functions f and g shown in the accompanying figure, for which values of x between -3 and 3 is $f(x) \ge g(x)$?
 - I. $-2 \le x \le 0$
 - II. $0 \le x \le 1$
 - III. $1 \le x \le 3$
 - (A) I only
 - (B) II only
 - (C) III only
 - (D) I and II



- f(2n) = 4f(n) for all integers n
- f(3) = 9
- 9. If function *f* satisfies the above two conditions for all positive integers *n*, which equation could represent function *f* ?
 - (A) f(n) = 9
 - (B) $f(n) = n^2$
 - (C) f(n) = 3n
 - (D) f(n) = 2n + 3



- 10. If in the accompanying figure (p, q) lies on the graph of y = f(x) and $0 \le p \le 5$, which of the following represents the set of corresponding values of q?
 - (A) $-5 \le q \le 15$
 - (B) $-5 \le q \le 10$
 - (C) $-5 \le q \le 5$
 - (D) $5 \le q \le 10$



- 11. The accompanying figure shows the graph of y = f(x). If function g is defined by g(x) = f(x + 4), then g(-1) could be
 - (A) -2
 - (B) 3
 - (C) 4
 - (D) 5



х	f(x)	g(x)		
1	2	3		
2	4	5		
3	5	1		
4	3	2		
5	1	4		


Questions 12–13 refer to the accompanying table, which gives the values of functions f and g for integer values of x from 1 to 5, inclusive.

- 12. According to the table, if f(5) = p, what is the value of g(p)?
 - (A) 1
 - (B) 2
 - (C) 3
 - (D) 4
- 13. Function h is defined by h(x) = 2f(x) 1, where function f is defined in the accompanying table. What is the value of g(k) when h(k) = 5?
 - (A) 1
 - (B) 2
 - (C) 3
 - (D) 4


х	0	1	4	5			2		
f(x)	-2	5	0	2	g(x)	2	-1	1	5

- 14. Some values of functions f and g are given by the tables above. What is the value of g(f(5))?
 - (A) -1
 - (B) 1
 - (C) 2
 - (D) 5
- 15. In 2012, a retail chain of fast food restaurants had 68 restaurants in California and started to expand nationally by adding 9 new restaurants each year thereafter. At this rate, which of the following functions *f* represent the number of restaurants there will be in this retail chain *n* years after 2012 assuming none of these restaurants close?
 - (A) f(n) = 2,012 + 9n
 - (B) f(n) = 9 + 68n
 - (C) f(n) = 68 + 9(n 2,012)
 - (D) f(n) = 68 + 9n
- According to market research, the number of magazine subscriptions that can be sold can be estimated using the function

$$n(p) = \frac{5,000}{4p-k},$$



- 5. What is the integer value of 2f(-1) + 3f(1)?
- 6. If *n* represents the number of different values of *x* for which f(x) = 2 and *m* represents the number of different values of *x* for which f(x) = 4, what is the value of *mn*?
- 7. Let *g* be the function defined by g(x) = x 1. If $\frac{1}{2}g(c) = 4$, what is the value of g(2c)?



8. The figure above shows the graph of function h. If function f is defined by f(x) = h(2x) + 1, what is the value of f(-1)?