

# BLT Meeting output (Validation) burnin run v1

Eduardo Zanette


20/04/2022

## **Abstract**

This document was produced for the purpose of presentation to Laurence, Eckhard and Ronald on 25/04/2022, the second BLT Model Meeting while in Germany. For documentation reasons, this run was made with the version v1 of the model. Here we analyze the comparison of empirical data with one run ( $n = 30$  days), but we also differentiate between the whole run (30 days) and the 20 last days, because the `step_forget` parameter in the model seemed to make the tamarin move more after a “burn-in period” of 10 days. The results show that the burn-in period does not make a difference in the v1 version of the model.


Guareí empirical (all data, 28 days) and July (6 days)

```
## Warning: Removed 276 rows containing missing values (geom_point).
```




```
## Warning: Removed 26 rows containing missing values (geom_point).
```


### Empirical (July only, 6 days)



## Guareí simulated



Simulated (20 days after burn in of 10 days – Resources from July)




## Validation Patterns

We are not checking seed dispersal patterns now, just movement and ranging patterns.

## Dayly Path Lenght (DPL)

Calculate and plot DPL



## Home Range

Comparison between methods: <https://doi.org/10.1111/2041-210X.13786>

## Turning Angles and Step Lengths

```
## [1] "Empirical (July, 6 d) Mean Relative Angle (degrees) = 0.06"
```

```
## [1] "Simulated (July, 30 d) Mean Relative Angle (degrees) 0.01"
```

```
## [1] "Simulated (July, 20 d, after burn-in run of 10 days) Mean Relative Angle (degrees) 0.01"
```

```
## [1] "Mean Relative Angle (degrees) : 3.60006621892628"
```


```
## [1] "Mean Relative Angle (degrees) : 0.386297408087265"
```

```
## [1] "Mean Relative Angle (degrees) : -0.119157674256658"
```


```
## [1] "Mean Step length (meters) : 16.88761495443"
```

```
## [1] "Mean Step lenght (meters) : 10.2508063826726"
```


```
## [1] "Mean Step lenght (meters) : 10.3488574898088"
```




**Empirical (July, 6 d)**



**Simulated (July, 30 d)**



**Simulated (July, 20 d,  
after burn-in run of 10 days)**



## Revisitations (coming soon)

### Conclusions

The *four* problems with this model are:

- 1) After the burnin they spend too much time foraging
- 2) Tamarins are still not properly bouncing at the border of the fragment (while foraging)
- 3) Foraging occurs for too long
- 4) Constant step lenght