STAT 315 Notes Fall 2026

Colorado State University

Preliminaries

Getting Started With R

Installing R

R is the base version of a free, open-source statistical computing program. RStudio is a nice graphic user interface (GUI) that works in conjunction with R. We will be using RStudio this semester as it is easier to learn how to use and has many nice shortcuts that aren't available in the standard R download.

Note: R and R Studio can not be installed on Chromebooks or other tablets/mini-computers. If you only have a tablet or a mini-computer, the free version of RStudio Cloud should work: R Studio Cloud. The free version allows 25 hours of use per month. I think this is likely enough for this course, but be sure to log out!

To begin using RStudio, you will need to install "R" first and then install "RStudio" on your computer.

Step 1: Download R

- Visit https://www.r-project.org/
- Click *CRAN* under *Download*
- Select any of the mirrors
- Click the appropriate link for your type of system (Mac, Windows, Linux)
- Download R on this next page.
  - For Windows, this will say *install R for the first time*
  - For Mac, this will be under *Latest release* and will be something like *R-4.1.0.pkg* – the numbers may differ depending on the most recent version
- Install R on your computer

Step 2: Download RStudio

- Visit https://posit.co/downloads
- Click to download
- Click the appropriate link for your type of system (Mac, Windows, Linux)
- Install RStudio on your computer

Step 3: Verify RStudio is working

- Open RStudio
- Let's enter a small dataset and calculate the average to make sure everything is working correctly.
- In the console, type in the following dataset of five homework scores:

```
# Assignment using =
x = c(90,75,100,85,90)

# Alternative assignment using <-
x <- c(90,75,100,85,90)

# More on the differences between = and <- at:
# https://www.r-bloggers.com/2014/01/difference-between-assignment-operators-in-r/
```

In the console, calculate the average of the five homework scores:

```
mean(x)
```

```
## [1] 88
```

Did you find the average of the five homework scores was 88? If so, you should be set up correctly!

Basic Concepts of Statistics

In this chapter, we'll introduce some of the basic concepts and definitions that are fundamental to the study of Probability and Statistics. In particular, sampling and sample statistics will be examined.

Populations, Samples, and Processes

A ***population*** is a well-defined complete collection of objects.


A ***sample*** is a subset of the population.

A ***population parameter*** is a numerical value that describes a characteristic of a population.

A ***sample statistic*** is a numerical value that describes a characteristic of a sample.

A student is interested in the average grade point average (GPA) of CSU students. She collects the GPAs of eight friends and finds an average of 2.87. For this example, determine the population, sample, parameter, and statistic.

Illustration of Population and Sample



Methods of Collecting a Sample

Simple Random Sampling (SRS) is a basic sampling method used in statistics where every individual or element in a population has an equal chance of being selected. This method ensures that the sample is unbiased.

Suppose we want to collect a simple random sample of CSU student GPAs. How might we do this?

Convenience Sampling is a sampling method where the sample is taken from a group of people that are easy to reach or readily available. This method relies on selecting individuals who are most conveniently accessible to the researcher, rather than using a random or systematic approach.

Suppose we want to collect a convenience sample of CSU student GPAs. How might we do this?

Stratified Sampling is a probability sampling method where the population is divided into distinct subgroups or "strata" that share similar characteristics, and a random sample is then taken from each stratum. This approach ensures that each subgroup is adequately represented in the final sample, leading to more precise and reliable estimates.

Suppose we want to collect poll data for the presidential election for 250 voters. Further, suppose a recent poll found that 36% of voters are registered with the Democratic Party, 31% of voters are registered with the Republican Party, and 33% are not registered for either major party. How might we collect a stratified sample and what's the advantage compared to SRS?

Note: While there are many other more complex techniques to sample, we'll generally assume that any sample collected in this class is a simple random sample.

Data can be either collected in an ***observational study*** or in an ***experiment***.

In an ***observational study***, data is collected without manipulating the environment or influencing how the data naturally occur.

How might one collect data to investigate a connection between smoking and cancer?

In an ***experiment***, participants are assigned to different treatment groups to determine if the treatment causes changes in the outcome. A key component is the ***control group***, where participants receive no treatment or a standard one, serving as a baseline for comparison. This setup helps establish causality by isolating the effect of the treatment from other variables.

Why might it be difficult to conduct a controlled experiment investigating smoking and cancer?

Suppose we want to set up a controlled experiment to investigate the efficacy of a new blood pressure reduction drug. We begin by randomly sampling 100 people with high blood pressure. How would we set up a controlled experiment?

In addition to the sampling method, bias may still occur if participants in an experiment are aware of the treatment they are receiving. This awareness can influence their behavior or responses, potentially skewing the results.

20 patients are being treated for depression, half are given an antidepressant and half are given a sugar pill (placebo). Suppose the patients know which pill they have been given. How might this cause the results to be biased?

Blinded Experiments

A ***blinded experiment*** is an experiment in which participants do not know which treatment they are receiving.

A ***double-blinded experiment*** is an experiment in which both the participants and the researchers do not know which treatment is being administered during the experiment.

A ***triple-blinded experiment*** is an experiment in which the participants, the researchers, and the analysts evaluating the data are all unaware of which treatment is being administered. This approach further reduces the potential for bias at all stages of the experiment.

Types of Data

We will break data into two categories: **quantitative** and **qualitative**.

Quantitative data is numeric data or numbers and can broken into two further categories, ***discrete*** and ***continuous***.

Discrete data is quantitative data with a finite or countably infinite number of values.

Continuous data is quantitative data with an uncountably infinite number of values or data taken from an interval.

Give examples of discrete and continuous data.

Qualitative data refers to names, categories, or descriptions and can also be broken down into two further categories, ***nominal*** and ***ordinal***.

Nominal data is qualitative data with no natural ordering.

Ordinal data is qualitative data with a natural ordering.

Give examples of nominal and ordinal data.

Summarizing Qualitative Data

While we will mostly focus on quantitative data in this class, there are a few simple methods for describing qualitative data that you should be aware of.\

Let's collect some data on eye color from members of the class and then explore some statistical and graphical summaries.

Data:

Frequency Table

Bar Graph

Proportions: $p = \frac{\#\text{ in category}}{\#\text{ total}}$

What proportion are brown?

Percentages: $P\% = p \cdot 100$

What percentage are blue?

Summary Statistics for Quantitative Data

For the following exercises, we will use the following data set. The heights of five students are measured to be (in cm): 183, 165, 165, 175, 187.

Notation: $\sum_{i=1}^{n} x_i = x_1 + x_2 + ... + x_n$

What is $\sum x_i$, the sum of all heights in the sample?

There are many ways to measure the "center" of our data set. We will look at mean, median, and mode.

The ***sample mean*** (or ***sample average***) $\bar{x}$ is given by: $\bar{x} = \frac{\sum_{i=1}^{n} x_i}{n}$.

What is the sample mean of the original heights data set, $\bar{x}$?

The ***sample median*** $\tilde{x}$ is obtained by first ordering the n observations from smallest to largest and then calculated by the following:\

$$\tilde{x} = \begin{cases} \text{the single middle value if } n \text{ is odd,} & \tilde{x} = x_{(n+1)/2} \\ \text{the average of the two middle values if } n \text{ is even,} & \tilde{x} = (x_{n/2} + x_{n/2+1})/2 \end{cases}$$

where x_i is the ith smallest value in the data set.

What is the sample median of the heights data set, $\tilde{x}$?

If we measure an additional person at 174cm, what is the new median of the data set?

The ***sample mode*** is the most commonly occurring data point.

What is the sample mode of the heights data set?

Measures of Variability

There are a few different ways to describe the variability of a data set, including range, variance, standard deviation, and interquartile range.

The ***range*** of a data set is the difference between the largest and the smallest values.

What is the range of the heights data set?

Typically, the most common method for measuring the spread of a data set is variance or standard deviation.

The ***sample variance***, denoted by s^2, is given by:

$s^2 = \sum_{i=1}^{n} \frac{(x_i - \bar{x})^2}{n-1}$

The ***sample standard deviation***, denoted by s, is given by:

$s = \sqrt{s^2} = \sqrt{\sum_{i=1}^{n} \frac{(x_i - \bar{x})^2}{n-1}}$

Using the heights data set, calculate the sample variance and standard deviation.

i	1	2	3	4	5	Σ
x_i						
$x_i - \bar{x}$						
$(x_i - \bar{x})^2$						

The ***z-score*** measures the number of standard deviations an observation is from the mean: $z_i = \frac{x_i - \bar{x}}{s}$

A ***percentile*** is a measure of relative standing. The pth percentile is the number where at least p% of the data values are less than or equal to this number.

Special percentiles:

1. 25th percentile = 1st quartile = Q_1
2. 50th percentile = 2nd quartile = Q_2 = median
3. 75th percentile = 3rd quartile = Q_3

Computing Quartiles
1. Order the list of data values from smallest to largest
2. Compute the median. This is Q_2.
3. Split the data in half at the median. If there are an odd number of data values, the median is included in neither half.
4. Q_1 is computed by finding the median of the lower half.
5. Q_3 is computed by finding the median of the upper half.

Calculate Q_1 and Q_3 of the heights data set.

The ***interquartile range (IQR)*** is a measure of statistical dispersion, representing the difference between the first quartile (Q1) and the third quartile (Q3). It captures the range of the middle 50% of the data, indicating the spread of the central portion of the dataset. It is given by: $Q_3 - Q_1$.

Calculate IQR for the heights data set.

An ***outlier*** is a data point that lies far outside of the normal range of the rest of the data.

Rule of thumb: Data points smaller than $Q_1 - 1.5 \cdot IQR$ or larger than $Q_3 + 1.5 \cdot IQR$ are outliers.

The ***five-number summary*** is a statistical description of a dataset, consisting of the minimum, first quartile (Q1), median (Q2), third quartile (Q3), and maximum. It provides a quick overview of the data's distribution, highlighting its center and spread.

The five numbers are: minimum, Q_1, median, Q_3, maximum.

What is the five number summary for the heights data set?

Visualizing Quantitative Data


A ***histogram*** visualizes the distribution of a dataset by grouping data into bins and plotting the frequency of data points in each bin. The x-axis represents the data range, and the y-axis shows the frequency of occurrences. Histograms are useful for understanding the shape, central tendency, and spread of data.

14 students were randomly sampled and were asked how many credits they are taking this semester.

Data: 12, 14, 15, 12, 9, 18, 22, 15, 16, 14, 14, 12, 3, 17

Class	Frequency	Relative Frequency	Percentage Frequency
$[0, 3)$			
$[3, 6)$			
$[6, 9)$			
$[9, 12)$			
$[12, 15)$			
$[15, 18)$			
$[18, 21)$			
$[21, 24)$			
Total			

Histogram





```
## Loading required package: gridExtra
```

```
##
## Attaching package: 'gridExtra'
```


```
## The following object is masked from 'package:dplyr':
##
##     combine
```

Histogram shapes


1. Unimodal


IQ Test Scores
count
10000
7500
5000
2500
0
50
100
150
Score
Minutes Late for Class
count
7500
5000
2500
0
0
5
10
15
20
Minutes


2. Bimodal


Heights of American Adults
count
5000
4000
3000
2000
1000
0
55
60
65
70
75
Height (in inches)


3. Bell-shaped




4. Positively (right) skewed - tail is on the right



5. Negatively (left) skewed - tail is on the left




6. Symmetric




Box Plots

A ***box plot***, or box-and-whisker plot, visualizes data distribution using a five-number summary: minimum, first quartile, median, third quartile, and maximum [more on these later]. The box shows the interquartile range (IQR), and the whiskers extend to values within 1.5 times the IQR. Outliers are plotted as individual points. Box plots are useful for comparing distributions and spotting outliers.



Here are some examples of box plots.



Notation for Population Parameters and Sample Statistics

Some care must be given when reporting a mean, variance, or standard deviation, as it is more important to distinguish between population and sample.


	Mean	Median	Variance	Standard Deviation
Population	μ	$\tilde{\mu}$	σ^2	σ
Sample	$\bar{x}$	$\tilde{x}$	s^2	s

Descriptive Statistics vs. Inferential Statistics

After a sample is taken, there are a variety of ways to describe the sample. This is the realm of ***Descriptive Statistics***. We can describe our sample with numbers (statistics) or pictorially.

Later in the semester, we will examine ***Inferential Statistics***. In Inferential Statistics, our goal is to use characteristics of a sample to say something about the full population (see Inferential Statistics).

Connection between Probability and Statistics



If we know the characteristics of a population, the study of Probability will help us determine the likelihood that a sample will have certain characteristics.

Likewise, Statistics can potentially help us use the characteristics of a sample to say something about the population.

R Companion for Chapter 1

Example 1: Let's manually enter the heights data set (183, 170, 160, 175, 187) from chapter 1 and the following R commands will do the calculations we completed by hand.

Note: Lines of code that begin with # are comments and are not executed by the computer. Lines of code that begin with # denote the output.

```
# manually enter your data by using the "c" command
heights = c(183,165,165,175,187)

# calculate sample statistics on data set
mean(heights)
```

```
## [1] 175
```

```
median(heights)
```

```
## [1] 175
```

```
var(heights)
```


```
## [1] 102
```

```
sd(heights)
```

```
## [1] 10.0995
```

We can also create a histogram in R. Note that we can specify the breaks for the bins. In this case, the breaks are defined at 160, 170, 180, 190.

```
hist(heights,breaks=c(160,170,180,190))
```



Example 2: Let's create a randomized dataset of eye colors and analyze the data.

Note: When we generate random data in this class, we will use the **set.seed** function, so that we all generate the same random data. If we don't use the **set.seed** command before generating random data, we will all get different random data.

Here, we generate 47 random eye colors and take a look at the first few observations.

```
set.seed(2020)
eye.colors = sample(c("amber", "blue", "brown", "gray", "green",
                      "hazel", "red"), 47, replace = T)

# the "head" function will give the first few data values
# or the "header" of the dataset
head(eye.colors)
```


```
## [1] "gray"  "gray"  "red"   "hazel" "amber" "amber"
```

We can create a frequency table of these values and create a barplot.

```
table(eye.colors)
```

```
## eye.colors
## amber  blue brown  gray green hazel   red
##     3    10     3    10     7     8     6
```

```
barplot(table(eye.colors), xlab = "eye color",
        ylab = "frequency", main = "Bar plot of eye color occurance")
```



We can also calculate proportion and percentage tables.

```
table(eye.colors) # shows how many of each value exist
```

```
## eye.colors
## amber  blue brown  gray green hazel   red
##     3    10     3    10     7     8     6
```

```
length(eye.colors) # tells how many data points there are
```

```
## [1] 47
```

```
table(eye.colors)/length(eye.colors) # proportion
```

```
## eye.colors
##      amber       blue      brown       gray      green      hazel        red
## 0.06382979 0.21276596 0.06382979 0.21276596 0.14893617 0.17021277 0.12765957
```

```
table(eye.colors)/length(eye.colors)*100 # percent
```

```
## eye.colors
##     amber      blue     brown      gray     green     hazel       red
##  6.382979 21.276596  6.382979 21.276596 14.893617 17.021277 12.765957
```

Probability

In this chapter, we'll delve into Probability, the branch of mathematics that deals with the study of randomness and uncertainty, helping us understand and quantify the likelihood of different outcomes.

Sample Space and Events

An ***experiment*** is any activity or process whose outcome is subject to uncertainty.

The ***sample space*** of an experiment, denoted by Ω or $\mathcal{S}$, is the set of all possible outcomes of that experiment.

An ***event*** is any collection (subset) of outcomes contained in the sample space Ω.

For the following examples, define the sample space and give an example of an event. Code for running the experiment in R is given.

Experiment 1: Roll a fair six-sided die.

```
# roll one fair six-sided die in R
sample(x=1:6,size=1)
```

```
## [1] 5
```

Experiment 2: Roll two fair six-sided dice.

```
# roll two fair six-sided dice in R
sample(x=1:6,size=2)
```

```
## [1] 1 3
```

Experiment 3: Randomly select a real number in the closed interval from 0 to 1.

```
# generate one random number between 0 and 1
runif(n=1)
```

```
## [1] 0.6137196
```


Set Theory

Note: For the following examples, suppose a fair six-sided die is rolled, so $\Omega = \{1, 2, 3, 4, 5, 6\}$.

Define the following events: A = roll an even $= \{2, 4, 6\}$, B = roll a 3 or higher $= \{3, 4, 5, 6\}$, C = roll a prime number $= \{2, 3, 5\}$.


A ***complement*** of an event A, denoted by A^c or A', is the set of all outcomes in Ω that are not contained in A.

Illustration




The ***union*** of two events A and B, denoted by $A \cup B$ and read "A or B", is the event consisting of all outcomes that are either in A or B or in both.

Illustration




The ***intersection*** of two events A and B, denoted by $A \cap B$ and read "A and B", is the event consisting of all outcomes that are in both A and B.

Illustration



The ***set difference*** of two events A and B, denoted by A\B and read "difference of A and B", is the event consisting of all outcomes that are in A but not in B.

Illustration



The ***null or empty set*** is the set of no events (or elements), denoted $\emptyset$.


When $A \cap B = \emptyset$, A and B are said to be ***mutually exclusive (or disjoint)*** events.

Suppose a fair six-sided die is rolled, so $\Omega = \{1, 2, 3, 4, 5, 6\}$.


Define the following events: A = roll an even $= \{2, 4, 6\}$, B = roll a 3 or higher $= \{3, 4, 5, 6\}$, C = roll a prime number $= \{2, 3, 5\}$.

Determine the following sets and fill in the Venn diagrams representing these sets.


Draw and determine $A \cup B^c$.




Draw and determine $A \cap B \cap C^C$.



Draw and determine $[A \cup B] \cap C$.



Draw and determine $A \cap B \cap C$.



Axioms and Rules of Probability

The probability of an event, A, is denoted by $P(A)$ and represents the likelihood or chance that the event A will occur. Note that $0 \leq P(A) \leq 1$.

Axioms of Probability

The axioms of probability were introduced by the mathematician Andrey Kolmogorov in 1933. These axioms form the foundation of probability theory, providing a rigorous, axiomatic framework for quantifying uncertainty. In mathematics, an axiomatic approach means that all rules and theorems are built from a set of fundamental principles, or axioms.

Axiom 1: For any event A, $P(A) \geq 0$

Axiom 2: $P(\Omega) = 1$

Axiom 3: If $A_1, A_2, A_3, \ldots$ is an infinite collection of disjoint events, then

$$P(\cup_{i=1}^{\infty} A_i) = P(A_1 \cup A_2 \cup A_3 \cup \ldots) = \sum_{i=1}^{\infty} P(A_i)$$

Probability Rules

The following can be derived from the Axioms of Probability.

1. $P(\emptyset) = 0$

2. For any event A, $P(A) + P(A^c) = 1$

3. $P(A^c) = 1 - P(A)$

4. For any event A, $0 \leq P(A) \leq 1$

5. For any two events A and B, $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

6. For any three events A, B, and C,

$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$.

7. $[A \cup B]^C = A^C \cap B^C$

8. $[A \cap B]^C = A^C \cup B^C$

Suppose our class has the following breakdown:

70% Math students
20% seniors
30% Born in Denver
10% Math students and seniors
15% Math students and born in Denver
10% Seniors and Born in Denver
5% Math students, seniors, and born in Denver

Define events as follows: A = Math student, B = senior, C = Born in Denver.

Suppose a random student from the class is selected. Calculate the following probabilities.

(a) $P(A \cup B)$

(b) $P(C \backslash B)$

(c) $P([A \cap C]^c)$

(d) $P(A \cup B \cup C)$

Counting Techniques

Equally Likely Outcomes

If $\Omega = \{x_1, x_2, ...x_N\}$ and all outcomes are equally likely, then $P(A) = |A|/N$, where $|A|$ is the number of outcomes contained in the event A.

You roll a fair six-sided die. Calculate $P(even)$.

(Product Rule for Ordered Pairs) If the first element of an ordered pair can be selected in n_1 ways and for each of the these n_1 ways the second element of the pair can be selected in n_2 ways, then the number of pairs is $n_1 \cdot n_2$.

You flip two fair coins. Calculate the probability that all are heads or all are tails.

(Generalized Product Rule) Suppose a set consists of k elements (k-tuples) and that there are n_1 possible choices for the first element, n_2 possible choices for the second element, ... , and n_k possible choices for the kth element, then there are $n_1 \cdot n_2 \cdot \cdots n_k$ possible k-tuples.

Quartiles is a word game where one of the goals is to find all five words that contain four tiles each. Suppose we have found 3 of the quartiles. How many total attempts are there to find the fourth quartile?



The factorial function, $n!$, is important in counting methods and probability. It is the product of all positive integers less than or equal to n, that is, $n! = n \cdot (n-1) \cdot (n-2) \cdot \ldots \cdot 2 \cdot 1$ for $n \in \mathbb{Z}^+$.

Note: $0! = 1$

An ordered subset is called a ***permutation***. The number of permutations of size k that can be formed from the n elements in a set will be denoted by $P_{n,k}$.

$$\boxed{P_{n,k} = \frac{n!}{(n-k)!}}$$

An unordered subset is called a ***combination***. The number of combinations of size k that can be formed from the n elements in a set will be denoted by $\binom{n}{k}$ or $C_{n,k}$.

$$\boxed{\binom{n}{k} = C_{n,k} = \frac{n!}{k! \cdot (n-k)!}}$$

Suppose there are five competitors in an Olympic competition. A gold, a silver, and a bronze medal are awarded for the top three competitors. Are combinations or permutations more appropriate for this situation? How many possible ways can these three awards be given out?

Suppose there are five competitors in an Olympic competition and only three can advance to the final competition. Are combinations or permutations more appropriate for this situation? How many possible ways can you choose three people from a group of five?

Conditional Probability

Conditional probability is the likelihood of an event occurring given that another event has already occurred. It refines our understanding of probability by taking into account additional information that might affect the outcome. Understanding conditional probabilities is crucial in many areas of statistics and decision-making, as it allows us to update our predictions and assessments based on new evidence.

For any two events A and B with $P(B) > 0$, the conditional probability of A given B is defined by:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

The following is a contingency table from a survey on congressional approval.

Political Affiliation	Approves of Congress	Disapproves of Congress	Total
Republican	100	20	120
Democrat	20	80	100
Independent	40	40	80
Total	160	140	300

(a) What is the probability that a randomly selected individual from the survey approves of Congress?

(b) What is the probability that a randomly selected individual approves of Congress given that they are Democrat?

(c) Given that the randomly selected person approves of the Congress, what is the probability that they are Republican?


Multiplication Rule: $P(A \cap B) = P(A|B) \cdot P(B)$

Suppose that people with a genetic variation have a 50% chance of getting a rare disease. The probability of having the genetic variation is 1%. What is the probability that a randomly chosen person has the genetic variation and also has the disease?

Law of Total Probability: Let $A_1, A_2, \ldots, A_k$ be mutually exclusive and exhaustive events (i.e., the events form a partition). Then for any event B,

$$P(B) = P(B|A_1) \cdot P(A_1) + P(B|A_2) \cdot P(A_2) + \ldots + P(B|A_k) \cdot P(A_k)$$

Illustration of a Partition



Box #1 has ten computer chips (five working and five broken). Box #2 has five (one working and four broken). Suppose we randomly select one of the two boxes and then randomly select a chip. What is the probability that the chip is broken?

Bayes' Theorem: Suppose that $A_1 \cap A_2 = \emptyset$ and $A_1 \cup A_2 = \Omega$, then if $P(B) > 0$,

$$P(A_1|B) = \frac{P(B|A_1)P(A_1)}{P(B|A_1)P(A_1) + P(B|A_2)P(A_2)}$$

Let $A_1, A_2, \ldots, A_k$ be a collection of mutually exclusive and exhaustive events, then if $P(B) > 0$,

$$P(A_j|B) = \frac{P(A_j \cap B)}{P(B)} = \frac{P(B|A_j) \cdot P(A_j)}{\sum_{i=1}^{k} P(B|A_i) \cdot P(A_i)}$$

Suppose a test to detect a rare disease is positive 95% of the time for a person with a disease and is positive 1% of the time for a person without the disease. Further, assume that 0.5% of the population has the disease. Given that a person tests positive for a disease, what is the probability that they actually has the disease?

Independence

Recall:

When $A \cap B = \emptyset$, A and B are said to be ***mutually exclusive (or disjoint)*** events.

Two events A and B are ***independent*** if $P(A|B) = P(A)$ and ***dependent*** otherwise.

Also, A and B are independent if and only if $P(A \cap B) = P(A) \cdot P(B)$.

Recall from the previous example that a test to detect a rare disease is positive 95% of the time for a person with a disease and is positive 1% of the time for a person without the disease. Further, assume that 0.5% of the population has the disease. Let D be the event that one has the disease and let P be the event that one has a positive test.

(a) Are D and P mutually exclusive?

(b) Are D and P independent?

Objective vs. Subjective Probability

There are two different interpretations of Probability. Both follow the Axioms of Probability.

Objective Probability refers to the probability of an event as determined by the long-term frequency of its occurrence, based on repeated trials or observations. It is rooted in empirical data and reflects the likelihood of an event as the proportion of times it occurs in a large number of trials.

I flip a coin 100 times and it lands on head 53 times. I estimate the probability of flipping a heads to be $P_{100}(head) = \frac{53}{100} = 0.53$. As we flip the coin more and more, we would expect that $P_n(head) \to 0.5$ as $n \to \infty$ if the coin is actually a fair coin.

Subjective Probability refers to the probability of an event as determined by an individual's personal judgment or belief, rather than objective data. It reflects the degree of confidence one has in the occurrence of an event. A well-known approach to subjective probability is Bayesian probability, which updates these beliefs based on new evidence.

I am given a coin which I have no reason to believe it is unfair (i.e, $P(head) = 0.5$). I flip the coin 100 times and 3 land on heads. Since this is very unlikely to happen if the coin is fair, I update my estimate to be $P(head) = 0.1$ (i.e., somewhere in between $3/100 = 0.03$ and $1/2 = 0.5$).

R Companion for Chapter 2


Let's simulate a fair six-sided die roll. Recall that we need to use the **set.seed** function so that we all generate the same random data. For consistency, I always set the seed to 2020, i.e., set.seed(2020). The sample function will generate 100 random die rolls with each possible outcome (1,2,3,4,5,6) being equally likely.

```
set.seed(2020)
data = sample(x = 1:6,size = 100,replace = T); data
```

```
##   [1] 4 4 6 1 1 4 2 6 1 5 2 2 6 5 2 3 2 5 4 2 6 6 4 6 4 2 4 5 4 4 3 6 2 2 6 3 5 4 5 5 2 5 1
##  [52] 3 2 6 2 1 3 2 1 6 5 5 2 5 6 4 3 6 4 4 2 2 1 6 1 4 6 2 2 2 1 4 1 1 4 2 2 5 5 4 4 4 6 5
```

Let's make a histogram of the simulated data. Recall that each outcome (1,2,3,4,5,6) is equally likely in theory but in our simulation, some numbers may come up more than others. To make the histogram look better, I define the boundary breaks to be 0:6 (i.e., 0,1,2,3,4,5,6).

```
hist(data,breaks=0:6)
```




With only 100 random die rolls, the distribution isn't as flat, uniform, and symmetric as we would expect. The number 3 came up less than we would anticipate. This is OK.

We can also generate a frequency table and use this to create a barplot. This looks a little nicer than the standard histogram and doesn't require use to use the **breaks** argument.

```
table(data)
```

```
## data
##  1  2  3  4  5  6
## 15 21  8 21 18 17
```

```
barplot(table(data))
```




If we want a better representation of the true distribution of the fair six-sided die rolls, we could simulate 10000 die rolls instead of 100.

```
set.seed(2020)
data = sample(x = 1:6,size = 10000,replace = T)
table(data)
```

```
## data
##    1    2    3    4    5    6
## 1630 1657 1733 1655 1629 1696
```

```
barplot(table(data))
```



Simulation can be helpful for us to verify hand calculations and to estimate quantities that are hard to calculate by hand. Our simulation results won't exactly match our hand calculations, but if the sample size is large enough, they should be close.\

For instance, we know that P(Roll=1) = 1/6 for the population of all fair six-sided die rolls. If you refer to the table at the bottom of the previous page, you see that 1 came up 1630 times out of 10000 die rolls. In the simulation, P(1) = 1630/10000 = 0.1630 which is pretty close the the theoretical value of P(1) = 1/6 = 0.1667.\

Here is an example of how to calculate the probability of a die roll less than 3 in our simulation.

```
# Calculate P(Roll < 3)
# From theory, we know that P(Roll < 3) = P(1) + P(2) = 1/6 + 1/6 = 1/3

# count up the number of rolls less than 3
num.under3 = sum( data < 3 )

# divide the number of rolls less than 3 by the number of rolls
prob.under3 = num.under3 / 10000
prob.under3
```

```
## [1] 0.3287
```

```
# alternatively:
prob.under3 = mean(data < 3)
prob.under3
```

```
## [1] 0.3287
```

From theory, we expect P(Roll < 3) $= 1/3 = 0.3333$. In our simulation, we found P(Roll < 3) $=$ 0.3287.

Let's simulate rolling two fair six-sided dice 10,000 times and calculate some probabilities. **d1** will represent the first die roll and **d2** will represent the second die roll.

```
set.seed(2020)
d1 = sample(x = 1:6,size = 10000,replace = T)
d2 = sample(x = 1:6,size = 10000,replace = T)
```

From our simulated data, let's calculate P(D1 = 1 and D2 = 2), that is, the probability that first die roll is 1 and the second die roll is 2. From theory, this probability should be $1/36 = 0.0278$. (We use the **&** for **AND** in R.)

```
num.events = sum(d1 == 1 & d2 == 2)
prob = num.events / 10000; prob
```

```
## [1] 0.0256
```

Next, let's calculate P(D1 < 2 or D2 > 2), that is, the probability that the first die roll is less than 2 or the second die roll is greater than 2. From theory, this probability should be 0.7222. We use the | for the **OR** in R.)

```
num.events = sum(d1 < 2 | d2 > 2)
prob = num.events / 10000; prob
```


```
## [1] 0.7183
```

Finally, let's look at the frequency table and distribution of the sum of the two die rolls.

```
table(d1+d2)
```

```
##
##    2    3    4    5    6    7    8    9   10   11   12
##  271  540  824 1154 1373 1703 1360 1149  814  525  287
```

```
barplot(table(d1+d2),xlab="Sum of Dice",ylab="Counts")
```


Counts
0
500
1000
1500
2
3
4
5
6
7
8
9
10
11
12
Sum of Dice

Discrete Random Variables

In this chapter, we'll explore random variables and their associated probability distributions, with a focus on discrete random variables. These concepts are the backbone of probability theory, enabling us to model and analyze uncertainty. We'll also briefly cover the differences between discrete and continuous random variables—discrete variables take on specific, countable values, while continuous variables can assume any value within a range.

Random Variables

For a given sample space Ω of an experiment, a ***random variable (RV)*** is a rule that assigns a number to each outcome in Ω. Mathematically, a random variable is a function with the sample space as its domain and the set of real numbers as its range.

A ***discrete random variable*** is a random variable whose possible values either constitute a finite set or a countably infinite set.

A ***continuous*** random variable is a random variable whose possible values are uncountably infinite or defined over an interval. Note: $P(X = c) = 0$ for a continuous random variable X and a constant c.

For the following cases, determine whether the random variable is discrete or continuous and state the sample space, Ω.

(a) Roll a fair six-sided die. Let X be the number on the top of the die.

(b) Pick a random person and let Y be the number of keys in their pocket.

(c) Suppose a student is given 50 minutes to complete an exam. Let Z be the amount of time it takes the student to complete the exam.

Probability Distributions for Discrete Random Variables

The ***probability mass function (pmf)*** or ***probability distribution*** of a discrete random variable X is defined for every number x by $p(x) = P(X = x)$.

Note: $\sum_{x \in X} p(x) = 1$

The ***cumulative distribution function (CDF)*** $F(x)$ of a discrete random variable X with pmf $p(x)$ is defined for every number x by: $F(x) = P(X \leq x) = \sum_{y:y \leq x} p(y)$

- $F(x)$ gives you the probability that a random variable will be at most equal to x.
- $F(x)$ must be right-continuous and non-decreasing.
- $F(x)$ must satisfy two limit laws:
  - (i) $\lim_{x \to -\infty} F(x) = 0$
  - (ii) $\lim_{x \to \infty} F(x) = 1$


For any two numbers a and b with $a \leq b$, $P(a \leq X \leq b) = F(b) - F(a^-)$.

Roll a fair six-sided die. Let X be the number of the top of the die. Find the pmf and cdf of X and sketch them.





Flip two fair coins. Let Y be the number of coins that land on heads. Find the pmf and cdf of Y and sketch both.



Expected Values (Means)

Let X be a discrete random variable with set of possible outcomes D and pmf $p(x)$. The ***expected value*** or ***mean value*** of X, denoted $E[X]$ or μ, is:

$$E[X] = \sum_{x \in D} x \cdot p(x)$$

$E[h(X)] = \sum p(x) \cdot h(x)$

$E[aX + b] = a \cdot E[X] + b$

Let X have a pmf $p(x)$ and expected value μ. Then the ***variance*** of X, denoted by $Var(X)$ or σ^2, is:

$$Var(X) = \sigma^2 = \sum_{x \in D} (x - \mu)^2 \cdot p(x) = E[(X - \mu)^2]$$

The ***standard deviation***, denoted SD or σ is:

$$SD(X) = \sigma = \sqrt{Var(X)}$$

$Var(X) = E[X^2] - (E[X])^2$

$Var(aX + b) = a^2 \cdot \sigma^2$

Roll a fair six-sided die. Let X be the number of the top of the die. Find $E[X]$ and $Var(X)$.

x	1	2	3	4	5	6	Σ
$p(x)$							
$x * p(x)$							
$x^2 * p(x)$							

Flip two fair coins. Let Y be the number of coins that land on heads. Find $E[Y]$ and $Var(Y)$.

y	0	1	2	Σ
$p(y)$				
$y * p(y)$				
$y^2 * p(y)$				

Suppose $Z = 3 \cdot Y + 2$, where Y is defined above. Find $E[Z]$ and $Var(Z)$.

Bernoulli and Binomial Random Variables

Suppose $p(x)$ depends on a quantity that can be assigned any one of a number of possible values, with each different value determining a different probability distribution. Such a quantity is called a ***parameter*** of the distribution.

The collection of all probability distributions for different values of the parameter is called a ***family*** of probability distributions.

A ***Bernoulli(p)*** random variable X is a discrete random variable with two possible outcomes (typically, these outcomes are 0 and 1).

The PMF of a Bernoulli(p) is:

$$p(x) = \begin{cases} 1-p & x = 0 \\ p & x = 1 \end{cases}$$

If $X \sim$ Bernoulli(p)) random variable, then $E[X] = p$ and $Var(X) = p(1-p)$

Show that if X is a Bernoulli(p) random variable, then $E[X] = p$ and $Var(X) = p(1-p)$.

X is a ***Binomial(n,p)*** random variable if X is a discrete random variable that satisfies the following conditions:

- The experiment consists of n Bernoulli trials, where n is fixed.
- The trials are independent.
- The probability of success, p, is constant from trial to trial.
- X is the number of successes

The PMF of a Binomial(n, p) is:


$p(x) = \binom{n}{x} p^x (1-p)^{n-x}, x = 0, 1, 2, \ldots, n$

If $X \sim$ Binomial(n, p) random variable, then $E[X] = np$ and $Var(X) = np(1-p)$.

Basketball player Lebron James has a career free throw percentage of 73.1% (i.e., there's a 73.1% chance he will make a basket from the free throw line). Suppose Lebron has six free throw attempts in a game and assume all free throw shots are independent. Answer the following questions.

(a) Let X be the number of free throws that Lebron makes. Find and sketch the pmf of X.

x	0	1	2	3	4	5	6
$p(x)$							



(a) On average, how many free throws is Lebron expected to make? What is the variance?

(c) What is the probability that Lebron makes at least five shots?

Geometric, Discrete Uniform, and Poisson Random Variables

X is a ***Geometric(p)*** random variable is X is a discrete random variable with the following properties.

- The experiment consists of independent trials.
- Each trial can result in a success or a failure.
- The probability of success is p and is constant across all trials.
- The experiment continues until a successful trial is observed.
- X is the number of total trials

The PMF of a Geometric(p) is:

$p(x) = p(1-p)^{x-1}, x = 1, 2, 3, \ldots$

If $X \sim$ Geometric(p) random variable, then $E[X] = \frac{1}{p}$ and $Var(X) = \frac{1-p}{p^2}$.

Basketball player Lebron James has a career free throw percentage of 73.1%. Suppose he takes shots until makes one. Calculate the probability that 3 or more shots will be required to make his first shot.

X is a ***Discrete Uniform(a,b)*** random variable is X is a discrete random variable such that the outcomes $a, a+1, \ldots, b$ are equally likely. Let $n = b - a + 1$.

The PMF of a Uniform(a, b) is:

$p(x) = \frac{1}{n}, x = a, a+1, \ldots, b$

If $X \sim$ Uniform(a, b) random variable, then $E[X] = \frac{a+b}{2}$ and $Var(X) = \frac{(b-a+1)^2-1}{12}$.

Using the formulas above, calculate the expected value and variance of a fair six-sided die roll.

A ***Poisson random variable*** is a discrete random variable with parameter λ and the following pmf:

$$p(x) = \frac{e^{-\lambda}\lambda^x}{x!}, x = 0, 1, 2, \ldots$$

If $X \sim$ Poisson(λ) random variable, then $E[X] = \lambda$ and $Var(X) = \lambda$.

Note: Poisson random variables are often used to model the number of events that occur in a finite period of time.

Jihan is going fishing and on average, she catches 2 fish per day. Assume that the time between successive fish caught is independent. What is the probability that Jihan catches less than 3 fish in a given day?

R Companion for Chapter 3

We can calculate probabilities for common discrete random variables easily in R. There are two parts to finding probabilities using the PMF or CDF.

R uses four prefixes to reference difference elements of a random variable. These are:

- ***p*** for "probability", the cumulative distribution function (CDF)
- ***q*** for "quantile", the inverse CDF
- ***d*** for "density", the probability mass function (PMF)
- ***r*** for "random", a random variable having the specified distribution

In addition, we have suffixes for common random variables: ***binom*** (binomial), ***pois*** (Poisson), and ***geom*** (Geometric). Note that ***geom*** is defined in terms of failures instead of total trials.

For instance, suppose that $X \sim Binomial(n = 10, p = 0.5)$, that is, X is a binomial random variable that counts the number of heads in 10 fair coin flips. We can calculate P(X=5), the probability of exactly 5 heads in 10 fair coin flips, as follows.

```
dbinom(x=5,size=10,prob=0.5)
```

```
## [1] 0.2460938
```

If we want the probability of at most five heads in 10 coin flips, $P(X \leq 5)$, we can use ***pbinom}.

```
pbinom(q=5,size=10,prob=0.5)
```

```
## [1] 0.6230469
```

Now, suppose that $Y \sim Poisson(\lambda = 2)$ and we want to know $P(Y > 3)$. First, we note that $P(Y > 3) = 1 - P(Y \leq 3)$.

```
1-ppois(q=3,lambda=2)
```

```
## [1] 0.1428765
```

If we use the prefix ***r***, we can generate simulated data according to a particular random variable. Let's generate 10,000 observations from a Poisson distribution with $\lambda = 3$, look at the first few values, and then calculate some summary statistics. This will require us to use the ***rpois*** function to generate random Poisson data.

```
rand.data = rpois(n=10000,lambda=3)
head(rand.data) # look at the first few values
```

```
## [1] 3 6 2 4 1 4
```

```
mean(rand.data) # calculate the mean
```

```
## [1] 3.0088
```

```
var(rand.data) # calculate the variance
```


```
## [1] 3.047827
```

From theory, we know that if $Y \sim Poisson(\lambda = 3)$, then $E[Y] = Var(Y) = \lambda = 3$. Our simulated results are pretty close to the theoretical values.

```
table(rand.data)
```

```
## rand.data
##    0    1    2    3    4    5    6    7    8    9   10   11
##  520 1458 2252 2213 1676 1031  514  211   79   28   15    3
```

```
barplot(table(rand.data))
```




How would you describe the distribution above? It seems to be asymmetric, skewed right (tail on the right), and unimodal.

Let's simulate some data and calculate the probabilities in the sample. Suppose we flip 10 fair coins and we count the number of heads. Let X be the number of heads. We can model X as a binomial random variable with parameters n=10 (number of coin flips) and p=0.5 (fair coin). Let's simulate running this experiment 100 times.

```
set.seed(2020)
coins = rbinom(n=100,size=10,prob=0.5)
table(coins)
```

```
## coins
##  1  2  3  4  5  6  7  8  9
##  3  1 12 19 32 19  8  5  1
```

```
barplot(table(coins))
```



In our simulation, what was the probability that 5 heads occurred out of 10 flips? In the above table, we can see we got 5 heads on 32 of the 100 simulations, so $P(5) = 32/100 = 0.32$.\

We could also calculate as follows:

```
p5 = sum(coins == 5)/100; p5
```

```
## [1] 0.32
```

What is the probability that we got more than 7 heads?

```
p = sum(coins > 7)/100; p
```

```
## [1] 0.06
```

Continuous Random Variables

In this chapter, we will explore continuous random variables, which are types of random variables that can take any value within an uncountable set, such as an interval or a combination of intervals. We will also delve into some of the most commonly encountered continuous random variables, with a particular focus on the Normal distribution.

A ***continuous random variable*** is a random variable whose possible values either constitute an interval of real numbers or a union of intervals of real numbers.

Probability Density Function (pdf)

Let X be a continuous random variable. The ***probability distribution*** or ***probability density function (pdf)*** of X is a function $f(x)$ such that for any two numbers a and b with $a \leq b$, $P(a \leq X \leq b) = \int_a^b f(x)dx$.


Note: All valid pdfs must satisfy two conditions:
(1) $f(x) \geq 0$ ($f(x)$ is non-negative).
(2) $\int_{-\infty}^{\infty} f(x)dx = 1$ (area under $f(x)$ is equal to 1)

Suppose X is a random variable with the following pdf:

$$f(x) = \begin{cases} 0 & x < 0 \\ \frac{x}{2} & 0 \leq x \leq 2 \\ 0 & x > 2 \end{cases}$$

(a) Is $f(x)$ a valid pdf?

(b) What is $P(X < 1)$?



(c) What is $P(1/2 \leq X \leq 3/2)$?



(d) What is $P(X = 1)$?

Caution! For a continuous random variable X, $P(X = c) = 0$ for all constants c. While it may be possible for X to be exactly equal to the constant c, it can only happen with probability zero. In general, when dealing with continuous random variables, we consider the probability that X takes on a value over an interval rather than a specific value.\

Cumulative Distribution Functions

The ***cumulative distribution function (CDF)*** of a continuous random variable X is defined for every number x by $F(x) = P(X \leq x) = \int_{-\infty}^{x} f(y)dy$. F(x) represents the area under the curve to the left of x.

Note: Conditions on $F(x)$:

- $0 \leq F(x) \leq 1$
- $\lim_{x \to -\infty} F(x) = 0$
- $\lim_{x \to \infty} F(x) = 1$

If X is a continuous random variable with pdf $f(x)$ and CDF $F(x)$, then at every x at which the derivative $F'(x)$ exists, $F'(x) = f(x)$.


Let X be a continuous random variable with pdf $f(x)$ and CDF $F(x)$. Then we have the following:

- For any number a, $P(X > a) = 1 - F(a)$
- For any two numbers a and b with $a \leq b$, $P(a \leq X \leq b) = F(b) - F(a)$.

Suppose X is a random variable with the following pdf:

$$f(x) = \begin{cases} 0 & x < 0 \\ \frac{x}{2} & 0 \leq x \leq 2 \\ 0 & x > 2 \end{cases}$$

(a) Find the CDF of X and sketch it.



(b) Use the CDF to calculate $P(X \leq 1)$.

Expected Values

The ***expected value*** or ***mean*** of a continuous random variable X with pdf $f(x)$ is:

$$E[X] = \mu = \int_{-\infty}^{\infty} x \cdot f(x)dx$$

If X is a continuous random variable with pdf $f(x)$ and $h(X)$ is any function of X, then:

$$E[h(X)] = \int_{-\infty}^{\infty} h(x) \cdot f(x)dx$$

The ***variance*** of a continuous random variable X with pdf $f(x)$ is:

$$Var(X) = \sigma^2 = \int_{-\infty}^{\infty} (x-\mu)^2 \cdot f(x)dx = E[(X-\mu)^2] = E[X^2] - (E[X])^2$$

The ***standard deviation*** of a continuous random variable X is:

$$SD(X) = \sigma = \sqrt{Var(X)}$$

If X is a continuous random variable and $Y = aX + b$, then:

$$E[Y] = aE[X] + b \text{ and } Var(Y) = a^2 Var(X)$$

Suppose X is a random variable with the following pdf:

$$f(x) = \begin{cases} 0 & x < 0 \\ \frac{x}{2} & 0 \leq x \leq 2 \\ 0 & x > 2 \end{cases}$$


Determine $E[X]$, $E[X^2]$, $Var(X)$, $SD(X)$.

Normal Distribution

A continuous random variable X is said to be ***Normal(μ,σ^2)*** or $\mathcal{N}(\mu, \sigma^2)$ random variable if the pdf of X is:

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/(2\sigma^2)}, -\infty < x < \infty$$

If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $E[X] = \mu$ and $Var(X) = \sigma^2$, that is, the expected value is μ and the variance is σ^2.



The normal distribution with parameter values $\mu = 0$ and $\sigma = 1$ is called the ***standard normal distribution***.

Note: $X \sim \mathcal{N}(\mu, \sigma^2)$, F(x) is denoted $\Phi(x)$. There is no closed form solution for $\Phi(x)$ since $\Phi(x) = F(x) = \int_{-\infty}^{x} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/(2\sigma^2)} = (?)$. Instead, we use software to assist in these calculations.

If $X \sim \mathcal{N}(\mu, \sigma^2)$, that is, X is a normal distribution with mean μ and standard deviation σ, then $Z = \frac{X-\mu}{\sigma}$ has a standard normal distribution.

If $Z \sim \mathcal{N}(0, 1)$, that is, X is a standard normal distribution, then $X = \sigma \cdot Z + \mu$ is a normal distribution with mean μ and standard deviation σ.


The ***Empirical Rule*** states that if X is (approximately) normally distributed then:

- Approximately 68% of the values are within 1 SD of the mean.
- Approximately 95% of the values are within 2 SD of the mean.
- Approximately 99.7% of the values are within 3 SD of the mean.


Suppose the weights of house cats are approximately normally distributed with mean of 10 pounds and a standard deviation of 3 pounds. Between what two bounds do approximately 95% of house cat weights lie?

Suppose $Z \sim \mathcal{N}(0, 1)$, that is, Z is a standard normal distribution. Calculate the following and draw an accompanying picture:


(a) $P(Z < 0.68)$




(b) $P(Z \geq -1.21)$



(c) $P(0.43 \leq Z < 0.68)$




(d) Suppose $P(Z < c) = 0.89$. Find c.




Suppose the weights of house cats are approximately normally distributed with mean of 10 pounds and a standard deviation of 3 pounds. Calculate the following and draw an accompanying picture.


(a) $P(X < 15)$



(b) What is the probability that a randomly chosen house cat will weigh between 8 and 12 pounds?



(c) What is the 99th percentile of house cat weights?



The Exponential and Uniform Random Variables

A continuous random variable X is said to be an ***Exponential(λ)*** random variable or an exponential random variable with parameter $\lambda > 0$ if the pmf of X is:

$$f(x|\lambda) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & otherwise \end{cases}$$

If $X \sim exp(\lambda)$, then $E[X] = \frac{1}{\lambda}$ and $Var(X) = \frac{1}{\lambda^2}$.

Suppose $X \sim exp(\lambda)$, find the CDF of X.

Suppose that $X \sim exp(\lambda = 2)$. Find $P(X > 2)$.

(Continuous) Uniform[a,b] Random Variable


A continuous random variable X is said to be ***Uniform[a,b]*** random variable if the pdf of X is:

$$f(x|a,b) = \begin{cases} \frac{1}{b-a} & a \leq x \leq b \\ 0 & otherwise \end{cases}$$

If $X \sim$ Uniform(a,b) random variable, then $E[X] = \frac{a+b}{2}$ and $Var(X) = \frac{(b-a)^2}{12}$.

Show that if $X \sim Uniform[a,b]$, then $E[X] = \frac{a+b}{2}$.

Suppose your local bus shows up once every 60 minutes, but you don't have a schedule, so you assume it's equally likely to show up at any time in the next 60 minutes. Let X be the amount of time you have to wait for the bus. Sketch the pmf and calculate the probability that your bus shows up in the next 10 minutes.



R Companion for Chapter 4

Similar to last chapter, we can calculate probabilities for common continuous random variables in R and generate random data according to continuous distributions.

R uses four prefixes to reference difference elements of a random variable. These are:

p for "probability", the cumulative distribution function (CDF) **q** for "quantile", the inverse CDF **d** for "density", the probability mass function (PMF) **r** for "random", a random variable having the specified distribution

Suffixes for continuous random variables include: **norm** (normal/Gaussian), **exp** (exponential), and **t** (t).

Suppose $X \sim \mathcal{N}(\mu = 100, \sigma = 15)$, the distribution often used to model IQ scores. Suppose we want to know the probability of a randomly chosen IQ score below 110.

```
pnorm(q=110,mean=100,sd=15)
```

```
## [1] 0.7475075
```

In other words, about 75% of IQ scores are less than 110.\

Let's calculate the probability that a randomly chosen score is above 130. Note that $P(X > 130) = 1 - P(X \leq 130)$.

```
1-pnorm(q=130,mean=100,sd=15)
```

```
## [1] 0.02275013
```

In other words, about 2.3% of IQ scores are greater than 130.\

We can also do inverse normal calculations using the prefix **q** which stands for quantile. Suppose we want to know the 90th percentile of IQ scores. We can calculate this using the **qnorm** function.


```
qnorm(p=0.9,mean=100,sd=15)
```

```
## [1] 119.2233
```

This result shows that the 90th percentile of IQ scores corresponds to an approximate IQ of 119.

In the second half of the course, we will be using the **t-distribution** frequently when dealing with statistical applications. Let's generate 10,000 random values for a t-distribution with 20 degrees of freedom, i.e., $Z \sim t(df = 20)$.

```
set.seed(2020)
rand.data = rt(n=10000,df=20)
hist(rand.data)
```



Based on the plot above, this distribution is approximately symmetric, bell-shaped, and unimodal.\

Let's calculate some summary statistics for this data.

```
mean(rand.data)
```

```
## [1] -0.009077007
```

```
median(rand.data)
```

```
## [1] -0.01514699
```

The mean and median are both approximately equal to zero and since these values are nearly equal, we can confirm that the distribution is approximately symmetric.

Finally, let's calculate the probability in our simulated data so that $P(Z > 1)$.

```
num.events = sum(rand.data > 1)
prob = num.events / 10000; prob
```

```
## [1] 0.1623
```

Approximately 16% of our randomly generated t-distribution values are greater than 1.

Multivariate Random Variables

These notes will explore pairs of random variables that may be related. To understand the relationship between these variables, we will start by defining joint probability distributions. Following that, we will examine the concepts of covariance and correlation to quantify and describe the nature of the relationship between the two variables.

Joint Probability Distributions

Let X and Y be two discrete random variables defined on the sample space Ω of an experiment. The ***joint probability mass function***, $p(x, y)$, is defined for each pair of numbers (x,y) by:

$$p(x, y) = P(X = x \text{ and } Y = y)$$

Note: $p(x, y) \geq 0$ and $\sum_x \sum_y p(x, y) = 1$

Let X and Y be two continuous random variables defined on the sample space Ω of an experiment. The ***joint probability density function*** $f(x, y)$ for variables X and Y is a function satisfying:

1. $f(x, y) \geq 0$
2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dy dx = 1$

Note: For any two-dimensional set A, $P((X, Y) \in A) = \int \int_A f(x, y) dy dx$.

Note: If A is a two-dimensional rectangle

$\{(x, y) : a \leq x \leq b, c \leq y \leq d\}$, then: $P((X, Y) \in A) = \int_a^b \int_c^d f(x, y) dy dx$

Let $f(x, y) = cxy, 0 \leq x \leq 1, 0 \leq y \leq 1$. Find c so that $f(x, y)$ is a valid joint pdf.

Marginal Probability Functions

The ***marginal probability mass function*** of a discrete random variable X with joint pmf $p(x, y)$ is denoted by $p_X(x)$ and is given by: $p_X(x) = \sum_y p(x, y)$

The ***marginal probability density function*** of a continuous random variable X with joint pdf $f(x, y)$ is denoted by $f_X(x)$ and is given by: $f_X(x) = \int_y f(x, y)dy$

An apartment complex has the following joint distribution of X bedrooms and Y bathrooms. Determine the marginal pmfs and calculate the probability a randomly selected apartment has more bedrooms than bathrooms (i.e., $P(X > Y)$).

Y | X	1	2	3	Σ
1	0.3	0.3	0	
2	0	0.3	0.1	
Σ				

Let X and Y have the joint pdf, $f(x, y)$, as follows.

$$f(x,y) = \begin{cases} \frac{4}{3}(xy + x) & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases} \text{ :::}$$

(a) Confirm $f(x, y)$ is a valid pdf
(b) Determine the marginal pdfs
(c) Calculate $P(X < 1/2, Y < 1/2)$

Independent Random Variables

Two discrete random variables are ***independent*** if $p(x, y) = p_X(x) \cdot p_Y(y)$.

Two continuous random variables are ***independent*** if $f(x, y) = f_X(x) \cdot f_Y(y)$.

If two random variables are not independent, they are ***dependent***.

Note: You can also check independence by seeing if the conditional distribution equals the marginal distribution, i.e., $p_{Y|X}(y|x) = p_Y(y)$ OR $p_{X|Y}(x|y) = p_X(x)$.

Let X and Y have the joint pdf, $f(x)$, as follows. Are X and Y independent?

$$f(x, y) = \begin{cases} \frac{4}{3}(xy + x) & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

An apartment complex has the following joint distribution of X bedrooms and Y bathrooms. Are X and Y independent?

Y | X	1	2	3	Σ
1	0.3	0.3	0	
2	0	0.3	0.1	
Σ				

Conditional Distributions

Let X and Y be two discrete distributions with joint pmf $p(x, y)$ and marginal pmf of X $p_X(x)$. Then for any X value of x for which $p_X(x) > 0$, the ***conditional probability mass function*** of Y given that $X = x$ is:

$$p_{Y|X}(y|x) = \frac{p(x, y)}{p_X(x)}$$

Let X and Y be two continuous distributions with joint pdf $f(x, y)$ and marginal pdf of X $f_X(x)$. Then for any X value of x for which $f_X(x) > 0$, the ***conditional probability density function*** of Y given that $X = x$ is:

$$f_{Y|X}(y|x) = \frac{f(x, y)}{f_X(x)}$$

For the apartment example, determine the conditional pmf $p_{Y|X}(y|x = 2)$. The joint pmf, $p_{X,Y}(x, y)$, is provided below.

Y | X	1	2	3	Σ
1	0.3	0.3	0	
2	0	0.3	0.1	
Σ				

Let X and Y have the joint pdf, $f(x, y)$, as follows. Determine the conditional pdf, $f_{Y|X}(y|x)$.

$$f(x, y) = \begin{cases} \frac{4}{3}(xy + x) & 0 \le x \le 1, 0 \le y \le 1 \\ 0 & \text{otherwise} \end{cases}$$

Expected Values, Covariance, and Correlation

Let X and Y be jointly distributed pmf $p(x, y)$ (if X and Y are discrete) or pdf $f(x, y)$ (if X and Y are continuous). The expected value of the function $h(X, Y)$, $E[h(X, Y)]$, is given by:

$$E[h(X,Y)] = \sum_x \sum_y h(x,y) \cdot p(x,y) \text{ (if X and Y are discrete)}$$

$$E[h(X,Y)] = \int_x \int_y h(x,y) \cdot f(x,y) dy dx \text{ (if X and Y are continuous)}$$

The ***covariance*** between two random variables is given by:

$$Cov(X,Y) = E[XY] - E[X]E[Y]$$

The ***correlation coefficient*** between two random variables is given by:

$$Corr(X,Y) = \rho_{XY} = \rho = \frac{Cov(X,Y)}{\sigma_X \cdot \sigma_Y}$$

For any two random variables X and Y, $-1 \leq \rho_{XY} \leq 1$.

If X and Y are independent then $\rho_{XY} = 0$

Note: $\rho_{XY} = 0$ does not imply that two random variables are independent!

For the apartment example, determine $E[X]$, $E[Y]$, $E[XY]$, and $Cov(X, Y)$. The joint pmf, $p_{X,Y}(x, y)$, is provided below.

Y | X	1	2	3	Σ
1	0.3	0.3	0	
2	0	0.3	0.1	
Σ				

Let X and Y be random variables and a and b be constants.

- $E[aX + bY] = aE[X] + bE[Y]$
- $Var(aX + bY) = a^2 Var(X) + b^2 Var(Y) + 2ab \cdot Cov(X, Y)$
- $Var(aX + bY) = a^2 Var(X) + b^2 Var(Y)$, if X and Y are independent

Bivariate Normal Distribution

The ***bivariate normal distribution*** of (X,Y) has the following joint probability distribution function:

$$f(x,y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \cdot exp\Bigg[-\frac{1}{2(1-\rho^2)}\Bigg(\Big(\frac{x-\mu_X}{\sigma_X}\Big)^2 + 2\rho\frac{(x-\mu_X)(y-\mu_Y)}{\sigma_1\sigma_2} + \Big(\frac{y-\mu_Y}{\sigma_Y}\Big)^2\Bigg)\Bigg]$$

$$-\infty < x < \infty, -\infty < y < \infty$$

where $E[X] = \mu_X$, $Var(X) = \sigma_X^2$, $E[Y] = \mu_Y$, $Var(Y) = \sigma_Y^2$, $\rho = Corr(X,Y)$




Figure 1: Bivariate Normal Distribution

We won't work directly with the multivariate normal distribution, but there are some useful properties of normal random variables which we will use.

If $X \sim \mathcal{N}(\mu_X, \sigma_X^2)$ and $Y \sim \mathcal{N}(\mu_Y, \sigma_Y^2)$, then: $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \sigma_X^2 + \sigma_Y^2 + 2Cov(X,Y))$, that is, the sum of normal random variables is a normal random variable.


If $X \sim \mathcal{N}(\mu_X, \sigma_X^2)$ and $Y \sim \mathcal{N}(\mu_Y, \sigma_Y^2)$ and X and Y are independent, then: $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \sigma_X^2 + \sigma_Y^2)$.

Suppose X and Y are independent normal random variables with means $\mu_X = 2$ and $\mu_Y = 3$ and variances $\sigma_X^2 = 1$ and $\sigma_Y^2 = 4$. Determine the distribution of $Z = 4X - Y$.

R Companion for Chapter 5

Let's return to the Hubble dataset. Let's load the dataset into R and then plot the data. Note the plot parameter **pch=16** fills in the data points for a nicer looking plot.

```
hubble = read_csv(file = "hubble.csv")
plot(Velocity~Distance,data=hubble, pch=16)
```



There appears to be a positive relationship between Distance and Velocity. We can calculate the correlation and covariance between the two variables as well.

```
cor(hubble$Distance,hubble$Velocity)
```

```
## [1] 0.789032
```

```
cov(hubble$Distance,hubble$Velocity)
```

```
## [1] 189.159
```

The correlation is +0.789 indicating a positive, moderately strong linear relationship between Distance and Velocity. The covariance is 189.159 which is less interpretable since covariance is dependent on the units of measurement. For this reason, we typically prefer to use correlation rather than covariance.

The **mtcars** dataset is provided in the base version of R. To find out more info about this dataset, you can get details by calling **?mtcars**.

```
# get info on the mtcars dataset
?mtcars
```

In the side window of R Studio, it should display the following description after running **?mtcars**.

The data were extracted from the 1974 Motor Trend US magazine, and comprises fuel consumption and 10 aspects of automobile design and performance for 32 automobiles (1973–74 models).

Let's load the dataset and take a look at the first few values.

```
data(mtcars)
head(mtcars)
```

```
##                   mpg cyl disp  hp drat    wt  qsec vs am gear carb
## Mazda RX4         21.0   6  160 110 3.90 2.620 16.46  0  1    4    4
## Mazda RX4 Wag     21.0   6  160 110 3.90 2.875 17.02  0  1    4    4
## Datsun 710        22.8   4  108  93 3.85 2.320 18.61  1  1    4    1
## Hornet 4 Drive    21.4   6  258 110 3.08 3.215 19.44  1  0    3    1
## Hornet Sportabout 18.7   8  360 175 3.15 3.440 17.02  0  0    3    2
## Valiant           18.1   6  225 105 2.76 3.460 20.22  1  0    3    1
```

In the code below, we extract just these two variables from **mtcars** and create a new dataset **m**. From this new dataset **m**, we can construct a two-way table of counts for the combinations of **cyl** and **gear**.

```
# creating new dataset m
m = mtcars[ , c("cyl", "gear")]

# construct a two-way table of counts for cyl and gear
m.table = table(m$cyl, m$gear)
m.table
```

```
##
##      3  4  5
##   4  1  8  2
##   6  2  4  1
##   8 12  0  2
```

We can also construct the joint probability mass function by dividing our table by the total number of observations.

```
# calculate the number of total observations
n = sum(m.table); n
```

```
## [1] 32
```

```
# construct the joint pmf
m.pmf = m.table/n; m.pmf
```

```
##
##           3       4       5
##   4 0.03125 0.25000 0.06250
##   6 0.06250 0.12500 0.03125
##   8 0.37500 0.00000 0.06250
```

Using the table, we can see that P(gear=3 and cyl=8) = 0.375 for example. What is P(gear=4 and cylinder = 4)? The middle entry of the table tells us this value is 0.125. In other words, 12.5% of the cars in this dataset have 4 gears and 4 cylinders.

Next, we are going to look at the joint probability mass function (PMF) of the variables **cyl** and **gear**. We can get the marginal PMFs by summing across rows and columns of the joint pmf.

```
# marginal pmf of gear
colSums(m.pmf)
```

```
##       3       4       5
## 0.46875 0.37500 0.15625
```

```
# marginal pmf of cyl
rowSums(m.pmf)
```

```
##       4       6       8
## 0.34375 0.21875 0.43750
```

From the tables above, you should see that P(gear=3)=0.46875 and P(cyl=4)=0.34375.

Finally, we can get the correlation as follows. Note that gear and cyl have a negative, moderate linear relationship.

```
cor(m$cyl,m$gear)
```

```
## [1] -0.4926866
```

Sampling Distributions, Central Limit Theorem, and Estimation

In this chapter, we will explore sampling distributions, focusing on the variability of sample statistics, which is crucial for inference. We will then dive into the Central Limit Theorem (CLT), which demonstrates how the distribution of sample means approaches normality as the sample size grows, forming the backbone of many statistical methods. Finally, we will introduce point estimation, highlighting how to derive reliable population parameter estimates through sample data and discussing properties such as bias and standard error.

Statistics and Their Distributions

A ***statistic*** is any quantity whose value can be calculated from sample data.

The random variables $X_1, X_2, \ldots, X_n$ are said to form a ***random sample*** of size n if X_i are ***independent and identically distributed (IID)***.

In chapter 1, we came across a few statistics. We can measure the center of a sample with the sample mean, $\bar{x}$ and the sample median, $\tilde{x}$. If we want to describe the spread of a sample, there are the sample variance (s^2) and sample standard deviation (s).

In fact, since each random sample could potentially be different, sample statistics have a probability distribution associated with them.

The Distribution of the Sample Mean


Suppose a researcher claims that Radon-222 has a half-life (median decay time) of 6 days and the the time between particles decaying can be modeled as an exponential($\lambda = 0.1155$) random variable. This random variable has a mean time of 8.66 days between decays. To test the accuracy of this claim, we measured the decay times for 50 Radon-222 particles and found a sample mean of $\bar{x} = 5.67$ and sample variance of $s^2 = 28.94$. Do we have enough information to reject the claim?

To answer this question, we need to know the distribution of $\bar{X}$ and specifically, we would like to calculate the probability of observing such a sample mean assuming the historical estimate is accurate.

Let's first explore the sampling distribtuion on $\bar{X}$ using simulations.

(a) Simulate 10,000 Radon-222 decay times and plot the data as a histogram.

```
set.seed(2026)
sims = rexp(n=10000,rate=0.1155)
hist(sims,main="10,000 simulations of Radon-222 decay times")
```




```
mean(sims)
```

```
## [1] 8.916651
```

```
var(sims)
```

```
## [1] 77.21487
```

(b) Simulate four exponential($\lambda = 0.1155$) random variables five times and calculate the sample mean of each sample.

```
set.seed(1)
sample1 = rexp(n=4,rate=0.1155); mean1 = mean(sample1)
sample2 = rexp(n=4,rate=0.1155); mean2 = mean(sample2)
sample3 = rexp(n=4,rate=0.1155); mean3 = mean(sample3)
sample4 = rexp(n=4,rate=0.1155); mean4 = mean(sample4)
sample5 = rexp(n=4,rate=0.1155); mean5 = mean(sample5)
```

x_1	x_2	x_3	x_4	$\bar{x}$
6.538371	10.230673	1.261530	1.210349	4.810231
3.775486	25.064663	10.645559	4.672579	11.039572
8.281970	1.273125	12.040997	6.597661	7.048438
10.715182	38.302461	9.130244	8.963151	16.777759
16.242729	5.668802	2.917173	5.095063	7.480942

(c) Compute the mean and variance of the five sample means.

```
means = c(mean1,mean2,mean3,mean4,mean5)
mean(means)
```

```
## [1] 9.431388
```

```
var(means)
```


```
## [1] 21.8483
```

(d) Repeat the experiment 10,000 times, that is, simulate 10,000 samples of 4 exponential($\lambda = 0.1155$) random variables. Create a histogram of the sample means and calculate the mean and variance of the sample means.

```
set.seed(2026)
nsims = 10000
means = rep(0,nsims)
simdata = rep(0,4)

for(i in 1:nsims){
  simdata = rexp(n=4,rate=0.1155)
  means[i] = mean(simdata)
}

hist(means,main="10,000 simulations of sample means of 4 Radon-222 decay times")
```




```
mean(means)
```

```
## [1] 8.649116
```

```
var(means)
```

```
## [1] 18.9354
```

Let $X_1, X_2, \ldots, X_n$ be a random sample from a distribution with mean value μ and standard deviation σ and let $\bar{X} = \frac{1}{n}\sum X_i$. Then:

- $E[\bar{X}] = \mu$
- $Var(\bar{X}) = \sigma^2_{\bar{X}} = \frac{\sigma^2}{n}$
- $\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$

Show that $Var(\bar{X}) = \frac{\sigma^2}{n}$.

If $\hat{\theta}$ is an estimator for θ, then the ***standard error*** of $\hat{\theta}$ is: $SE(\hat{\theta}) = \sqrt{Var(\hat{\theta})}$. It is the magnitude of a typical or representative deviation between an estimate and the value of θ. If the standard error is a function of unknown parameters, then we can estimate it with the ***estimated standard error*** which is denoted $s_{\hat{\theta}}$


If $X_1, X_2, \ldots, X_{50}$ are iid Uniform[10,20] random variables, then determine $E[\bar{X}]$ and $SE(\bar{X})$.

Let's go back to our simulations to explore the shape of the sampling distribution of sample means.

```
set.seed(2026)
nsims = 10000
n = c(4, 10, 30, 100)
means = cbind(rep(0,nsims), rep(0,nsims), rep(0,nsims), rep(0,nsims))
count = 1

for(j in n){
for(i in 1:nsims){
  simdata = rexp(n=j,rate=0.1155)
  means[i,count] = mean(simdata)
}
  count = count +1
}

par(mfrow = c(1, 4))
hist(means[ , 1], main = "",
     sub = "n = 4",
     xlab = "Sample means", prob = TRUE)
lines(density(means[ , 1]),lwd=2,lty=2)
hist(means[ , 2], main = "",
     sub = "n = 10",
     xlab = "Sample means", prob = TRUE)
lines(density(means[ , 2]),lwd=2,lty=2)
hist(means[ , 3], main = "",
     sub = "n = 30",
     xlab = "Sample means", prob = TRUE)
lines(density(means[ , 3]),lwd=2,lty=2)
hist(means[ , 4], main = "",
     sub = "n = 100",
     xlab = "Sample means", prob = TRUE)
lines(density(means[ , 4]),lwd=2,lty=2)
```



	n = 4	n = 10	n = 30	n = 100
Mean of Means	8.649116	8.627836	8.625484	8.667377
Var of Means	18.935399	7.571442	2.497474	0.754795

Central Limit Theorem: Let $X_1, X_2, \ldots, X_n$ be a random sample from a distribution with mean μ and variance σ^2. If n is sufficiently large, $\bar{X}$ has approximately a normal distribution with mean $\mu_{\bar{X}} = \mu$ and variance $\sigma^2_{\bar{X}} = \sigma^2/n$. In other words, $\bar{X} \dot{\sim} \mathcal{N}(\mu, \sigma^2)$.

Note: The symbol, $\dot{\sim}$, stands for "approximately distributed as".

This result is nontrivial and extremely useful! It allows us to determine how likely or unlikely a sample mean is if we are given a hypothesized population mean.

Suppose that $X_1, X_2, \ldots, X_{50}$ is a random sample from an exponential($\lambda = 0.1155$) random variable (the historical claim). Determine the approximate sampling distribution of $\bar{X}$.

Suppose the decay time of Radon-222 follows an exponential distribution with $\lambda = 0.1155$ decays/day (the historical claim). What is the probability that a random sample of 50 Radon particles results in a sample mean of less than 5.67 days/decay? Do you believe the historical claim?

Suppose $X_1, X_2, \ldots, X_{50}$ are iid Uniform[10,20] random variables. Determine the sampling distribution of $\bar{X}$ and calculate $P(\bar{X} < 14)$.

Several General Concepts of Point Estimation

A ***point estimate*** of a parameter θ is a single number that can be regarded as a sensible value for θ and is often denoted with a hat (i.e., $\hat{\theta}$). A point estimate is obtained by selecting a suitable statistic and computing its value from the given sample data. The selected statistic, $\hat{\theta}$, is called the ***point estimator*** of θ.

1. $\bar{X} = \frac{1}{n}\sum_{i=1}^{n} X_i$
2. $S^2 = \frac{1}{n-1}\sum_{i=1}^{n}(X_i - \bar{X})^2$

The ***bias*** of a point estimator $\hat{\theta}$ for a parameter θ is defined as $Bias[\hat{\theta}] = E[\hat{\theta}] - \theta$.

A statistic $\hat{\theta}$ is called an ***unbiased estimator*** of the parameter θ if $E[\hat{\theta}] = \theta$.

$\bar{X}$ is an unbiased estimator for μ.

Which of the following two estimators for σ^2 is unbiased for σ^2?

- $S^2 = \frac{1}{n-1}\sum_{i=1}^{n}(X_i - \bar{X})^2$
- $\gamma = \frac{1}{n}\sum_{i=1}^{n}(X_i - \bar{X})^2$

The ***mean squared error (MSE)*** of an estimator $\hat{\theta}$ of a parameter θ is defined by $E[(\hat{\theta} - \theta)^2] = Var + Bias^2$.

Typically, unbiased estimators are preferred though some advanced methods consider biased estimators that minimize MSE.

Suppose $X_1, X_2, \ldots, X_n$ are exponential(λ) random variables. Is $\bar{X}$ an unbiased estimator for λ?

Suppose $X_1, X_2, \ldots, X_n$ are Poisson(λ) random variables. Is $\bar{X}$ an unbiased estimator for λ?

Methods of Point Estimation

There are a variety of ways to obtain a point estimate. These methods include:

- Method of Moments
- Maximum Likelihood Estimation
- Bayesian Estimation

While we won't cover how these estimation methods are completed, we will compare different estimators. Each of these methods could produce a different point estimate.

Suppose X_1, X_2, X_3 are a random sample of $\mathcal{N}(\mu = 10, \sigma^2 = 4)$. Four estimators are given below. Determine if each estimator is unbiased and calculate its standard error. Which of the four estimators is preferred for estimating μ?

(a) $\hat{Y_1} = \frac{X_1+X_2+X_3}{3}$

(b) $\hat{Y_2} = \frac{X_1}{2} + \frac{X_2}{4} + \frac{X_3}{4}$

(c) $\hat{Y_3} = X_1$


(d) $\hat{Y_4} = \frac{X_1+X_2+X_3}{2}$

R Companion for Chapter 6

What does the distribution of the sample means of independent (continuous) uniform random variables look like? The Central Limit Theorem tells us that for a large sample size, the distribution of such means will be approximately normally distributed. Let's see this in action.

Suppose we know a bus arrives once per hour at a designated stop but we don't know when the next bus is and we don't know when the last bus came. We will model this as a continuous uniform[0,60] random variable. In other words, the next bus could come in anywhere from 0 to 60 minutes with all possibilities being equally likely. In the code below, we simulate 10,000 uniform[0,60] random variables and look at the histogram of results.

```
# Generate 10,000 uniform[0,60] random varibles
set.seed(2020)
bus = runif(10000,0,60)
hist(bus)
```



The mean and variance of this sampling distribution is calculated below.

```
mean(bus)
```

```
## [1] 29.79179
```

```
var(bus)
```

```
## [1] 298.1569
```

From chapter 4, we know that if $X \sim Uniform[a,b]$, then $E[X] = \frac{b+a}{2}$ and $Var(X) = \frac{(b-a)^2}{12}$. For a uniform[0,60] random variable, we get the theoretical values of $E[X] = \frac{60+0}{2} = 30$ and $Var(X) = \frac{(60-0)^2}{12} = 300$. Our simulated values are very close to these theoretical values.

Now, let's suppose we have to wait for two buses, each being modeled as an independent uniform[0,60]. Let's look at the sampling distribution for the mean of the two bus wait times.

```
set.seed(2020)
bus2 = rep(0, 10000) # initialize this vector to zeros

# 10,000 simulations of 2 bus waits
for (i in 1:10000){
x1 = runif(2,0,60) # n=2
bus2[i] = mean(x1)
}


mean(bus2)
```

```
## [1] 29.90972
```

```
var(bus2)
```

```
## [1] 149.6154
```

```
hist(bus2)
```



Notice that the average of two bus waits is no longer uniform and instead has more of a triangular shape. Our simulations of two bus waits had a mean of 30.05955 and variance of 151.2226. From chapter 6, we know that $E[\bar{X}] = \mu$ and $Var(\bar{X}) = \frac{\sigma^2}{n}$.

For this simulation, the theoretical values for these quantities are $E[\bar{X}] = \mu = 30$ and $Var(\bar{X}) = \frac{\sigma^2}{n} = \frac{300}{2} = 150$. Again, our simulations give values very close to the theoretical values.

Finally, let's suppose that we repeat the experiment for 30 days straight and look at the distribution of sample means of 30 days. We will simulate these 30 day periods 10,000 times.

```
set.seed(2020)
bus30 = rep(0, 10000) # initialize this vector to zeros

# 10,000 simulations of 2 bus waits
for (i in 1:10000){
x2 = runif(30,0,60) # n=30
bus30[i] = mean(x2)
}


mean(bus30)
```

```
## [1] 29.97879
```

```
var(bus30)
```

```
## [1] 10.19619
```

```
hist(bus30)
```



Our simulations of 30 bus waits had a mean of 30.00598 and variance of 10.08975. The theoretical values for this simulation are $E[\bar{X}] = \mu = 30$ and $Var(\bar{X}) = \frac{\sigma^2}{n} = \frac{300}{30} = 10$.

Notice that the distribution of sample means is narrower for n=30 than n=2 and the variance is lower when a larger sample size is used. Also note that the distribution is somewhat bell-shaped.

In the coming chapters on Inferential Statistics, we will use the fact that the Central Limit Theorem states that the sampling distribution of sample means is approximately normally distributed.

Confidence Intervals for One Population Parameter

In this chapter, we will explore how to construct confidence intervals for one-parameter population values, such as the population mean or proportion. In Chapter 6, we discussed how to derive a point estimate for a population parameter. Since each sample may produce a different point estimate, it's essential to account for the inherent randomness in this process. Confidence intervals provide a way to quantify the uncertainty associated with these estimates by creating a range of plausible values for the parameter, rather than relying on a single estimate. This approach gives us a more reliable method to assess the precision of our estimates.

Basic Concepts of Confidence Intervals

In Chapter 6, we examined how to find a point estimate for an unknown population parameter. When reporting point estimates, it is important to note the amount of uncertainty associated with the point estimate. This is especially key when we want to do Statistical Inference.

We will examine two methods of Statistical Inference:

- Confidence Intervals
- Hypothesis Tests

A ***confidence interval*** is a range of values, derived from sample data, that reflects the uncertainty associated with an estimate of an unknown population parameter. It provides an interval within which the parameter is likely to fall, with a specified level of confidence (e.g., 95%). The confidence level indicates the probability that the interval will contain the true parameter in repeated sampling.

There are many flavors of confidence intervals:

- Some are one-sided while others are two-sided
- Some assume population variance is known while others do not
- Some assume a higher confidence level while others assume a lower confidence level

General Form of a Confidence Interval (Two-Sided)

Confidence Intervals for a Population Mean when the Population Standard Deviation is Known.

Recall that if we take a random sample from a normal population or if n is 'large' enough (by the Central Limit Theorem) then $\bar{X} \sim N(\mu, \frac{\sigma^2}{n})$. Further, from Chapter 4, we know that $Z = \frac{(\bar{X}-\mu)}{\frac{\sigma}{\sqrt{n}}}$ has a standard normal distribution.

We can use this result to produce a confidence interval.

If our random sample comes from a normal distribution or n is 'large' and σ is known, then a $100(1-\alpha)\%$ confidence interval for μ is given by: $\bar{x} \pm z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}$

α is the probability that a confidence interval from repeated sampling does not contain the true population parameter. It represents the likelihood of the confidence interval failing to capture the parameter in repeated samples, and is equal to $1 -$ confidence level.

Suppose we collect a random sample and use it to create a confidence interval. Further, suppose that we do this process 100 times and create 100 confidence intervals.

Suppose we want to create a two-sided 95% confidence interval for μ.

Writing and Interpreting a Confidence Interval

Here's how to write confidence intervals:

Here's how to interpret confidence intervals:

A random sample of 16 CPU chips has a total of 342 defects. Suppose we know the population standard deviation is 4.52 and also that the number of defects/chip is approximately normally distributed.

(a) Calculate a point estimate for μ, the true mean number of defects/chip.

(b) Calculate a 95% CI for μ.

Suppose that GPAs are approximately normally distributed with population variance $\sigma^2 = 1.2$. If we take a random sample of 12 students and find a sample mean of 2.87. Construct a 90% confidence interval for μ.

Confidence Intervals for a Population Mean when the Population Standard Deviation is Unknown.

In this section, we will assume that both μ and σ are unknown.

Recall that if we take a random sample from a normal population or if n is 'large' enough (by the Central Limit Theorem) then $\bar{X} \sim N(\mu, \frac{\sigma^2}{n})$. Further, from Chapter 4, we know that $Z = \frac{(\bar{X}-\mu)}{\frac{\sigma}{\sqrt{n}}}$ has a standard normal distribution.

If σ is unknown, then $\frac{(\bar{X}-\mu)}{\frac{S}{\sqrt{n}}}$ does ***not*** have a normal distribution.

The ***t-distribution*** is a probability distribution used in statistics that is similar to the normal distribution but has heavier tails. It is typically used when estimating population parameters when the sample size is small and/or the population standard deviation is unknown. The t-distribution becomes closer to the normal distribution as the sample size increases. t_k denotes a t-distribution with k degrees of freedom.

Properties of the t-distribution

- Each t_k curve is bell-shaped and centered at zero.
- Each t_k curve has more spread than the standard normal curve.
- As k increases, the spread of the corresponding t_k curve decreases.
- As $k \to \infty$, the sequence of t_k curves approaches the standard normal curve.

Find t^* such that the area under the curve to the right of t^* is 0.025 for $df = 10$.

Find t^* such that the area under the curve to the left of t^* is 0.05 for $df = 14$.

Find the two values, t_1 and t_2, such that the area under the curve is the middle 95% for $df = 24$.

Suppose $t^* = 2.1$ and $df = 9$. Calculate $P(t(9) \geq t^*)$.

If $\bar{X}$ is the mean of a random sample of size n from an (approximately) normal distribution (or n large enough) with mean μ and sample standard devitation s, then $t = \frac{(\bar{X}-\mu)}{(\frac{S}{\sqrt{n}})}$ follows a t-distribution with (n-1) degrees of freedom.

Let $\bar{x}$ and s be the sample mean and sample standard deviation from a random sample from a (approximately) normal population (or n is large enough) with mean μ. Then a $100(1-\alpha)\%$ confidence interval for μ is given by: $\bar{x} \pm t_{\alpha/2} \cdot \frac{s}{\sqrt{n}}$, df=n-1

Suppose a random sample of 30 Hummers have a mean of 9.8 miles/gallon (MPG) and a sample standard deviation of 2.2. Compute a 95% confidence interval for μ.

Suppose a random sample of 17 Priuses have a mean of 52.3 MPG and variance of 14.9. Construct a 99% confidence interval for μ.

Confidence Intervals for a Single Population Proportion.

Provided that n is sufficiently large, a $100(1-\alpha)\%$ confidence interval for a population proportion p is given by $\hat{p} \pm z_{\alpha/2} * \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$.

Suppose a random sample of 80 students finds 45 of them spend more than 20 hours/week on homework. Construct a 95% confidence interval for the true proportion of students that spend more 20 hours/week on homework.

One-Sided Confidence Intervals

Suppose a random sample of 30 Hummers have a mean of 9.8 miles/gallon (MPG) and a sample standard deviation of 2.2. Find the upper bound for a 95% lower confidence interval for μ.

Suppose a random sample of 17 Priuses have a mean of 52.3 MPG abd a sample variance of 14.9. Find the lower bound for a 99% upper confidence interval for μ.

R Companion for Chapter 7

The following example uses the dataset **faithful** which contains the duration of eruption (in minutes) and waiting times between eruptions for Old Faithful geyser in Yellowstone National Park, Wyoming, USA.

```
# look up info on the faithful dataset
?faithful

#load the faithful dataset
data(faithful)
```

We can use the **t.test** function to run a hypothesis test (see Chapter 8) and to create a confidence interval for an unknown parameter.

Suppose we want to create a 95% confidence interval for the true duration of eruptions for Old Faithful. By default, R will produce a 95% confidence interval unless otherwise specified. Since the population variance of eruption times for Old Faithful is unknown, the confidence interval will be created using the t-distribution.

```
t.test(faithful$eruptions)
```

```
##
##  One Sample t-test
##
## data:  faithful$eruptions
## t = 50.397, df = 271, p-value < 2.2e-16
## alternative hypothesis: true mean is not equal to 0
## 95 percent confidence interval:
##  3.351534 3.624032
## sample estimates:
## mean of x
##  3.487783
```

In the output, we see that the 95% confidence interval for the true mean eruption duration is $3.351534 < \mu < 3.624032$. If someone were to claim that the average eruption duration of Old Faithful is 4 minutes, we would reject this claim.

We could create a 99% confidence interval for the true mean eruption duration as well. This will be a wider confidence interval since we are requiring a higher confidence.

```
t.test(faithful$eruptions, conf.level = 0.99)
```

```
##
##  One Sample t-test
##
## data:  faithful$eruptions
## t = 50.397, df = 271, p-value < 2.2e-16
## alternative hypothesis: true mean is not equal to 0
## 99 percent confidence interval:
##  3.308257 3.667309
## sample estimates:
```

```
## mean of x
##  3.487783
```

The 99% confidence interval for the true mean eruption duration is $3.308257 < \mu < 3.667309$.

We can also create one-sided confidence intervals by using the **alternative** argument with either **greater** or **less**. Below, a 90% upper one-sided confidence interval is created.

```
t.test(faithful$eruptions, conf.level = 0.9, alternative = "greater")
```

```
##
##  One Sample t-test
##
## data:  faithful$eruptions
## t = 50.397, df = 271, p-value < 2.2e-16
## alternative hypothesis: true mean is greater than 0
## 90 percent confidence interval:
##  3.398876      Inf
## sample estimates:
## mean of x
##  3.487783
```

The 90% upper one-sided confidence interval is $3.398876 < \mu$. We are 90% confident that the true mean eruption duration of Old Faithful is at least 3.339 minutes.

Hypothesis Tests for One Population Parameter

In the previous chapter, we explored Statistical Inference through the application of confidence intervals, which allowed us to estimate population parameters within a range of plausible values. In this chapter, we will turn our attention to hypothesis testing, another fundamental approach in Statistical Inference. Hypothesis testing provides a structured framework to evaluate specific claims or assumptions about a population parameter by analyzing sample data. Through this method, we can determine whether the evidence supports or refutes a given hypothesis, helping us make informed decisions in the presence of uncertainty.

Basic Concepts of Hypothesis Testing

Imagine a company claims that its employees spend, on average, 5 hours per week using social media during work hours. The management is concerned that this may be affecting productivity and wants to verify if the average time is indeed 5 hours. To test this claim, a random sample of employees is surveyed, and the average time spent on social media is calculated. By formulating and testing a hypothesis, the company can assess whether the claim holds up under scrutiny or if the evidence suggests a different average usage.

- One way to test this claim would be to collect a sample and see if 5 is contained in the confidence interval for μ.
- Another approach to testing this claim is to again collect a sample and see how consistent that sample is with initial hypothesis of $\mu = 5$. This is called hypothesis testing.

Goal of Hypothesis Testing: Test competing claims about a population parameter.

The ***null hypothesis***, denoted H_0, represents the initial claim or assumption that is presumed to be true in the absence of evidence to the contrary.

The ***alternative hypothesis***, denoted H_a, is the statement that contradicts H_0, often described as the opposite of the null hypothesis, and represents what we are testing for in the analysis.

- The null hypothesis will be rejected in favor of the alternative hypothesis if the sample evidence suggests that H_0 is false.
- The two possible conclusions of a hypothesis test are **Reject** H_0 or **Fail to Reject** H_0.

Formulating Hypotheses

1. H_0 contains $=, \leq, \geq$.
2. H_a contains $\neq, >, <$
3. Usually, the claim we are attempting to show is more plausible is H_a

Jorge claims that homework for this class takes, on average, six hours per week. Aaron doesn't think the average is six hours, so he collects a random sample of eight students and records the time it takes them to complete the homework.

Brenda claims that newly graduated data science majors from CSU have an average salary of \$63,000, but Rawand believes it is smaller.

Rosa believes less than 50% of college students have consumed alcohol. You collect a random sample of 20 students to test the claim.

Different Types of Hypothesis Tests

There are three different types of hypothesis tests.

1. Left-tailed test

2. Right-tailed test

3. Two-tailed test

Errors in Hypothesis Testing

After collecting a sample, we will decide which hypothesis is supported by the data.

Since we don't know the true parameter value, there is a chance we will make an error.

Two Types of Error

1. Type I error (α): Reject H_0 when H_0 is true.
2. Type II error (β): Fail to reject H_0 when H_0 is false.

Note: α is also called the level of significance for a test.

Jury trials, H_0 : Defendant is innocent vs. H_A : Defendant is guilty

Six Steps for Hypothesis Testing

Different textbooks will give a different number of steps or outline on how to complete a hypothesis test. For our class, stick to the following method.

Step 1: State the hypotheses.

Step 2: Determine the level of significance.

Step 3: Compute a test statistic.

Step 4: Calculate a p-value and plot the results.

The ***p-value*** is the probability of observing a test statistic as extreme as, or more extreme than, the one obtained from the sample data, assuming the null hypothesis is true.

The p-value does not measure the probability that the null hypothesis is true.

ASA Statement on p-values: https://www.amstat.org/asa/files/pdfs/p-valuestatement.pdf

Step 5: Make a statistical decision.

Step 6: Interpret the statistical decision (in the context of the problem).

Tests About a Population Mean, μ

Case 1: σ known

If σ is known, then the test statistic is the following:

$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}}$$

A materials scientist is developing a new alloy intended for use in aerospace engineering, where the average tensile strength is claimed to be 1500 MPa (megapascals). Based on previous research, the tensile strength of this alloy is known to be normally distributed with a population variance of 40 MPa^2.

To validate the claim, the scientist tests a sample of 10 specimens and finds a sample mean tensile strength of 1495 MPa. The goal is to perform a hypothesis test at a significance level of $\alpha = 0.05$ to determine whether the average tensile strength is actually different from the claimed 1500 MPa.

An engineering firm claims that a new fuel-efficient engine design provides an average fuel efficiency of at least 40 miles per gallon (mpg). The design specifications guarantee this minimum efficiency, and engineers want to verify that the engine consistently meets this standard. It is known from previous tests that fuel efficiency for this type of engine follows a normal distribution with a population variance of 9 mpg^2.

A random sample of 8 engines is tested, and the sample shows an average fuel efficiency of 38.5 mpg. The engineers decide to conduct a one-sided hypothesis test at a significance level of $\alpha = 0.05$ to determine if the true average fuel efficiency is actually lower than the claimed 40 mpg.

In addition, state what Type I and Type II errors are in the context of the problem.

Case 2: σ unknown

If σ is unknown, then the test statistic is the following:

$$t_{stat} = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}, df = n - 1$$

A chemist claims that a newly synthesized chemical compound has an average melting point of 250°C. However, the chemist does not know the population variance for the melting point of this compound. To test the consistency of the compound, a sample of 31 batches is produced and their melting points are measured. The sample has a mean melting point of 247°C, with a sample standard deviation of 8.4°C.

The chemist wants to perform a hypothesis test at a significance level of $\alpha = 0.10$ to determine whether the true average melting point is different from the claimed 250°C.

A software engineer claims that a new web application backend has an average response time of less than 200 milliseconds under typical user load. However, the team wants to verify this claim to ensure that the system meets performance expectations.

To test this, a sample of 50 response times is collected from the application under standard load conditions. The sample mean response time is 210 milliseconds, with a sample standard deviation of 25 milliseconds. The engineer performs a one-sided hypothesis test at a significance level of $\alpha = 0.05$ to determine if the true average response time is actually greater than the claimed 200 milliseconds.

Tests Concerning a Population Proportion, p

We can run a hypothesis test for a population proportion using the following formula:

$$z = \frac{\bar{p} - p_0}{\sqrt{p_0(1-p_0)/n}}$$

A mathematics department claims that at least 80% of its students pass a standardized math proficiency exam required for graduation. A new curriculum is introduced, and the department wants to evaluate whether the pass rate has fallen below this benchmark.

To test this, a random sample of 150 students who took the exam under the new curriculum is selected, and 112 students passed. The department conducts a one-sided hypothesis test at a significance level of $\alpha = 0.05$ to determine if the true pass rate is now less than the claimed 80%.

R Companion for Chapter 8

The **faithful** dataset of Old Faith Geyser data is used again for this example. Since the population variance of Old Faithful eruption duration is unknown, the t-distribution is used to complete the following hypothesis tests.

Let μ denote the true mean eruption duration of Old Faithful and we want to test $H_0 : \mu = 3$ vs. $H_a : \mu \neq 3$ at $\alpha = 0.05$.

```
t.test(faithful$eruptions,mu = 3)
```

```
##
##  One Sample t-test
##
## data:  faithful$eruptions
## t = 7.0483, df = 271, p-value = 1.502e-11
## alternative hypothesis: true mean is not equal to 3
## 95 percent confidence interval:
##  3.351534 3.624032
## sample estimates:
## mean of x
##  3.487783
```

From the output, we see that the p-value is $1.5 \cdot 10^{-11}$. This is less than $\alpha = 0.05$, so we would reject $H_0 : \mu = 3$ and conclude the true mean eruption duration of Old Faithful differs from 3 minutes.

Suppose we want to test $H_0 : \mu = 3.5$ vs. $H_a : \mu < 3.5$ at $\alpha = 0.05$. In other words, we would like to test to see if the mean is less than 3.5 minutes.

```
t.test(faithful$eruptions,mu=3.5,alternative = "less")
```

```
##
##  One Sample t-test
##
## data:  faithful$eruptions
## t = -0.17653, df = 271, p-value = 0.43
## alternative hypothesis: true mean is less than 3.5
## 95 percent confidence interval:
##      -Inf 3.602007
## sample estimates:
## mean of x
##  3.487783
```

Here, the p-value is 0.43, so we would fail to reject $H_0 : \mu = 3.5$. In other words, we don't have sufficient evidence to conclude the true mean eruption duration is less than 3.5 minutes.

Inference Based On Two Samples

In Chapters 7 and 8, we explored hypothesis tests and confidence intervals for making statistical inferences about a single population parameter. In this chapter, we will expand these methods to compare two population parameters. We'll cover hypothesis tests for both independent and paired samples using z-tests and t-tests, as well as tests for comparing two population proportions. Additionally, we will introduce confidence intervals for the difference between two parameters and discuss the concept of statistical power in the context of two-sample inference.

z-tests and Confidence Intervals for a Difference Between Two Population Parameters

A manufacturer produces two types of industrial batteries, Model A and Model B, which are used in renewable energy systems. The company claims that the lifespan of both battery models is approximately the same, but an engineering team wants to verify this by conducting a study. They know the population standard deviations of the lifespans for both models from previous extensive testing.

A sample of 15 Model A batteries has an average lifespan of 8.5 years with a known population standard deviation of 1.2 years. A sample of 15 Model B batteries has an average lifespan of 9.1 years with a known population standard deviation of 1.5 years. The engineers perform a hypothesis test to determine if there is sufficient evidence to suggest a difference in the average lifespan between the two battery models.

(a) One way to test this claim is create a confidence interval for the difference in means $(\mu_1 - \mu_2)$.
(b) Another approach to testing this claim is to see how consistent that sample is with initial hypothesis of $\mu_1 - \mu_2 = 0$.

Note: We must make an assumption of whether the samples from the two populations are independent or dependent. In this section (9.1) and the next section (9.2), we will assume the samples are independent.

Assumptions for Comparison of Independent Samples

(1) $X_1, X_2, \ldots, X_{n_1}$ are a random sample from a distribution with mean μ_1 and variance σ_1^2.
(2) $Y_1, Y_2, \ldots, Y_{n_2}$ are a random sample from a distribution with mean μ_2 and variance σ_2^2.
(3) The X and Y samples are independent.

Notation

	True Population Parameters			Sample Statistics		
Group	Mean	Variance	SD	Mean	Variance	SD
Group #1	μ_1	σ_1^2	σ_1	$\bar{x}_1$	s_1^2	s_1
Group #2	μ_2	σ_2^2	σ_2	$\bar{x}_2$	s_2^2	s_2

If X and Y are independent, then

(a) $E[\bar{X} - \bar{Y}] = \mu_1 - \mu_2$

(b) $Var(\bar{X} - \bar{Y}) = \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}$ (σ_1^2 and σ_2^2 may be unknown)

Proof:

Writing Hypotheses for Difference in Means

(a) Testing $\mu_1 \neq \mu_2$

(b) Testing $\mu_1 > \mu_2$

(c) Testing $\mu_1 < \mu_2$

We can complete a hypothesis test for the difference in means (assuming the variances or standard deviations of the two populations are known) by using the following test statistic.

$$z = \frac{(\bar{x_1} - \bar{x_2}) - D_0)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

A sample of 15 Model A batteries has an average lifespan of 8.5 years with a known population standard deviation of 1.2 years. A sample of 15 Model B batteries has an average lifespan of 9.1 years with a known population standard deviation of 1.5 years. The engineers perform a hypothesis test to determine if there is sufficient evidence to suggest a difference in the average lifespan between the two battery models.

A two-sided confidence interval for the difference in means can also be completed with the following formula, assuming again that the population standard deviations are known.

$$(\bar{x}_1 - \bar{x}_2) \pm z_{\alpha/2} \cdot \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

The Two-Sample t-test and Confidence Interval

If the population standard deviations of the two populations are unknown, we can estimate them and use similar methods as in section 9.1.

A confidence interval for the difference in means can be completed with the following formula if the population standard deviations are unknown.

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2,\nu} \cdot \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

For the above confidence interval, you must calculate the degrees of freedom (ν). Round the value of ν to the next lowest integer if it is not a whole number.

$$\nu = \frac{\left(s_1^2/n_1 + s_2^2/n_2\right)^2}{(s_1^2/n_1)^2/(n_1 - 1) + (s_2^2/n_2)^2/(n_2 - 1)}$$

It has been claimed that people in Okinawa live longer on average than people on the Japanese mainland. To investigate this, a study is conducted. A random sample of 25 people from Okinawa shows an average lifespan of 84.7 years, with a sample variance of 25 years^2. Meanwhile, a sample of 22 people from the Japanese mainland finds an average lifespan of 83.2 years, with a sample variance of 20 years^2.

We will use this data to calculate a 95% confidence interval for the difference in average lifespan between Okinawa and the Japanese mainland. Given the sample sizes and variances, the degrees of freedom for the test can be approximated as 43. The goal is to determine whether there is a statistically significant difference in the average lifespan between the two populations.

We can complete a hypothesis test for the difference in means (assuming the variances or standard deviations of the two populations are unknown) by using the following test statistic. This is called a **two-sample t-test**.

$$t = \frac{(\bar{x}_1 - \bar{x}_2 - D_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}, df = \nu$$

$$\nu = \frac{\left(s_1^2/n_1 + s_2^2/n_2\right)^2}{(s_1^2/n_1)^2/(n_1 - 1) + (s_2^2/n_2)^2/(n_2 - 1)}$$

Using the data from the Okinawa life expectancy study, conduct a hypothesis test at a significance level of $\alpha = 0.05$ to determine if the average lifespan in Okinawa (μ_1) is significantly greater than the average lifespan on the Japanese mainland (μ_2).

Analysis of Paired Data

In the previous two sections, we assumed that the two samples were independent. However, this assumption does not always hold true in practice.

Examples of paired samples include:

- **Before/after studies**: The same individuals are measured before and after an intervention.
- **Matched-pair designs**: For example, comparing blood pressure in the right and left arms of the same individuals.
- **Sibling studies**: Measuring certain traits (e.g., height or test scores) in siblings to see if there are significant differences.
- **Repeated measurements**: Taking the same measurement under different conditions (e.g., testing reaction time under different levels of noise for the same participants).

A sample is **paired** if the same subjects or closely matched subjects are measured twice under different conditions.

A **paired t-test** helps control for differences between individuals by focusing on the differences between the paired measurements.

Advantage: A paired t-test allows us to use a **smaller sample size** because it reduces the variance that comes from differences between individuals, leaving only the variation between the paired measurements themselves.

Disadvantage: It's not always **feasible** to collect paired data. If this is the case, you can use a two-sample t-test under the assumption that the samples are independent.

CI and Hypothesis Tests for Paired Samples (μ_d)

When doing inference for the true average change (μ_d), we compare "treatments" (e.g. before/after) on an individual level.

Create a "new", single sample based on the sample differences:

1. For each individual, define d_i = difference between treatments for subject i.
2. Calculate $\bar{d} = \frac{1}{n} \cdot \sum d_i$
3. Calculate $s_d = \sqrt{\frac{\sum (d_i - \bar{d})^2}{n-1}}$

Using the "new' ' sample, use the same procedures for inference as for a single population.

$$t = \frac{\bar{d} - \mu_{d_0}}{s_d/\sqrt{n}}, df = n - 1$$

$$\bar{d} \pm t_{\alpha/2, n-1} \cdot \frac{s_d}{\sqrt{n}}$$

A pharmaceutical company is testing a new drug designed to lower blood pressure. They select a group of 10 patients with high blood pressure and measure their systolic blood pressure before and after administering the drug. The company wants to know if the drug significantly lowers blood pressure. Assume that the differences in blood pressure before and after the treatment are approximately normally distributed. Complete a six-step hypothesis test to test the claim the average systolic blood pressure is less after receiving the drug at $\alpha = 0.05$.

Patient	1	2	3	4	5	6
BP Before (mmHg)	150	160	155	162	148	170
BP After (mmHg)	140	152	148	150	140	160
d_i						
$d_i - \bar{d}$						
$(d_i - \bar{d})^2$						

Inferences Concerning a Difference Between Population Proportions

We can complete a hypothesis test for the difference in proportions.

$$z = \frac{(\bar{p_1} - \bar{p_2}) - D_0}{\sqrt{(\hat{p} \cdot (1 - \hat{p}) \cdot (\frac{1}{n_1} + \frac{1}{n_2})}}$$

$$\hat{p} = \frac{n_1}{n_1 + n_2}\bar{p_1} + \frac{n_2}{n_1 + n_2}\bar{p_2}$$

We can also create a confidence interval for the difference in proportions.

$$(\bar{p_1} - \bar{p_2}) \pm z_{\alpha/2} \cdot \sqrt{\frac{\bar{p_1}(1 - \bar{p_1})}{n_1} + \frac{\bar{p_2}(1 - \bar{p_2})}{n_2}}$$

A study is conducted to compare the proportion of college students who binge drank in the last year at two campuses: CSU-Fort Collins and CSU-Pueblo. A random sample of 80 students from CSU-Fort Collins found that 44 of them reported binge drinking in the past year. Meanwhile, a sample of 75 students from CSU-Pueblo found that 45 of them reported binge drinking.

Create a 90% confidence interval for the difference in proportions of binge drinking between the two campuses. Additionally, test whether there is significant evidence to suggest that one campus has a higher proportion of binge drinkers than the other.

Statistical Power

Statistical power is an essential concept in hypothesis testing, representing the probability that a test correctly rejects a false null hypothesis. In other words, it quantifies how likely we are to detect an effect when there is one.

Definitions

- **Type I error (α)**: The probability of rejecting the null hypothesis when it is actually true. For instance, if $\alpha = 0.05$, there is a 5% chance of incorrectly rejecting the null hypothesis when it is true.
- **Type II error (β)**: The probability of failing to reject the null hypothesis when it is false.
- **Statistical Power**: Defined as the probability of rejecting the null hypothesis when it is false, i.e., correctly detecting a true effect.

$$\text{Power} = 1 - \beta = P(\text{Reject } H_0 \mid H_0 \text{ is false})$$

Power depends on several factors:

- The effect size (how large the true difference is between the groups).
- The sample size (larger sample sizes increase power).
- The significance level (α).
- The variability in the data (lower variability increases power).

Consider a study to test the effectiveness of a new online learning platform for improving exam scores. The platform's developers claim that it will increase the average exam score by 5 points compared to traditional learning methods. Previous studies show that the standard deviation of exam scores is 10 points. You decide to conduct a study with 50 students in each group (online and traditional learning).

- **Null Hypothesis (H_0)**: The mean exam score is the same for both groups.
- **Alternative Hypothesis (H_a)**: The mean exam score for the online group is higher by 5 points.

We want to know the statistical power of detecting this difference if it exists, using an online power calculator instead of hand calculations.

Step-by-Step Power Analysis

1. **Determine effect size**: Using Cohen's d, the effect size can be calculated as:

$$d = \frac{\mu_1 - \mu_2}{\sigma} = \frac{5}{10} = 0.5$$

   This represents a moderate effect size.

2. **Choose significance level (α)**: Set $\alpha = 0.05$ for a 5% risk of Type I error.

3. **Calculate Power in R**: Rather than performing hand calculations, you can input these parameters into R. Simply provide the effect size, sample size, significance level, and test type (two-sample t-test).

4. **Interpret the results**: R will output the statistical power. For this example, the power may be around 0.75, meaning there is a 75% chance of detecting a 5-point difference in exam scores if the platform truly has this effect.

Cohen's d and Effect Size Guidelines

- **Small effect size**: $d < 0.2$
- **Moderate effect size**: $0.2 < d < 0.8$
- **Large effect size**: $d > 0.8$

Key Takeaways

- **Larger sample sizes** increase statistical power, as they provide more data to detect true differences.
- **Larger effect sizes** (i.e., bigger differences between the groups) also increase power.
- **Smaller variability** (lower standard deviation) enhances the power of a test by making the effect easier to detect.

A tech company has released a new version of their app and wants to compare user engagement time (in minutes) between the new version and the old version. Suppose the standard deviation of engagement time is 20 minutes, and they expect the new version to increase engagement by 10 minutes. They plan to randomly select 100 users for each version.

- **Null Hypothesis** (H_0): There is no difference in engagement time between the two versions.
- **Alternative Hypothesis** (H_a): Engagement time is higher for the new version by 5 minutes.

Using R, determine the probability of detecting this 5-minute increase with 100 users per group at $\alpha = 0.05$, that is, calculate the statistical power.

```r
library(pwr)

# Parameters
effect_size = 0.25  # Given effect size
n1 = 100  # Sample size for Group 1
n2 = 100  # Sample size for Group 2
alpha = 0.05  # Significance level

# Power calculation
pwr.t2n.test(n1 = n1, n2 = n2, d = effect_size,
                      sig.level = alpha, alternative = "greater")
```

```
##
##      t test power calculation
##
##              n1 = 100
##              n2 = 100
##               d = 0.25
##       sig.level = 0.05
##           power = 0.5465182
##     alternative = greater
```

R Companion for Chapter 9

The dataset **chickwts** is included in the base version of R. This dataset include 71 observations of the growth rate of chickens using a variety of feed supplements.


```
data(chickwts)

# pick out some random indices of the dataset and look at the elements
set.seed(2020)
chickwts[sample(x=71,size=10,replace = F),]
```

```
##    weight      feed
## 28    250   soybean
## 22    271   linseed
## 65    318    casein
## 17    148   linseed
## 36    248   soybean
## 42    226 sunflower
## 49    325  meatmeal
## 45    334 sunflower
## 56    242  meatmeal
## 8     124 horsebean
```

We can create a boxplot of the weights for the various types of feed.

```
boxplot(weight~feed,data=chickwts)
```



Based on the boxplot, there appears be a difference in weights between **casein** and **horsebean**. To test this, let's run the hypothesis test $H_0 : \mu_1 - \mu_2 = 0$ vs. $H_a : \mu_1 - \mu_2 \neq 0$, where μ_1 is the true mean weight of chicks given casein and μ_2 is the true mean weight of chicks given horsebean. We will use $\alpha = 0.05$ for the following examples.

```
# create new dataset of chicks given casein
chicks1 = chickwts[chickwts$feed=="casein",]
chicks2 = chickwts[chickwts$feed=="horsebean",]
t.test(chicks1$weight,chicks2$weight,mu=0)
```

```
##
##  Welch Two Sample t-test
##
## data:  chicks1$weight and chicks2$weight
## t = 7.3423, df = 18.36, p-value = 7.21e-07
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  116.6982 210.0685
## sample estimates:
## mean of x mean of y
##  323.5833  160.2000
```

The test statistic is $t = 7.34$ and corresponds to p-value of $7.21 \cdot 10^{-7}$. Since the p-value is less than $\alpha = 0.05$, we would reject H_0 and conclude that the true mean chick weights differ between casein and horsebean feed.

Suppose we want to test if the true mean weight of casein feed chicks is more than 200 units greater than the true mean weight of horsebean feed chicks, i.e., $H_0 : \mu_1 - \mu_2 = 200$ vs. $H_a : \mu_1 - \mu_2 > 200$.

```
t.test(chicks1$weight,chicks2$weight,mu=200,alternative = "greater")
```

```
##
##  Welch Two Sample t-test
##
## data:  chicks1$weight and chicks2$weight
## t = -1.6455, df = 18.36, p-value = 0.9416
## alternative hypothesis: true difference in means is greater than 200
## 95 percent confidence interval:
##  124.8371      Inf
## sample estimates:
## mean of x mean of y
##  323.5833  160.2000
```

The test statistic for this hypothesis test is $t = -1.6455$ with a p-value of 0.9416. At $\alpha = 0.05$, we would fail to reject $H_0 : \mu_1 - \mu_2 = 200$ and conclude there is not sufficient evidence to say that the true mean weight of casein feed chicks is more than 200 units greater than the true mean weight of horsebean feed chicks.