

Cell-cycle-gated feedback control mediates desensitization to interferon

Edmund Miller

2021-04-22 Thu

Introduction

Hypothesis

- Does pretreatment with IFN- α result in desensitization?
- Can a computational model be created to predict IFN responses?
- What role does the cell cycle play in the IFN response?

Results

IFN- α pretreatments confer
opposite effects depending on
their durations

Fig 1A. HeLa Reporter Cell Line

_20210422_091025screenshot.png

Fig. 1D Schematic of IFN- α pretreatment experiments

_20210422_104851screenshot.png

Fig. 1D A diagram of the microfluidic set-up

_20210422_105123screenshot.png

Fig. 1B Time Lapse of Cells treated for IFN- α for 48 hours

_20210422_091431screenshot.png

Fig. 1B Time traces of nuclear/cytoplasmic STAT1-mCherry and PIRF9-YFP signals of the cell

_20210422_091808screenshot.png

Fig. 1C Averaged time traces of nuclear/cytoplasmic STAT1-mCherry, PIRF9-YFP

_20210422_100656screenshot.png

Fig. Supp 1D Time course western blots showing the dynamics of phosphorylation (pY701), IRF9 and expression of STAT1

_20210422_093026screenshot.png

Fig. 1E PIRF9-driven YFP induction response to the second IFN- α treatment

_20210422_102211screenshot.png

USP18 is responsible for
desensitization induced by the
prolonged IFN- α pretreatment

Fig. 2A Time-lapse images of STAT1 nuclear translocation

_20210422_112910screenshot.png

Fig. Supp. 2A Western blots of USP18 expression in WT and USP18-KD cells

_20210422_121123screenshot.png

Fig. 2B YFP induction response to the second IFN- α treatment in USP18-KD

_20210422_111043screenshot.png

Fig. 1F Amounts of PIRF9-YFP induction by the second IFN- α stimulation

_20210422_111453screenshot.png

Fig. 2C Amounts of PIRF9-YFP induction in USP18-KD cells by the second IFN- α stimulation

_20210422_111412screenshot.png

Computational modeling
suggests a delayed negative
feedback loop through USP18

Fig. 3A Simple kinetic model of the IFN-driven gene regulatory network

_20210422_113839screenshot.png

Fig. Supp 3A kinetic model of the IFN-driven gene regulatory network with parameters

_20210422_115407screenshot.png

Fig. 3C Amounts of PIRF9-YFP induction by the second IFN- stimulation Predicted by model simulations

_20210422_114424screenshot.png

Fig. Supplement 3B Model fitting results

_20210422_115253screenshot.png

Fig 3D. Experimental design with repetitive IFN pulses

_20210422_115744screenshot.png

Fig. 3E Model prediction of the responses to pulse versus sustained IFN inputs

_20210422_115809screenshot.png

Fig. 3F Experimental data of the responses to pulse versus sustained IFN inputs

_20210422_115854screenshot.png

The kinetics of USP18
upregulation by IFN is
heterogeneous in single cells

Fig. 4A Dual reporter cell line schematic

_20210422_120826screenshot.png

Fig. 4B Time traces of PIRF9-YFP and PUSP18-CFP of a single cell in response to IFN-

_20210422_120948screenshot.png

Fig 4C. Distributions of PIRF9 and PUSP18 activation times in single cells

_20210422_121313screenshot.png

Fig 4D. Distributions of delay times in single cells

_20210422_121358screenshot.png

Fig 4E. Representative time traces of PIRF9 and PUSP18 in a single cell from each group

_20210422_121444screenshot.png

Cell cycle phases differentially
regulate USP18 expression

Fig 4E. Delay times as a function of the percentages of cell cycle progression upon IFN treatment onset

_20210422_122543screenshot.png

Fig 5A. Delay times in cells treated with different cell cycle perturbation

_20210422_121847screenshot.png

Fig 5B. CDK2 activity reporter Schematic

_20210422_122704screenshot.png

Fig 5C. Nuclear DHB and PUSP18-driven gene expression

_20210422_122941screenshot.png

Fig 5D. Effect of decitabine on DNA methylation and nucleosome occupancy

_20210422_123028screenshot.png

Fig 5D. Distribution of delay times upon decitabine treatment

_20210422_123104screenshot.png

Cell-cycle-gated feedback control shapes single-cell responses to repetitive IFN inputs

Fig. 6A simple model of the IFN-driven gene regulatory network

_20210422_123240screenshot.png

Fig. 6C cell-cycle gated feedback control simulated responses under different pretreatment conditions

_20210422_123535screenshot.png

Fig. 6D cell-cycle gated feedback control experimental responses under different pretreatment conditions

_20210422_123547screenshot.png

Higher levels of USP18 expression by the prolonged pretreatment lead to reduced IRF9 induction upon the second stimulation at the single-cell level, qualitatively in agreement with our experimental data

Conclusion

Conclusion

- The effects of IFN pretreatments depend on their input durations
- The G1 and early S phases enable an open window for immediate USP18 upregulation upon the IFN treatment
 - If they miss the window the USP18 induction has to wait for G1 of the next cell cycle
- SARS-CoV-2 is especially sensitive to type I IFNs
 - IFN pretreatment a potential strategy to prevent SARS-CoV-2 infection