

Course

Course > Week 2: Perceptrons > Week 2 Exercises > Week 2 Exercises: Perceptrons

< Previous



Next >

Week 2 Exercises: Perceptrons

Bookmark this page

Week 2 Exercises: Perceptrons

(1.0 / 1.0 points)

For these exercises, it will be helpful to review the notes on [Linear Classifiers](#) and the [Perceptron](#). You may also find it helpful to write some test code with a local python installation or in a [google colab notebook](#).

1) Classification

Consider a linear classifier through the origin in 4 dimensions, specified by

$$\theta = (1, -1, 2, -3)$$

Which of the following points x are classified as positive, i.e. $h(x; \theta) = +1$?

1. $(1, -1, 2, -3)$
2. $(1, 2, 3, 4)$
3. $(-1, -1, -1, -1)$
4. $(1, 1, 1, 1)$

Enter a Python list with a subset of the numbers 1, 2, 3, 4:

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

2) Classifier vs Hyperplane

Consider another parameter vector

$$\theta' = (-1, 1, -2, 3)$$

Ex2a

Does θ' represent the same hyperplane as θ does?

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

Ex2b

Does θ' represent the same classifier as θ does?

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

3) Linearly Separable Training

As [discussed in lecture and in the lecture notes](#), note that $\mathcal{E}_n(\theta, \theta_0)$ refers to the training error of the linear classifier specified by θ, θ_0 , and $\mathcal{E}(\theta, \theta_0)$ refers to its test error. What does the fact that the training data are *linearly separable* imply?

Select "yes" or "no" for each of the following statements:

Ex3a

There must exist θ, θ_0 such that $\mathcal{E}(\theta, \theta_0) = 0$

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

Ex3b

There must exist θ, θ_0 such that $\mathcal{E}_n(\theta, \theta_0) = 0$

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

Ex3c

A separator with 0 training error exists

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

Ex3d

A separator with 0 testing error exists, for all possible test sets

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

Ex3e

The perceptron algorithm will find θ, θ_0 such that $\mathcal{E}_n(\theta, \theta_0) = 0$

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

4) Separable Through Origin?

Provide two points, (x_0, x_1) and (y_0, y_1) in two dimensions that are linearly separable but not linearly separable through the origin. If you get stuck try drawing a picture and review the notes on [offsets](#).

Enter a Python list with two entries of the form `[[x0, x1], label]` where label is 1 or -1. (So each entry represents a point with 2 dimensions and its label)

Submit


View Answer

100.00%

You have infinitely many submissions remaining.

This page was last updated on Saturday January 18, 2020 at 01:16:11 PM (revision 043d9ae).

Sec None Release 1900-01-01 00:00:00 Due 9999-12-31 23:59:59



Powered by CAT-SOOP v14.0.0.dev144.

CAT-SOOP is [free/libre software](#), available under the terms of the [GNU Affero General Public License, version 3](#).

[Download Source Code](#)

[Javascript License Information](#)

< Previous

Next >