

Week 3 Exercises: Features

 [Bookmark this page](#)

Week 3 Exercises: Features

(1.0 points possible)


For these exercises, it will be helpful to review the notes on [Linear Classifiers](#)

1) Feature representation

For the following feature, pick what might be the best encoding for linear classification. The assumption is that there are other features in the data set.

The point of this question is to think about alternatives; there are many options, many not mentioned here.

Car make, e.g. Chevy, Ford, Toyota, VW, for predicting gas mileage (lo, hi).

4 unary features (one-hot): 1000, 0100, 0010, 0001 

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

2) Feature mapping

Consider the following, one-dimensional, data set. It is not linearly separable in its original form.

- 1.  $x^{(1)} = -1, y^{(1)} = +1$
- 2.  $x^{(2)} = 0, y^{(2)} = -1$
- 3.  $x^{(3)} = 1, y^{(3)} = +1$

**Ex2.a.:** Which of these feature transformations leads to a separable problem?

- 1.  $\phi(x) = 0.5 * x$
- 2.  $\phi(x) = |x|$
- 3.  $\phi(x) = x^3$
- 4.  $\phi(x) = x^4$
- 5.  $\phi(x) = x^{2k}$  for any positive integer  $k$

Enter a Python list with a subset of the numbers 1, 2, 3, 4, 5.

Submit

View Answer

100.00%

You have infinitely many submissions remaining.

**Ex2.b.:** Your friend Kernelius uses feature transformation  $\phi(x) = (x, x^2)$  on the data above. In the new space, the linear classifier with  $\theta = (0, 1)$  and  $\theta_0 = -0.25$  achieves perfect accuracy. What points from the original space  $\mathbb{R}$  map to this linear classifier in  $\mathbb{R}^2$ ? (It may be helpful to find the equation of the separator.)

Enter a Python list with all values of  $x$  which constitute this separator.

You have infinitely many submissions remaining.


Solution: [-0.5, 0.5]

**Explanation:**

With  $\theta = (0, 1)$  and  $\theta_0 = -0.25$ , our separator is the line  $y = 0.25$  in  $\mathbb{R}^2$ . From our original space  $\mathbb{R}$ , points mapped onto the separator are those for which  $x^2 = 0.25$ , leading to  $x = \pm 0.5$

This page was last updated on Saturday January 18, 2020 at 01:16:11 PM (revision 043d9ae).

Sec None Release 1900-01-01 00:00:00 Due 9999-12-31 23:59:59



Powered by [CAT-SOOP](#) v14.0.0.dev144.

CAT-SOOP is [free/libre software](#), available under the terms of the [GNU Affero General Public License, version 3](#).

[Download Source Code](#)  
[Javascript License Information](#)