MAGNETIC-A: COMPREHENSIVE WEB-BASED TOOLBOX FOR PALEOMAGNETIC ANALYSES



VERSION 2.0

By Edoardo Dallanave

Department of Earth Sciences, University of Milan (20133 Milan, Italy)

ALP-CIMaN Alpine Palaeomagnetic Laboratory (12016 Peveragno, Italy)

Email: edoardo.dallanave@unimi.it
Personal webpage: https://edoardodallanave.wixsite.com/mysite
Repository: https://doi.org/10.5281/zenodo.18220000

Table of Contents

MAGNETIC-A 3
1. Starting with Magnetic-A: Web version and local version 3
1.1. *Web version* 3
1.2. *Run Magnetic-A locally* 3
2. Directions analysis 4
2.1. *Vector end-points interpolation* 4
Figure 2.1. Vector end-points interpolation main window. 4
2.2. *Export figure with all diagrams* 5
Figure 2.2: window showing all details of the sample selected in the "Vector end-points interpolation" panel. 6
2.3. *All saved directions* 6
Figure 2.3. Window showing all saved interpolated directions and planes. 7
3. Directions display & average 7
3.1. *Directions display & average* 7
Figure 3.1. Directions display & average window. 7
3.2. *Plot multiple directions sets* 9
Figure 3.2. Plot multiple directions sets window. 9
3.3. *Add average directions* 10
Figure 3.3. Add average directions window. 10
Figure 3.4. Pop-up window for manual entry of average direction and 95% confidence. 11
4. Non-parametric statistics 11
4.1. *Confidence ellipses* 11
Figure 4.1. Confidence ellipses sub-page of the bootstrap statistics. 12
4.2. *Reversal test* 12
Figure 4.2. Bootstrap test for a common mean direction (aka reversal test) sub-page. 12

4.4. Fold test ... 13
Figure 4.3. Bootstrap-based fold test. ... 13

5. Distribution reliability (SVEI test) ... **14**

5.1. SVEI test ... 14
Figure 5.1. SVEI test for consistency with the THG24 GGP model. ... 14

5.2. Inclination flattening estimate ... 15
5.2.1. Suggested routine for the inclination flattening test ... 15
Figure 5.2. Preliminary (explorative) paleomagnetic inclination flattening detection. ... 15
Figure 5.3. Detailed inclination flattening calculation process. ... 16
Figure 5.4. The f value is applied to the original dataset. ... 16

6. TK03.GAD E/I Test ... **16**

Figure 6.1. TK03.GAD-based inclination shallowing detection and correction. ... 17

7. VGP and Pmag Poles ... **17**

Figure 7.1. VGP and Pmag Poles page. The red rectangles envelop the commands for the projection details, while the blue rectangle envelops the Action Buttons. ... 18

7.1. Projection commands ... 18
7.2. Action buttons ... 18
7.2.1. Internal ... 18
Figure 7.2.1. Internal VGPs pop-up window ... 19

7.2.2. External ... 19
Figure 7.2.2. External VGPs pop-up window. ... 20

7.2.3. Simulate ... 20
Figure 7.2.3. Fisherian VGPs simulator pop-up window. ... 21

7.2.4. Merge ... 21
Figure 7.2.4. VGPs Merging tool pop-up window. ... 22

7.2.5. Pmag Pole ... 22
Figure 7.2.5. Add standard paleomagnetic pole pop-up window. ... 22

7.2.6. Rotate ... 23
Figure 7.2.6. Rotate VGPs and Poles pop up window. ... 23

7.2.7. Fisher A95 ... 23
Figure 7.2.10. Pop-up window allowing the Fisher mean and associated A95 calculation of the entries selected in the list of loaded VGPs and Poles. ... 23

7.2.8. APWP ... 24
Figure 7.2.7. Apparent polar wander pop-up window ... 24

7.2.9. Locality ... 24
Figure 7.2.8. Pop-up window showing the plotting details and the list of the loaded localities. ... 24

7.2.10. Small circle ... 24
Figure 7.2.9. Pop-up window showing the plotting details and the list of the loaded circles. ... 25

7.2.11. Great circle ... 25
Figure 7.2.11. Pop-up window with the details of the great circle interpolating the selected geographic points (VGPs average, poles, localities). ... 25

7.2.12. Other commands within the VGPs and Pmag Poles window ... 25

8. Magnetic polarity ... **26**

Figure 8.1. Magnetic polarity stratigraphy page ... 26

9. Site map ... **27**

Figure 9.1. Site map page. ... 27

Glossary ... **27**
References ... **28**
Appendix 1: Demagnetization data file accepted formats ... **30**
Appendix 2: Average directions and 95% confidence files format ... **31**
Appendix 3: Custom APWP file format ... **33**
Appendix 3: Direction file formats for plotting magnetostratigraphy ... **34**
Appendix 4: Sites file for mapping tool ... **34**

MAGNETIC-A

This manual introduces the user to the main functionalities of Magnetic-A, describing the calculations and plotting options. Figures are designed to be easily downloaded as vector pdf files and used with minimal manipulation. Help buttons (⃝) are present in different key locations of the pages: pop-up window will open with useful information and references.

1. Starting with Magnetic-A: Web version and local version

1.1. Web version

Magnetic-A is compiled using the R-Package Shiny, specifically designed to develop web-apps. The web version of Magnetic-A is deployed in the shinyapps server (https://www.shinyapps.io/). To access Magnetic-A, please navigate your browser to either:

- https://edoardodallanave.shinyapps.io/MagneticA/

or alternatively:

- https://edoardodallanave.shinyapps.io/MagneticA2/

1.2. Run Magnetic-A locally

If an extensive use of Magnetic-A is envisaged, it is advantageous to run Magnetic-A locally (faster bootstrap, no risk of losing connection). To do so, download and install both R and RStudio:

- R = https://www.r-project.org/
- RStudio = https://www.rstudio.com/products/rstudio/download/

Magnetic-A requires a series of packages that need to be installed only once. Most of the packages are stored in the CRAN, which makes it easy to proceed. Type the following commands in the RStudio console and press return after each:

- install.packages("plyr")
- install.packages("dplyr")
- install.packages("shiny")
- install.packages("shinyWidgets")
- install.packages("DT")
- install.packages("shinyhelper")
- install.packages("shinyjqui")
- install.packages("glue")
- install.packages("tidyverse")

The last (but essential!) package is called PmagDiR (Dallanave, 2024) and it is stored in GitHub. The easiest way to install it is to use a package called “devtools”. Type the following commands:

- install.packages("pak")
- pak::pak(“edoardo-paleomag/PmagDiR”)

PmagDiR is often updated. Just check for updates in GitHub, and type the last two commands for installing updated versions.

Once all of these is done, you can run Magnetic-A locally by typing:

- shiny::runGitHub("Magnetic-A", "edoardo-paleomag")

Alternativley, you can run Magnetic-A offline after pulling the GitHub content in a folder of your local device and type:

- shiny::runApp("/Path_to_the _folder/Name_of_the_folder/")

The instructions are also available in https://github.com/edoardo-paleomag/Magnetic-A/blob/main/README.md or https://doi.org/10.5281/zenodo.18220000

2. Directions analysis

2.1. Vector end-points interpolation




Figure 2.1. Vector end-points interpolation main window.

This page is designed to isolate paleomagnetic components of the NRM. VEPs can be selected by either dragging the mouse onto the diagram, or by selecting the demagnetization steps on the right-hand side of the screen. If you want to preserve the selection also after shifting to a different coordinate system or projection, then you are recommended to select the points by clicking on the list (dragged selection is otherwise lost). Selected VEPs are highlighted by different color: projection of VEPs onto the horizontal plane change from black to red, while projection onto the vertical plane change from white to yellow. The same applies for the equal area projection, but in this case black and white symbols correspond respectively to projection onto the lower and upper hemisphere of the diagram.

To operate the software:

Load demag data: click on it to upload the demagnetization data file from your device.

File type: different data formats are accepted. Please refer to the Appendix 1 of this manual to check examples of data files. The file format currently accepted are also described in the help pup-up window shown by clicking on the help button. By selecting "Example data" at the bottom of the list, some real examples of VEPs data from Dallanave et al. (2012) are available to familiarize with the functions. The list can be updated with new formats upon request.

Diagram: select the preferred projection between N-Right (north axis point to the right of the screen), N-Up (north axis points up), and equal area diagram.

Coordinates: select the preferred VEPs projection coordinates between Specimen (data are plotted in the sample coordinate system), Geographic (data are plotted with respect to the geographic reference system), and Tilt Corrected (Tilt Corr. – Data are plotted restoring the original horizontality of the strata). The data in the three different projections are taken from the demagnetization file uploaded, so pay attention to the file format (Appendix 1) to avoid any problem.

UNITS: this button opens a pop-up window where the user can define the demagnetization unit, NRM unit, volume and mass of the specimens if normalization is required. For some formats (e.g. LASA), the window is pre-compiled, but all values can be updated.

TAGS: this button opens a pop-up window with the labels spacing and position. It is advised to set all parameters as desired before exporting the figure (sub-page 1.3, see below) for publication purposes.

Interpolation: The selected VEPs can be interpolated using different methods. These are:

- PCA Free: PCA free from origin of the demagnetization axes.
- PCA Anch.: PCA anchored (*sensu* Kirschvink, 1980).
- PCA Or. Incl.: the origin of the demagnetization axes is included as VEP.
- PCA Constr.: PCA constrained (sensu Heslop & Roberts, 2016).
- Fisher: VEPs are averaged by using standard spherical statistics (Fisher, 1953).
- G. Circle: VEPs are interpolated by a plane (Kirschvink, 1980) expressed as pole (*i.e.*, declination and inclination of the line perpendicular to the plane).

Component name: The name associated to each component saved (multiple NRM components can be saved during the same session!).

List of loaded specimens: interactive list of the specimens loaded; by selecting a specimen, all operation options are activated.

Delete selection: The selected VEPs will be removed from the figure.

Restore data: All VEPs will be restored and the current interpolation cancelled (this does not affect interpolation already saved).

Run Probabilistic PCA test: this button performs the Bayesian model selection introduced by Heslop & Roberts (2016) which provide the probability that the selected VEPs support the use of an anchored, unanchored, or constrained PCA solution. Please cite the original Heslop & Roberts (2016) paper if using this method (recommended).

READ CARFULLY: Generally speaking, the use of **PCA anchored** is generally **WARMLY DISCOURAGED** (unless justified by the probabilistic PCA test) as it results in **ARTIFICIALLY LOW MAD**. VEPs failing to trend linearly toward the origin of the demagnetization axes should be interpolated with Fisher (1953) mean.

Run interpolation: performs the VEPs interpolation. The interpolation line is plotted on the diagram as blue dotted line.

Save interpolation: save the interpolated components in the list of directions (interactive list in "All saved samples" page).

Export all saved directions: export all saved directions in your local device.

READ CARFULLY: Magnetic-A DOES NOT SAVE AUTOMATICALLY the interpolated directions in your personal device. It is **WARMLY RECOMMENDED** to download regularly the saved directions to avoid annoying loss of data.

2.2. Export figure with all diagrams

In this page a figure with all diagrams, units, and the interpolated component (if present) is shown. By clicking on *Export figure* the figure is downloaded as editable vector file (pdf) (Figure 1.2). The name of the exported file is the name of the selected sample.




Figure 2.2: window showing all details of the sample selected in the "Vector end-points interpolation" panel.

2.3. All saved directions

In this sub-page you find an interactive list of all saved directions. All directions selected in the table will be displayed in the equal area diagram on the main panel (Figure 1.3). The table is interactive, so any parameter can be used to sort the list by clicking on the header. For example, clicking on the header of the last column will sort the directions by component name, allowing to select and display more easily components with the same name.

Import saved directions: upload a file as exported from the "*Vector end-points interpolation"* page. This allows to work on data previously isolated using Magnetic-A.

Name of exported file: selected (highlighted in blue) directions can be exported as a new (.csv) file in your personal device, and the name of the exported file can be typed here.

Coordinates: in this diagram, directions can be displayed by using a geographic or a tilt corrected coordinate system frame.

Delete selection: Selected directions will disappear from the interactive list. If directions are from a loaded file, the original file stored in the personal device will stay unchanged.

Undo delete: restores last deleted directions.

Export selected directions: this will download in the local device the selected directions using the "*name of exported file*"; this option is useful to separate the NRM components in different files.

Export figure: equal area diagram is downloaded locally as vector (editable pdf) file.

Combine DI & GC: combines directions and great circle (or only great circle) by using the algorithm of McFadden & McElhinny (1988).

Add GC dirs to list: the directions lying on the great circles (shown as triangles in figure) estimated by using McFadden & McElhinny (1988) are saved as directions and added to the interactive list, indicated by "*Dir*" in the "*Type*" column.

Clear GC dirs from plot: the directions lying on the great circles (triangles) are deleted from the figure (not from the list if already saved).

Show dragged directions details: opens a pop-op window with details of the directions selected by dragging the mouse. This feature is convenient for identify outliers, but the buttons described above (e.g. *Delete selection*) do not operate on the directions selected by mouse-dragging (light blue rectangle in the figure panle).




Figure 2.3. Window showing all saved interpolated directions and planes.

3. Directions display & average

3.1. Directions display & average

This sub-page is designed to work on a set of interpolated directions. Directional data can be plotted with different style, and several statistical options to estimate the average directions are available (Figure 3.1). Directions can be either uploaded as a file or directly imported (internally) from the "*Directions analysis*" page.




Figure 3.1. Directions display & average window.

Load directions file: upload a directions file.

Site name: directions can be exported after processing, and the exported file will have the name typed here.

Directions file type: several (.csv) file formats are accepted (the details are listed within the helper pop-up window). **NOTE**: add a header to the data file, or it will skip the firs data point!

- Dec, Inc: two columns with declination and inclination, and optional third column with stratigraphic position.

- G_dec, G_inc, B_az, B_plunge: four columns in total, first two with declination and inclination in geographic coordinates, third and fourth with bedding (azimuth and dip of plunge; NO STRIKE); optional fifth column with stratigraphic position for magnetic polarity plot.
- G_dec, G_inc, B_dec, B_inc: four columns in total, first two with declination and inclination in geographic coordinates, third and fourth with declination and inclination in tilt corrected coordinates; optional fifth column with stratigraphic position for magnetic polarity plot.
- S_d, S_i, G_d, G_i, B_d, B_i: six columns in total, first two with declination and inclination in specimen coordinates, third and fourth with declination and inclination in geographic coordinates, fifth and sixth with declination and inclination in tilt corrected coordinates; optional seventh column with stratigraphic position.
- Magnetic-A: file as exported from the Directions analysis page.
- Internal file: all directions selected within the "All saved samples" sub-page.
- Example file: paleomagnetic ChRM directions from in geographic and tilt corrected coordinates, with associated stratigraphic position for magnetic polarity determination (see below), from Dallanave et al. (2012).

Coordinates: preferred reference frame (specimen, geographic, tilt corrected). Selection depends on the input file! Make sure to select the appropriate file type.

Site latitude, Site longitude: these are used in the next pages for the VGPs calculation and site map. Make sure to type in the proper coordinates if you want to use the "*VGP and Pmag Poles*" pages.

Flattening: a flattening factor ranging from 0 to 1 (applied to the directions by using tangent function of King, 1955) can be typed in and used to correct directions that are already known to be affected by paleomagnetic inclination shallowing.

Mean: the average direction and 95% confidence angles can be estimated by using different approaches.

- Fisher: calculate (Fisher, 1953) average and 95% cone of confidence (a95); if a distribution is bimodal, statistic is calculated automatically for the two modes, and for all directions flipped onto a common mode.
- Fisher (no split): same as above but without splitting a bimodal distribution. This is convenient for random distributions that the User wants to test for randomness (see *Watson's test of Randomness* button below).
- Elliptic: calculate average direction and 95% ellipses defined by the two semi-angles of confidence (a95dec and a95inc) as described by Deenen et al. (2011). Bimodal distributions are automatically split.
- Inc. only (single mode and bimodal): specifically designed for directions from azimuthally unoriented deep-sea drilled cores, provides average inclination and 95% confidence calculated by using the maximum likelihood solution of Arason & Levi (2010). *Single mode* considers all data as a single distribution. *Bimodal* split the directions in up-pointing and down-pointing.
- Arithmetic (single mode and bimodal): similarly to Inc. only, it calculates the arithmetic means, useful when the maximum likelihood solution fails to provide an estimate (see Arason & Levi, 2010).

Mode: if directions are originally split in two clusters (e.g., normal and reversed polarity) or are generically divided in up-pointing and down-pointing, different options can be applied. Bimodal distributions are evaluated by using the same computational background of PmagDiR (Dallanave, 2024).

- Bimodal: directions are plotted in their original form.
- Mode 1: all directions are flipped toward Mode 1, which is considered the one with down-pointing average direction.
- Mode 2: all directions are flipped toward Mode 2, which is considered the one with up-pointing average direction.
- All down: all directions flipped as down-pointing
- All up: all directions flipped as up-pointing.

Apply f: if yes, it applies to the directions the flattening correction factor typed in the "*Flattening*" box.

Color down; color up; symbol: colors and symbols of choice can be used.

GAD: adds to the plot the geocentric axial dipole direction (and circle if selected) for the latitude typed in the "*Site Latitude"* box.

Cut-off: applies different types of cut-offs to the direction distribution. The cut-offs defined as "*dynamic*" include in the reiterative process the correction for inclination flattening by using the TK03.GAD-based method (Tauxe & Kent, 2004). Although this method is now superseded by the recent SVEI method (Tauxe et al., 2024), inclusion in the cut-off reiterative of the TK03.GAD method provide a less-strict cut-off result. Details on this concept are provided in Dallanave (2024). *"Static"* cut-off does not include any inclination flattening evaluation.

READ CARFULLY: Be aware that for comparing the distribution with what expected from GGP model THG24 (Tauxe et al., 2024), it is recommended **NOT TO APPLY ANY CUT-OFF**. The explanation is provided in Tauxe et al. (2024).

- Vandamme: applied the cut-off proposed by Vandamme (1994).
- Fixed: applies a cut-off based on the VGP angular distance from the average paleomagnetic pole; the cut-off maximum angle is defined in the "*VGP filter-fixed radius*" box (default= 45°).

- Cut up-(down-) pointing: cuts all directions pointing either up or down.
- Cut fixed inc.: specifically designed for directions from azimuthally unoriented deep-sea drilled cores, it cuts all directions comprised within the angular interval defined in the "*Min inc. filt.*" and "*Max inc. filt.*" boxes (default values are both zero).

Export graph: download the equal area diagram as editable vector (pdf) file.

Export stat: download the table (.csv format) with the statistical parameters.

Export directions: download the directions as displayed in the equal area in a single file, declination and inclination in two columns (.csv format).

Watson's test of randomness: perform the Watson's test of randomness (Watson, 1956) to check for random distributions at a 95% confidence values. Results are shown in a pop-up window.

Delete dragged directions: directions selected within the figure panel are deleted.

Restore all directions: any deleted direction is restored.

3.2. Plot multiple directions sets

This sub-page is designed to plot different sets of paleomagnetic directions on the same equal-area diagram (Figure 3.2). The directions already loaded and processed in the "*3.1. Directions display & average"* can be imported directly by clicking on the *"ADD MAIN SET TO LIST",* while any other (external) set can be uploaded and plot. This sub-page is plot-only, for processing and averaging, please use the dedicated pages.



Figure 3.2. Plot multiple directions sets window.

Load directions file: upload a directions file. Only one format accepted: two columns (csv file) with header, with declination and inclination.

GAD Latitude: Type in the latitude for plotting the GAD direction.

GAD: adds to the plot the geocentric axial dipole direction (and circle if selected) for the latitude typed in the "*GAD Latitude"* box.

ADD MAIN SET TO LIST: import to the list of dets the directions as processed and plotted in the *"Directions display & average"* sub-page.

DELETE ENTRY: delete the set selected in the table.

The **Table of the loaded sets** is interactive. Selected sets are highlighted in blue, and the symbols and colors are editable.

N: Number of directions

Sym., Col. D., and Col. U.: symbol and color of the down pointing and up pointing directions.

Valid symbols: c= circle; d= diamond, t= triangle, s= square

3.3. Add average directions

This sub-page adds loaded average directions to the *"Plot multiple directions sets"* equal area (Figure 3.3). Also the non-parametric confidence ellipse of Heslop et al. (2023), as determined in the *"Bootstrap statistics"* page, can be here uploaded and plotted.




Figure 3.3. Add average directions window.

Load parametric file: this is used to upload files with average directions and parametric 95% confidence circle (Fisher, 1953) or ellipse (Deenen et al., 2011).

Six file formats are accepted and recognized automatically:

1) Standard (Fisher1953) average and confidence as exported from the "*Directions display & average*" sub-page of Magnetic-A (See Appendix A2.1).
2) Average direction with elliptic confidence (Deenen et al., 2011) as exported from the "*Directions display & average*" sub-page of Magnetic-A (See Appendix A2.2).

DO NOT CHANGE the columns name of the file before importing if using data exported from Magnetic-A!

The remaining 4 formats can be compiled manually, all as .csv file with (any) columns header:

3) Standard Fisher (1953): 3 columns with declination, inclination, a95 (colors and symbol are set automatically) (See Appendix A2.3).
4) Same format as above, but with 3 more columns (total columns= 6), with symbol, color of symbol, color of lines (See Appendix A2.4).
5) Elliptic (Deenen et al., 2011): 4 columns with declination, inclination, a95_dec, a95_inc (colors and symbol are set automatically) (See Appendix A2.5).
6) Same format as above, but with 3 more columns (total columns= 7), with symbol, color of symbol, color of lines (See Appendix A2.6).

Load non-parametric file: this is used to upload files as exported from the *"Confidence ellipse"* sub-page of the *"Bootstrap statistics"* page.

Two formats are accepted and recognized automatically:

1) 4 columns (csv), with (any) header. Columns 1 & 2, only one cell each, with averaged declination and inclination; columns 3 & 4 with declination and inclination of the 95% confidence ellipse points (no limit in the number of rows).
2) 8 columns (csv), with any header. Same as above, with 2 average directions (Dec1, Inc1, Ellips_dec1, Ellips_inc1, Dec2, Inc2, Ellips_dec2, Ellips_inc2

Add entry manually: this button opens a pop-up window to insert manually any average directions with either standard or elliptical 95% confidence (Figure 3.4).




Figure 3.4. Pop-up window for manual entry of average direction and 95% confidence.

4. Non-parametric statistics

This page performs non-parametric statistical analysis of the distribution. The first page calculates the 95% confidence ellipses around the average direction(s), while the second page performs the bootstrap-based reversal test (Heslop et al., 2023).

READ CAREFULLY: Please cite Heslop et al. (2023) if using this page for publication (the link to the publication is embedded in the bottom left of the page).

4.1. Confidence ellipses

This sub-page works on the directions loaded in the "*Directions display & average*" page. To perform the calculation of the non-parametric confidence ellipses, at least 10 directions (per mode, if distribution is bimodal) must be present. If not, a warning message will appear, and the simulation does not start (Figure 3.1).

Export name: name of the result file and graphic file to export.

Bootstrap n.: number of the simulated pseudosamples used to estimate the confidence ellipses. Default is 10.000 as suggested by Heslop et al. (2023).

Plot Dirs: add or remove the directions from the equal area (not the average and the ellipsis).

Run Mode 1 and *Run Mode 2:* perform the test for the two modes (if present) separately.

Clear plot: remove all directions and confidence.

Export graph: downloads equal area diagram

Export ellipsis: the confidence ellipsis is not defined by parameters but by a series of declination-inclination pairs. This command will generate and download a .csv file containing the average direction of the mode (or both modes if present) and the coordinate of the confidence ellipsis (for both modes if present).




Figure 4.1. Confidence ellipses sub-page of the bootstrap statistics.

4.2. Reversal test

This sub-page performs the bootstrap test for common mean direction of Heslop et al. (2023), applied to the two distribution modes (see e.g. the classic textbooks of Butler, 1992 or Tauxe, 2010 for details on the field tests for paleomagnetic stability).




Figure 4.2. Bootstrap test for a common mean direction (aka reversal test) sub-page.

Export name: name of the result file and graphic file to export.

Bootstrap n.: number of the simulated pseudosamples used to calculate the estimate the confidence ellipses. Default is 10.000 as suggested by Heslop et al. (2023).

Perform: starts test.

Export graph: downloads figure with histogram containing the critical v_3 value, the observed value, and the result of the simulations (Heslop et al., 2023), together with the equal area diagram with both modes plotted on a common polarity.

Export Common direction and ellipsis: if the test is positive, a common direction and the relative 95% confidence ellipsis is generated and exported.

4.3. Common mean direction test (CMDT) of two datasets

This sub-page performs the bootstrap test for common mean direction of Heslop et al. (2023), applied to the set loaded in the "*Directions display & average*" page and a second set loaded in this page

READ CAREFULLY: while the "*Reversal test*" page applies the CMDT to the same dataset assumed bimodal, this page applies the CMDT to two different datasets.

Load second set: upload a csv value with declination and inclination. Magnetic-A will isolate the first two columns of any file uploaded.

Show second set: opens a pop-up window with an equal-area diagram of the second set.

All other buttons are the same as the reversal test sub-page.

4.4. Fold test

This sub-page performs the fold test based on the eigen analysis approach of (Tauxe & Watson, 1994), which gives confidence limits on the degree of unfolding required to produce the tightest grouping of data and requires no assumptions about the polarity or distribution of data (i.e., it works on either single mode or bimodal distributions).




Figure 4.3. Bootstrap-based fold test.

The test is performed on paleomagnetic directions in geographic coordinates coupled with bedding data, expressed as azimuth of dip (i.e., strike +90°) and plunge (down= positive). The paleomagnetic direction file is selected in the *Directions display & average* page. The first format accepted is the *G_dec, G_inc, B_az, B_plunge* format (see paragraph *3.1-Directions display & average* of this manual). Alternatively, the test works on *Magnetic-A* files and *Internal* files (see paragraph *3.1-Directions display & average*). In these two latter cases, a file with sample code, azimuth of dip and plunge must be loaded (three columns, comma separated values, with header).

READ CAREFULLY: the sample codes of the Magnetic-A (or Internal) and the bedding files MUST be the same!

Upload bedding file: this is used to upload the bedding file if necessary. By clicking on the help button, details on the files format are provided.

Unfolding %: select the unfolding range of the test. Default is -10% to 120%. The range can be narrowed down to increase the accuracy of the maximum grouping of the data and the 95% confidence boundaries.

Bootstrap n.: number of bootstrapped pseudosamples.

Perform: Starts the test.

Export graph: the result of the fold test is graphical only. Best grouping and 95% confidence are shown numerically in the figure.

5. Distribution reliability (SVEI test)

This page performs the test for paleomagnetic directions distribution consistency with the THG24 GGP field model (Tauxe et al., 2024) (first sub-page), and the estimation of the inclination flattening for direction distribution derived by sedimentary rocks (second sup-page).

READ CAREFULLY: The original protocol is compiled in Python and published as Jupyter Notebook (Tauxe et al., 2024), and it has been translated into R (with minor modifications) to be applied within Magnetic-A. Please cite the original publication if using this method (the link to the publication is embedded in the bottom left of the page).

5.1. SVEI test

This sub-page compares a given directions distribution (loaded in the "*Directions display & average*" page) with simulated distributions drawn from the THG24 paleosecular variations model (Tauxe et al., 2024) for the same sampling site latitude. The empirical declination and inclination from the distribution are compared with the expected cumulative distribution functions using the Anderson-Darling "goodness-of-fit" test (Anderson & Darling, 1952; please refer to Tauxe et al., 2024 for details on the method). Similarly to the Elongation-Inclination method of Tauxe & Kent (2004), the elongation (E) of the distribution and its orientation, expressed as declination of elongation (E Decl.) are compared to the ones expected from simulated distributions (Tauxe et al., 2024). In the case of Figure 5.1, the directions declination do not fit the expected declination, and the declination of elongation is outside the confidence boundaries defined by the simulated distribution: the test is not passed. For consistency with Tauxe et al. (2024), this example is based on the paleomagnetic directions of Gilder et al. (2001) filtered for MAD≤10, also available at https://www2.earthref.org/MagIC/20098.



Figure 5.1. SVEI test for consistency with the THG24 GGP model.

Export name: name of the graphic file to export.

Kappa: assumed uncertainty. First run the test with kappa=infinite (no site scatter assumed), then 100 and 50 if test fails. For sedimentary records originally sampled for magnetostratigraphy, 50 is preferred (Tauxe et al., 2024).

Simulations: number of the simulated dataset drawn from the THG24 model.

Please cite the original Tauxe et al. (2024) publication if using this test!

5.2. Inclination flattening estimate

This sub-page gradually "unflattens" the paleomagnetic directions following the tangent function of King (1955):

$$\tan I_o = \frac{\tan I_f}{f}$$

Where I_o is the original (unflattened) inclination, I_f = is the flattened direction, and f = flattening factor ranging from 0 to 1. After each unflattening step, it applies the SVEI test for consistency with the THG24 paleosecular variations model. Please refer to Tauxe et al. (2024) for details on the routine.

Export name: name of the graphic file to export.

Simulations: number of the simulated dataset drawn from the THG24 model.

Initial f, Target f, and *Increment:* the series of unflattening *f* values can be compiled by using three intervals (see below for the fast and slow routines)

Check f list: opens a pop-up window with all *f* values that will be applied.

Perform flattening test: starts the test.

5.2.1. Suggested routine for the inclination flattening test

Following what suggested by (Tauxe et al., 2024), before starting the unflattening procedure on data from sedimentary rocks, it should be tested if the data (when bimodal) pass the reversal test (sub-page 3.1), and that they are not already consistent with the THG24 model (sub-page 4.2). After that, an initial inspection of the data can be applied setting *f* values ranging from 1 to 0.3 with 0.05 increments (red-line rectangle in Figure 5.2; analogous of the *Quick= True* option in the original Jupyter notebook of Tauxe et al. (2024)). The pink areas in all diagrams bracket the range of *f* where the data show consistency with the THG24 model.




Figure 5.2. Preliminary (explorative) paleomagnetic inclination flattening detection.

If the test shows promising results, the series of "*f*" increments can be adjusted to refine the results and maximize the reliability of the confidence boundaries. In the following example, after inspection of the results of Figure 5.2, *f* increments are reduced to 0.01 from *f*= 0.75 to *f*= 0.35, and left as 0.05 between 1 and 0.75, and between 0.35 and 0.3 (red-line rectangle in Figure 5.3). This is similar (not identical!) to the Quick=False option in the original Jupyter notebook of Tauxe et al. (2024).




Figure 5.3. Detailed inclination flattening calculation process.

The estimated best *f* value, indicated atop the right-down panel (pID) can be typed in within the "*flattening*" box of the "*Directions display & average*" panel (red-line rectangle in Figure 5.4) to work on the unflattened directions. This allows, for example, to calculate the VGPs by using the unflattened directions.




Figure 5.4. The f value is applied to the original dataset.

Please cite the original Tauxe et al. (2024) publication if using this test!

6. TK03.GAD E/I Test

This page performs the test for inclination flattening detection and correction of paleomagnetic directions based on the TK03.GAD PSV model of (Tauxe & Kent, 2004) (Figure 6.1). It is faster than the SVEI test, but it is based on an older PSV model and has less strict reliability criteria.

READ CAREFULLY: while this test can be used to quickly evaluate the expected degree of inclination shallowing of a distribution, it is warmly advised to use the more realistic SVEI test for publication purposes.

Export name: name of the downloadable files (buttons in main window).

Bootstrap?: for a very fast check the analysis can be performed without bootstrapping (no confidence boundaries determined).

Bootstraps number: number of simulated pseudosamples.

Perform: run test.

Export graph: downloads figure with test and histograms.

Export stats: download unflattened inclination and declination of E (average and 95% confidence boundaries).

For details on this adaptation of the original test please check Dallanave (Dallanave, 2024).




Figure 6.1. TK03.GAD-based inclination shallowing detection and correction.

7. VGP and Pmag Poles

The VGP and Pmag Poles page of Magnetic-A has a series of Action Buttons allowing a set of operations on VGPs, paleomagnetic poles, the average of them, and comparison with internal or loaded APWP. All Action Buttons open pop-up windows with different computations, producing numerical and graphical results displayed in the same stereographic projection (Figure 7.1).




Figure 7.1. VGP and Pmag Poles page. The red rectangles envelop the commands for the projection details, while the blue rectangle envelops the Action Buttons.

7.1. Projection commands

The commands enveloped by the red rectangles in Figure 7.1 are used to define the main features of the stereographic projection in the main panel.

Export name: name of the downloadable files (buttons in main window).

Type: the loaded VGPs sets can be displayed as VGP points, Fisher average, or bootstrapped pseudosamples mean.

Plot poles name: if "*yes*", it plots also the name of the VGPs sets onto the stereographic projection. Useful mostly if the sets are plotted as average pole with either A95 or B95.

Coastline: add or remove coastline from plot.

Center of plot: either "*Automatic*" or "*Manual*"; the first option centers the plot on the first selected entry. If "*manual*" is selected, latitude and longitude can be typed in the *Center lat* and *Center long* boxes, respectively.

Grid: defines angular distance between parallel and meridians.

Bigger and *smaller*: change size of plot.

7.2. Action buttons

7.2.1. Internal

Calculates, plots, and adds to *list of loaded VGPs and poles* (Figure 7.1) the VGPs from the directions plotted and manipulated in the *Directions display & average* page. Calculated VGPs can be exported in different formats.




Figure 7.2.1. Internal VGPs pop-up window.

Hemisphere: VGPs can be plotted either on the northern or the southern hemisphere.

Color and *symbol:* define the color and the symbol of the loaded VGPs.

VGPs to export: it defines the exportable VGPs file form. Three options are available: (1) "*Single mode"* = even with bimodal distribution, the VGPs are flipped on a common polarity; (2) "*Bimodal"* = if the distribution of paleomagnetic directions is bimodal, it keeps the VGPs in the original position; (3) "*Rotated*" = VGPs are bimodal but rotated to place the average paleomagnetic pole coinciding with the geographic north pole; this is used to determine (e.g.) the magnetic polarity stratigraphy.

Bootstrap n.: number of VGPs pseudosamples used for the B95 confidence angle (default is 2000).

Under this entry box, warning messages appear depending on unusual entry options.

Add to List of loaded VGPs: it loads the current VGPs onto the *List of loaded VGPs*, automatically calculating A95 and B95. Here the list of loaded VGPs is only visualized; the interactive list is shown in the next pages.

Export VGPs: downloads the VGPs as csv file in the format selected in the "*VGPs to export"* box.

7.2.2. External

Loads, plots, and manipulate existing VGPs sets, in the form of *Longitude* and *Latitude.* Accept csv files, with header, same as exported from previous window. Since the scope of the VGPs analysis window is to determine paleomagnetic poles, bimodal VGPs distributions are automatically converted into single mode.




Figure 7.2.2. External VGPs pop-up window.

VGPs name: name assigned to the distribution

Hemisphere: VGPs can be plotted either on the northern or the southern hemisphere.

Color and *symbol:* define the color and the symbol of the loaded VGPs.

Cut-off: Magnetic-A offers the possibility to apply a *cut-off*, either based on a given distance from the average pole (*VGP filter radius* window) or based on the method suggested by Vandamme (1994). Note that the *dynamic cut-off* is not possible as its calculation is performed by working on the paleomagnetic directions rather than on the VGPs.

Bootstrap n. and *Add to List of loaded VGPs*: as in 7.2.1

7.2.3. Simulate

This command is designed to generate a Fisher (1953) distribution of VGPs from known longitude, latitude, number of VGPs, and kappa precision parameter (Figure 7.2.3). The rationale behind using simulated VGPs rather than paleomagnetic pole when averaging multiple paleomagnetic poles is explained in (Vaes et al., 2023). The simulated VGPs can be uploaded in the *List of loaded VGPs* and processed as a real set of VGPs.




Figure 7.2.3. Fisherian VGPs simulator pop-up window.

Longitude, Latitude: longitude and latitude of the simulated VGPs center.

N, K: number of VGPs and Fisher (1953) precision parameter of the know paleomagnetic pole.

K tolerance: the precision parameter of the simulated VGPs might differ from the one typed in the box. Here, the maximum difference between the selected and the real simulated kappa can be typed in. The script reiterates the simulation until the difference is lower than the tolerance.

VGPs name, color, symbol Bootstrap n. and *Add to List of loaded VGPs*: as in 7.2.2

7.2.4. Merge

This sub-page (Figure 7.2.4) is designed to merge different VGPs file into one. This operation allows to calculate a weighted mean of different sets of VGPs, rather than simply averaging the paleomagnetic poles determined from the VGPs themselves. A similar (although with a more complex algorithm) is at the base of the GAPWP of Vaes et al. (Vaes et al., 2023).




Figure 7.2.4. VGPs Merging tool pop-up window.

M-VGPs name, color, symbol, Bootstrap n.: define name, color, and symbol of the merged VGPs, and number of bootstrapped pseudosamples to calculate the B95 parameter

Add merged VGPs to list: merges the VGPs selected from the main window into one.

Type: plots data as VGPs, fisher mean, or bootstrapped pseudosamples.

7.2.5. Pmag Pole

Add standard paleomagnetic poles either manually or from list file as exported from the main window (*Export poles table* button) or analogous format.




Figure 7.2.5. Add standard paleomagnetic pole pop-up window.

Pole long, lat, 95% confidence: Longitude, latitude, and 95% confidence angle of the external pole.

Pole color, symbol, name: color, symbol, and name of the pole as saved in the *List of loaded VGPs and Poles*:

Add to list of loaded VGPs and Poles: add external pole(s) to the main list.

7.2.6. Rotate

This command allows rotating any selected VGPs distribution, with the associated Fisher average and the bootstrapped pseudosamples, or paleomagnetic pole (Long, Lat, A95) by applying the rotational parameters typed in by the user (Figure 7.2.6).




Figure 7.2.6. Rotate VGPs and Poles pop up window.

Color, symbol: color and symbol all the rotated dataset (they can be updated later after double click on the main list).

Type: plots data as VGPs, fisher mean, or bootstrapped pseudosamples.

E_Pole Long, E_Pole_Lat, Rotation: rotational parameters applied to the datasets, which are then rotated by clicking on the *Rotate* button and added to the main list by clicking on the *Add to list of loaded VGPs and Poles* button.

7.2.7. Fisher A95

Adds Fisher average and 95% confidence angle (A95) of all selected poles and VGP averages to the *list of loaded VGPs and poles* (Figure 7.2.10).




Figure 7.2.10. Pop-up window allowing the Fisher mean and associated A95 calculation of the entries selected in the list of loaded VGPs and Poles.

7.2.8. APWP

Add an apparent polar wander path (APWP) to the plot. Magnetic-A has two global APWP internally loaded (Torsvik et al., 2012; Vaes et al., 2023; respectively called T2012 and V2023) (Figure 7.2.7) in all the different reference frames compiled by the authors in the original publication. A custom APWP can be compiled and uploaded (see Appendix 3).



Figure 7.2.7. Apparent polar wander pop-up window

7.2.9. Locality

Add any geographic point to the stereographic projection. The list is appended within the pop-up window, so the entries must be selected here for plotting.




Figure 7.2.8. Pop-up window showing the plotting details and the list of the loaded localities.

7.2.10. Small circle

This window (Figure 7.2.9) allows to plot circles with defined radius around any geographic point. The list is appended within the pop-up window, so the entries must be selected here for plotting.




Figure 7.2.9. Pop-up window showing the plotting details and the list of the loaded circles.

7.2.11. Great circle

Interpolate all selected VGPs, Poles, and (if specified) Localities with a plane, plotted on the stereographic projection as a great circle and expressed numerically as a pole (Figure 7.2.11). The pole direction is estimated by the orientation of the minimum eigenvector of the 3x3 distribution matrix interpolating the points. The list of the poles is appended within the pop-up window, and it must be selected here for plotting.




Figure 7.2.11. Pop-up window with the details of the great circle interpolating the selected geographic points (VGPs average, poles, localities).

Symbol: symbol of the pole of the plane (it can be omitted from the graph).

Color: color of both great circle and symbol associated to the interpolating plane.

Plot pole: includes or omit the pole from the figure.

Add locality to Plane: If yes, it includes any selected locality within the interpolation.

7.2.12. Other commands within the VGPs and Pmag Poles window

Delete: delete selected entries from List of Loaded VGPs and Poles

Export graph: export image as pdf vector file.

Export Pole Table: export the full *List of loaded VGPs and Poles* (for uploading and plotting as Pmag Poles file).

Export selected VGPs sets: export the sets of VGPs selected in the *List of loaded VGPs and Poles.*

Export selected stats: export the statistical parameters of the sets selected in the *List of loaded VGPs and Poles.*

8. Magnetic polarity

This page is designed to compile a classic magnetic polarity stratigraphy plot. To generate this plot, this page uses the paleomagnetic declination and inclination data uploaded in the *Directions display & average* page, so please refer to Chapter 2 of this guide to verify the data format (see also Appendix 3 below). In case of paleomagnetic directions isolated with Magnetic-A, the position of the specimens is missing from the directions file. In this case, a file with the sample position can be uploaded here (format is described in the pop-up window that opens clicking on the question mark).




Figure 8.1. Magnetic polarity stratigraphy page

Upload depth file: samples depth file. To date, two formats are accepted (see also online help):

- A simple two columns csv file with sample code and depth. Be aware that the depth is assigned by checking the sample name, so THE SAMPLE NAME MUST BE THE SAME in both this file and the demagnetization data file.
- Spinner or ex-spinner file downloaded from LIMS (https://web.iodp.tamu.edu/LORE/), in case of discrete specimen data generated on the JOIDES Resolution. This file is must be the same uploaded in the Vector end-points interpolation window.

The file type does not need to be selected, the script recognizes the file depending on the number of columns.

Export name: name of the downloaded graph or reversals file.

Base and *Top plot*: a specific stratigraphic interval can be selected for plotting by typing in the stratigraphic base and top. By leaving the cells empty, the script will use the entire stratigraphic interval of the data.

Reverse depth: the position of the samples will be trated as top-to-down depth (e.g., samples from deep-sea drilled cores).

Unit: depth unit that will appear in the graph.

Vertical grid: defines the spacing of the horizontal dotted line in the graph.

Point color: defines the color of the data points.

Decl. offset (°): defines the horizontal offset in the declination plot. In case of paleomagentic directions organize in two modes pointing quasi-north-and-south like the ones of the example (Figure 8.1), the declination plot would result in annoying left-and-right swings between 0° and 360° degree. Setting an offset of 90° avoid this graphical problem.

VGP/Incl.: the magnetic polarity stratigraphy is determined by using three possible strategy:

- *Use VGP*: when paleomagnetic directions are defined by both declination and inclination.
- *Use Inc (N.Hem)*: uses only inclination data, assuming down-pointing inclination as normal.
- *Use Inc (S.Hem)*: uses only inclination data, assuming up-pointing inclination as normal.

Export graph: download the vector graphic file in your device.

Export rev.: download a csv file with top and bottom of the normal polarity zones, defined by the arithmetic mean of the position of two consecutive samples with different polarity.

9. Site map

Magnetic-A includes also a very simple tool for plotting localities in a present-day geographic sketch, which might be useful for publication purposes. While it automatically plots the working site with the coordinates typed in the *Directions display, filter & average* page, a file with several localities name, coordinates, and symbol features can be uploaded (Appendix 3).




Figure 9.1. Site map page.

Internal site name: name of the site/localities which coordinates are typed in the *Directions display, filter & average* page (Chapter 2 of this manual).

Site file: a list of sites can be uploaded here, csv format, with five columns that include site name, longitude, latitude, symbol, color (see also online helper and Appendix 4).

Color and *symbol*: color and symbol of the internal site.

Map grid: angular distance between parallels and meridians.

Central meridian: the map can be centered either on Greenwich or 180°E meridian (useful for Pacific and peri-Pacific sites).

Land, Sea, Grid color: define the color of respectively land, sea and grid lines.

Plot – Refresh: plot or update map if anything has been changed.

Delete sites file: ignore the uploaded site file when plotting.

Glossary

A^2D and A^2I = Anderson-Darling test on declination and inclination (respectively) as a function of the flattening factor "*f*". In both cases, a value higher than 3.07 rejects the hypothesis that the considered directions distribution is drawn from the THG24 GGP model (please refer to Tauxe et al., 2024 for computational details).

A95= 95% confidence angle around the paleomagnetic pole estimated by using Fisher (1953) statistic.

APWP= Apparent polar wander path.

B95= 95% confidence angle around the paleomagnetic pole estimated by using bootstrap resampling. The routine that performs this calculation is described in Dallanave (2024).

ChRM= Characteristic remanent magnetization.

csv= comma separated values.

E= Elongation

GAPWP= global synthetic apparent polar wander path.

GGP= Giant Gaussian Process.

I= Inclination

MAD= maximum angular deviation.

NRM= natural remanent magnetization.

pID= combined *p*-values calculated from A^2D and A^2I. A value lower than 0.025 indicates that the distribution of directions is not drawn from the THG24 GGP model (please refer to Tauxe et al., 2024 for computational details).

PCA= principal component analysis.

PSV= paleosecular variations.

T2012= GAPWP of Torsvik et al., (2012).

THG24= Tauxe-Heslop-Gilder (2024) field model.

TK03.GAD= Tauxe and Kent (2004) field model.

V2023= GAPWP of (Vaes et al., 2023).

VEPs= Vector end-points.

VGPs= virtual geomagnetic poles.

References

Anderson, T. W., & Darling, D. A. (1952). Asymptotic theory of certain "goodness-of-fit" criteria based on stochastic processes. *The Annals of Mathematical Statistics*, *23*(2), 193–212. https://doi.org/10.1214/aoms/1177729437

Arason, P., & Levi, S. (2010). Maximum likelihood solution for inclination-only data in paleomagnetism. *Geophysical Journal International*, *182*(2), 753–771. https://doi.org/10.1111/j.1365-246X.2010.04671.x

Butler, R. F. (1992). *Paleomagnetism: Magnetic Domains to Geologic Terranes*. Blackwell Scientific Publication. https://www.geo.arizona.edu/Paleomag/

Dallanave, E. (2024). Assessing the reliability of paleomagnetic datasets using the R package PmagDiR. *Scientific Reports*, *14*(1666). https://doi.org/10.1038/s41598-024-52001-x

Dallanave, E., Agnini, C., Muttoni, G., & Rio, D. (2012). Paleocene magneto-biostratigraphy and climate-controlled rock magnetism from the Belluno Basin, Tethys Ocean, Italy. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *337–338*, 130–142. https://doi.org/10.1016/j.palaeo.2012.04.007

Deenen, M. H. L., Langereis, C. G., van Hinsbergen, D. J. J., & Biggin, A. J. (2011). Geomagnetic secular variation and the statistics of palaeomagnetic directions. *Geophysical Journal International*, *186*(2), 509–520. https://doi.org/10.1111/j.1365-246X.2011.05050.x

Fisher, R. (1953). Dispersion on a sphere. *Proceedings of the Royal Society of London*, *A217*, 295–305.

Gilder, S. A., Chen, Y., & Sen, S. (2001). Oligo-Miocene magnetostratigraphy and rock magnetism of the Xishuigou section, Subei (Gansu Province, western China) and implications for shallow inclinations in central Asia. *Journal of Geophysical Research*, *106*(12), 30505–30521.

Heslop, D., & Roberts, A. P. (2016). Analyzing paleomagnetic data: To anchor or not to anchor? *Journal of Geophysical Research: Solid Earth*, *121*(11), 7742–7753. https://doi.org/10.1002/2016JB013387

Heslop, D., Scealy, J. L., Wood, A. T. A., Tauxe, L., & Roberts, A. P. (2023). A Bootstrap Common Mean Direction Test. *Journal of Geophysical Research: Solid Earth*, *128*(8), 1–12. https://doi.org/10.1029/2023JB026983

King, R. F. (1955). The remanent magnetism of artificially deposited sediments. *Monograph of the Natural Royal Astronomica Society, Geophysical Supplement*, *7*, 115–134.

Kirschvink, J. L. (1980). The least-squares line and plane and the analysis of palaeomagnetic data. *Geophysical Journal of the Royal Astronomical Society*, *62*(3), 699–718. https://doi.org/10.1111/j.1365-246X.1980.tb02601.x

McFadden, P. L., & McElhinny, M. W. (1988). The combined analysis of remagnetization circles and direct observations in palaeomagnetism. *Earth and Planetary Science Letters*, *87*(1–2), 161–172. https://doi.org/10.1016/0012-821X(88)90072-6

Tauxe, L. (2010). *Paleomagnetic principle and practice*. Kluwer academic publishers.

Tauxe, L., Heslop, D., & Gilder, S. A. (2024). Assessing Paleosecular Variation Averaging and Correcting Paleomagnetic Inclination Shallowing. *Journal of Geophysical Research: Solid Earth*, *129*(8). https://doi.org/10.1029/2024JB029502

Tauxe, L., & Kent, D. V. (2004). A simplified statistical model for the geomagnetic field and the detection of shallow bias in paleomagnetic inclinations: Was the ancient magnetic field dipolar? In J. E. T. Channell, D. V. Kent, W. Lowrie, & J. G. Meert (Eds.), *Timescales of the Paleomagnetic Field, Geophys. Monogr.* (Vol. 145, pp. 101–115). American Geophysical Union. http://academiccommons.columbia.edu/catalog/ac:144298

Torsvik, T. H., Van der Voo, R., Preeden, U., Mac Niocaill, C., Steinberger, B., Doubrovine, P. V., van Hinsbergen, D. J. J., Domeier, M., Gaina, C., Tohver, E., Meert, J. G., McCausland, P. J. a., & Cocks, L. R. M. (2012). Phanerozoic polar wander,

palaeogeography and dynamics. *Earth-Science Reviews*, *114*(3–4), 325–368. https://doi.org/10.1016/j.earscirev.2012.06.007
Vaes, B., van Hinsbergen, D. J. J., van de Lagemaat, S. H. A., van der Wiel, E., Lom, N., Advokaat, E., Boschman, L. M., Gallo, L. C., Greve, A., & Guilmette, C. (2023). A global apparent polar wander path for the last 320 Ma calculated from site-level paleomagnetic data. *Earth-Science Reviews*, *245*(104547), 1–35. https://doi.org/10.1016/j.earscirev.2023.104547
Vandamme, D. (1994). A new method to determine paleosecular variation. *Physics of the Earth and Planetary Interiors*, *85*(1–2), 131–142. https://doi.org/10.1016/0031-9201(94)90012-4
Watson, G. S. (1956). A test for randomness of directions. *Geophysical Journal International*, *7*(s4), 160–161. https://doi.org/10.1111/j.1365-246X.1956.tb05561.x

Appendix 1: Demagnetization data file accepted formats

NOTE: JR6A expanded data can be retrieved under **Magnetism** in the following link: *https://web.iodp.tamu.edu/LORE/*

Magnetic-A.csv

```
sample,step,Sx,Sy,Sz,Gx,Gy,Gz,Bx,By,Bz
ar001,0,-0.00361,-0.01776,0.08674,0.06726,-0.00541,0.05743,0.07922,-0.00749,0.03898
ar001,80,-0.01496,-0.01582,0.04329,0.0044,-0.02745,0.0397,0.01375,-0.02909,0.03624
ar001,120,0.00811,-0.03025,-0.02536,-0.0314,-0.01094,-0.02277,-0.03605,-0.01013,-0.0149
ar001,200,0.01698,-0.03436,-0.04519,-0.04069,-0.0071,-0.04248,-0.04983,-0.0055,-0.03157
ar001,250,0.01511,-0.03578,-0.05307,-0.04665,-0.00285,-0.04626,-0.05653,-0.00118,-0.03357
ar001,300,0.01864,-0.03566,-0.05301,-0.04499,-0.00457,-0.04883,-0.05556,-0.00272,-0.03654
ar001,320,0.0249,-0.0313,-0.04975,-0.03725,-0.00637,-0.05144,-0.0487,-0.00435,-0.04103
ar001,350,0.01324,-0.03522,-0.04885,-0.04492,-0.00393,-0.04205,-0.05383,-0.00235,-0.02999
ar001,375,0.01688,-0.02799,-0.05272,-0.04009,0.00147,-0.04731,-0.05039,0.00326,-0.03602
ar001,400,0.01744,-0.02792,-0.05127,-0.039,0.00048,-0.04681,-0.04918,0.00223,-0.0359
ar001,425,0.01644,-0.02824,-0.03867,-0.03311,-0.0062,-0.0378,-0.04136,-0.00471,-0.02882
ar001,450,0.01519,-0.02489,-0.03601,-0.02985,-0.00473,-0.03513,-0.03753,-0.00335,-0.02698
ar001,475,0.01221,-0.02571,-0.04095,-0.03443,-0.00126,-0.03605,-0.04215,7e-05,-0.02665
ar001,500,0.01064,-0.02079,-0.03022,-0.02598,-0.00291,-0.02784,-0.03198,-0.00184,-0.0208
ar001,525,0.00836,-0.01786,-0.03602,-0.02791,0.00333,-0.02993,-0.03434,0.00446,-0.02206
ar001,550,0.00876,-0.01773,-0.02357,-0.02116,-0.00354,-0.02206,-0.0259,-0.00268,-0.0164
ar001,575,0.00535,-0.01784,-0.01889,-0.02034,-0.00458,-0.01641,-0.02373,-0.00397,-0.01117
ar001,600,0.00115,-0.01113,-0.01818,-0.01701,0.0017,-0.01279,-0.01959,0.00216,-0.0082
ar001,625,0.00491,-0.00549,-0.00893,-0.0064,-0.00124,-0.00956,-0.00854,-0.00087,-0.00776
ar001,650,0.00172,-0.0041,0.00294,-0.00065,-0.00525,0.00063,-0.00051,-0.00528,0.00055
```

Figure A1.1. Magnetic-A data file format for demagnetization data.

ardo1_LASA.txt

```
******************************** 22 July      2009 AzFile = ********************
LAT:  0.00  LON:   0.00 REC:2131 SAM: 115
!
!   ID     TREAT  IC CD    J    CDECL CINCL  GDECL GINCL  BDECL BINCL  SUSC  V/M
!J units 1e-4 emu (equivalent to 1e-2 A/m assuming 10 cc volume)
!_____________________________________________________________________________
ar001        0.0  Gh  0 0.08861 258.5  78.2  355.4  40.4  354.6  26.1 -99.0  0.0
ar001       80.0  Gh  0 0.04846 226.6  63.3  279.1  55.0  295.3  48.4 -99.0  0.0
ar001      120.0  Gh  0 0.04030 285.0 -39.0  199.2 -34.4  195.7 -21.7 -99.0  0.0
ar001      200.0  Gh  0 0.05925 296.3 -49.7  189.9 -45.8  186.3 -32.2 -99.0  0.0
ar001      250.0  Gh  0 0.06576 292.9 -53.8  183.5 -44.7  181.2 -30.7 -99.0  0.0
ar001      300.0  Gh  0 0.06655 297.6 -52.8  185.8 -47.2  182.8 -33.3 -99.0  0.0
ar001      320.0  Gh  0 0.06383 308.5 -51.2  189.7 -53.7  185.1 -40.0 -99.0  0.0
ar001      350.0  Gh  0 0.06166 290.6 -52.4  185.0 -43.0  182.5 -29.1 -99.0  0.0
ar001      375.0  Gh  0 0.06203 301.1 -58.2  177.9 -49.7  176.3 -35.5 -99.0  0.0
ar001      400.0  Gh  0 0.06093 302.0 -57.3  179.3 -50.2  177.4 -36.1 -99.0  0.0
ar001      425.0  Gh  0 0.05063 300.2 -49.8  190.6 -48.3  186.5 -34.7 -99.0  0.0
ar001      450.0  Gh  0 0.04634 301.4 -51.0  189.0 -49.3  185.1 -35.6 -99.0  0.0
ar001      475.0  Gh  0 0.04987 295.4 -55.2  182.1 -46.3  179.9 -32.3 -99.0  0.0
ar001      500.0  Gh  1 0.03819 297.1 -52.3  186.4 -46.8  183.3 -33.0 -99.0  0.0
ar001      525.0  Gh  0 0.04106 295.1 -61.3  173.2 -46.8  172.6 -32.5 -99.0  0.0
ar001      550.0  Gh  1 0.03077 296.3 -50.0  189.5 -45.8  185.9 -32.2 -99.0  0.0
ar001      575.0  Gh  1 0.02653 286.7 -45.4  192.7 -38.2  189.5 -24.9 -99.0  0.0
ar001      600.0  Gh  1 0.02135 275.9 -58.4  174.3 -36.8  173.7 -22.6 -99.0  0.0
ar001      625.0  Gh  1 0.01157 311.8 -50.5  191.0 -55.7  185.8 -42.1 -99.0  0.0
ar001      650.0  Gh  5 0.00533 292.8  33.5  262.9   6.8  264.5   5.9 -99.0  0.0
```

Figure A1.2. LASA data file format for demagnetization data.

COR_example_file.COR

Name	Field (mT)	Squid_x	Squid_y	Squid_z (emu)
U1507B001	0.0	-1.19518E-4	-5.55012E-5	-1.09887E-4
U1507B001	5.0	-1.13729E-4	-6.24101E-5	-1.07915E-4
U1507B001	10.0	-1.02764E-4	-6.26037E-5	-9.84553E-5
U1507B001	15.0	-9.09908E-5	-5.77228E-5	-8.78316E-5
U1507B001	20.0	-7.97847E-5	-5.17261E-5	-7.77476E-5
U1507B001	25.0	-6.98384E-5	-4.60461E-5	-6.90506E-5
U1507B001	30.0	-6.09014E-5	-4.06279E-5	-6.10760E-5
U1507B001	35.0	-5.41789E-5	-3.64687E-5	-5.48418E-5
U1507B001	40.0	-4.95503E-5	-3.33774E-5	-5.00430E-5
U1507B001	45.0	-4.44535E-5	-3.00566E-5	-4.54666E-5
U1507B001	50.0	-3.98352E-5	-2.81075E-5	-4.14571E-5
U1507B001	60.0	-3.35216E-5	-2.36917E-5	-3.49746E-5
U1507B001	70.0	-2.78695E-5	-1.93478E-5	-2.92240E-5
U1507B001	80.0	-2.38148E-5	-1.67695E-5	-2.58164E-5
U1507B001	90.0	-1.94629E-5	-1.39544E-5	-2.14934E-5
U1507B001	100.0	-1.57446E-5	-1.15818E-5	-1.77652E-5

Figure A1.3. Bremen (.cor) data file format for demagnetization data.

koa001a

```
    koa001a  koa001a
     0.0  78.0  49.0  68.1  35.1  11.0
TH  20  24.4 -31.5 357.2 -51.4 5.10E-07   1.1 316.6   2.3  0.0000  0.0000  0.0000
TH 100  34.6 -23.0  16.5 -49.6 6.82E-08   1.3 319.2  14.3  0.0000  0.0000  0.0000
TH 150  22.6 -30.3 356.5 -49.6 5.55E-08   3.5 314.7   2.2  0.0000  0.0000  0.0000
TH 250   9.3 -51.9 318.2 -55.8 5.74E-08   2.1 322.2 -20.3  0.0000  0.0000  0.0000
TH 300 355.9 -58.8 301.8 -52.2 6.63E-08   1.3 323.4 -30.5  0.0000  0.0000  0.0000
TH 325 352.7 -61.1 297.3 -51.9 6.40E-08   1.1 325.0 -33.0  0.0000  0.0000  0.0000
TH 350 350.6 -60.6 297.2 -50.7 6.28E-08   1.6 323.9 -33.7  0.0000  0.0000  0.0000
TH 375 351.5 -64.0 292.7 -52.5 5.74E-08   1.9 327.8 -34.7  0.0000  0.0000  0.0000
TH 400 357.1 -62.2 297.0 -54.2 5.20E-08   1.1 327.3 -31.7  0.0000  0.0000  0.0000
TH 425 359.3 -64.6 293.7 -55.9 4.51E-08   1.0 330.3 -32.0  0.0000  0.0000  0.0000
TH 450  11.1 -61.4 302.2 -60.3 3.52E-08   1.1 330.7 -25.7  0.0000  0.0000  0.0000
TH 475   1.3 -61.3 299.7 -55.7 3.00E-08   1.7 327.6 -29.5  0.0000  0.0000  0.0000
TH 500  14.8 -73.0 278.0 -62.0 1.60E-08   1.8 342.1 -31.9  0.0000  0.0000  0.0000
TH 525 220.7 -75.3 239.1 -41.4 5.01E-09   6.1  14.4 -51.9  0.0000  0.0000  0.0000
TH 550 193.6   2.8 186.9  21.9 7.95E-09   3.0 109.5 -17.1  0.0000  0.0000  0.0000
TH 575 264.6  49.6 317.3  78.7 2.21E-09  12.1 184.3   0.8  0.0000  0.0000  0.0000
TH 600   6.9 -75.0 274.7 -59.5 5.19E-09   2.7 342.6 -34.8  0.0000  0.0000  0.0000
```

Figure A1.4. CIT data file format for demagnetization data.

This is the *Sample Data File* format designed for *PaleoMag X*. For details, please follow this link:

https://cires1.colorado.edu/people/jones.craig/PMag_Formats.html

Appendix 2: Average directions and 95% confidence files format.

FISHER_2026-03-06_stat.csv

```
"","dec","inc","a95","N","k"
"Mode 1",353.47,35.03,3.91,82,16.97
"Mode 2",177.01,-36.13,2.44,157,22.26
"All",355.8,35.77,2.08,239,20.08
```

Figure A2.1. Average directions and Fisher (1953) confidence as exported from Magnetic-A (sub-page 3.1 above)



Figure A2.2. Average directions and elliptic confidence (Deenen et al., 2011) as exported from Magnetic-A (sub-page 3.1 above)




Figure A2.3. Example of average directions and confidence accepted by Magnetic-A: Fisher 95% confidence.




Figure A2.4. Example of average directions and confidence accepted by Magnetic-A: Fisher 95% confidence plus symbol and colour details.



Figure A2.5. Example of average directions and confidence accepted by Magnetic-A: elliptic 95% confidence.



Figure A2.6. Example of average directions and confidence accepted by Magnetic-A: elliptic 95% confidence plus symbol and colour details.

Appendix 3: Custom APWP file format





Figure A2.1. Example of external APWP file.


Appendix 3: Direction file formats for plotting magnetostratigraphy

Three types are shown: Dec, Inc, Pos (Figure A3.1); G_dec, G_inc, Bedding_azimuth, Bedding plunge, Position (Figure A3.2); G_dec, G_inc, B_dec, B_inc, Position (Figure A3.3). In all files: G_dec and G_inc = Declination and inclination in geographic coordinates; B_dec, B_inc= Declination and inclination in tilt corrected coordinates.


Ardo_PmagDiR.csv


```
"B_Dec","B_Inc","Position"
181.5,-34.1,-1
172.1,-46.5,-0.58
186.3,-43.3,-0.18
183.3,-38.8,0.18
184.4,-19.3,0.34
175.5,-13,0.4
170.9,-38.3,0.45
350.1,29.6,0.51
358,39.8,0.56
2.5,36.7,0.57
358.3,35,0.68
344.9,31.9,0.99
343.8,34.3,1
344.3,42.1,1.12
338.5,25.5,1.35
340.3,42.5,1.85
346,27.5,2.12
349.3,25.5,2.14
342.7,30.4,2.16
336.3,32.4,2.44
345.2,44.5,2.87
299,32.8,3.45
338.6,49.2,3.65
306.8,44.5,3.71
300.5,44.7,3.71
145.9,-36.4,3.8
176.6,-38.4,3.9
178.9,-27.7,3.95
175.3,-25.7,4.15
4.4,43.4,4.25
332.5,31.8,4.46
```

Figure A3.1


Ardo_Geo_bed_PmagDiR.csv

```
"G_dec","G_inc","B_az","B_plung","Position"
184.3,-48.1,350.1,14.3,-1
173,-60.8,350.1,14.3,-0.58
191.9,-56.8,350.1,14.3,-0.18
187.1,-52.7,350.1,14.3,0.18
186.3,-33.1,350.1,14.3,0.34
176,-27.2,350.1,14.3,0.4
171.3,-52.5,350.1,14.3,0.45
350.2,43.9,350.1,14.3,0.51
0.6,53.9,350.1,14.3,0.56
5.9,50.5,350.1,14.3,0.57
0.4,49.1,350.1,14.3,0.68
343.9,46.1,350.1,14.3,0.99
342.4,48.5,350.1,14.3,1
342.3,56.3,350.1,14.3,1.12
336.6,39.4,350.1,14.3,1.35
337,56.5,350.1,14.3,1.85
345.3,41.7,350.1,14.3,2.12
349.2,39.8,350.1,14.3,2.14
341.2,44.5,350.1,14.3,2.16
333.3,46.2,350.1,14.3,2.44
343.3,58.7,350.1,14.3,2.87
290.1,41,350.1,14.3,3.45
333.7,63.1,350.1,14.3,3.65
294.2,53.9,350.1,14.3,3.71
286.9,52.8,350.1,14.3,3.71
139.8,-49.1,350.1,14.3,3.8
178.7,-52.6,350.1,14.3,3.9
180.7,-41.8,350.1,14.3,3.95
176.2,-39.9,350.1,14.3,4.15
9.7,57.1,350.1,14.3,4.25
328.6,45.3,350.1,14.3,4.46
```


Figure A3.2


Ardo_Geo_TC_PmagDiR... — Edited

```
"G_dec","G_inc","B_dec","B_inc","Position"
184.3,-48.1,181.5,-34.1,-1
173,-60.8,172.1,-46.5,-0.58
191.9,-56.8,186.3,-43.3,-0.18
187.1,-52.7,183.3,-38.8,0.18
186.3,-33.1,184.4,-19.3,0.34
176,-27.2,175.5,-13,0.4
171.3,-52.5,170.9,-38.3,0.45
350.2,43.9,350.1,29.6,0.51
0.6,53.9,358,39.8,0.56
5.9,50.5,2.5,36.7,0.57
0.4,49.1,358.3,35,0.68
343.9,46.1,344.9,31.9,0.99
342.4,48.5,343.8,34.3,1
342.3,56.3,344.3,42.1,1.12
336.6,39.4,338.5,25.5,1.35
337,56.5,340.3,42.5,1.85
345.3,41.7,346,27.5,2.12
349.2,39.8,349.3,25.5,2.14
341.2,44.5,342.7,30.4,2.16
333.3,46.2,336.3,32.4,2.44
343.3,58.7,345.2,44.5,2.87
290.1,41,299,32.8,3.45
333.7,63.1,338.6,49.2,3.65
294.2,53.9,306.8,44.5,3.71
286.9,52.8,300.5,44.7,3.71
139.8,-49.1,145.9,-36.4,3.8
178.7,-52.6,176.6,-38.4,3.9
180.7,-41.8,178.9,-27.7,3.95
176.2,-39.9,175.3,-25.7,4.15
9.7,57.1,4.4,43.4,4.25
328.6,45.3,332.5,31.8,4.46
```

Figure A3.3

Appendix 4: Sites file for mapping tool


localities_file.csv

```
Name,Long,Lat,Symbol,Color
New York,-74,40.7,d,yellow
Dubai,55.2707828,25.2048493,s,light green
Perth,115.860457,-31.950527,t,cyan
Cape Town,18.424055,-33.924869,c,pink
```

Figure A4.1. Example of Sites file to upload on the Site map page.