

TITLE OF THE PAPER

EDUARDO SILVA
ESILVA@DIM.UCHILE.CL

ABSTRACT. abstract

Keywords: Keyword 1, Keyword 2, Keyword 3.

INTRODUCTION

Introduction.

We show:

Theorem ??. *Every countable group G has a non-empty, strongly aperiodic subshift on the alphabet $\{0, 1\}$.*

Theorem ??. *Every finitely generated group G has a non-empty G -effectively closed strongly aperiodic subshift.*

1. PRELIMINARIES

Throughout this article the groups G considered will always be countable; we denote their identity element 1_G . When G is finitely generated we associate a finite set $S \subset G$ of generators and the undirected right Cayley graph $\Gamma(G, S) = (G, \{\{g, gs\} \mid g \in G, s \in S\})$ so that (G, d) is a metric space where d is the distance induced on G by $\Gamma(G, S)$. If two words w_1, w_2 in S^* represent the same element in G , we write $w_1 =_G w_2$. We shall denote by $B(g, n) = \{h \in G \mid d(g, h) \leq n\}$ the ball of size n centered in $g \in G$. In general we denote $B_\Gamma(v, n)$ the ball of size n centered in v of an arbitrary graph Γ . For $g \in G$, we denote by $|g|$ the length of a shortest path from 1_G to g in $\Gamma(G, S)$, that is to say $|g| = d(1_G, g)$. We also denote by $\text{WP}(G) := \{w \in (S \cup S^{-1})^* \mid w =_G 1_G\}$ the set of words which can be written using elements from S and their inverses which are equal to 1_G in the group G . If $\text{WP}(G)$ is a decidable language we say that G has decidable word problem. For more references see [17].

We now give some basic definitions of symbolic dynamics. For a more complete introduction the reader may refer to [15, 8]. Let \mathcal{A} be a finite alphabet and G a countable group. The set $\mathcal{A}^G = \{x : G \rightarrow \mathcal{A}\}$ equipped with the left group action $\sigma : G \times \mathcal{A}^G \rightarrow \mathcal{A}^G$ defined by $(\sigma_g(x))_h = x_{g^{-1}h}$ is the G -fullshift. The elements $a \in \mathcal{A}$ and $x \in \mathcal{A}^G$ are called *symbols* and *configurations* respectively. By taking the discrete topology on \mathcal{A} we obtain that the set of configurations \mathcal{A}^G is compact and metrizable. In the case of a countable group, given an enumeration $1_G = g_0, g_1, \dots$ of G , the topology is generated by the metric $d(x, y) = 2^{-\inf\{n \in \mathbb{N} \mid x_{g_n} \neq y_{g_n}\}}$. If E is a subset of \mathcal{A}^G , we denote by \overline{E} its topological closure. In the case of a finitely generated group another possibility which is more practical is $d(x, y) = 2^{-\inf\{|g| \mid g \in G : x_g \neq y_g\}}$. This topology is generated by a clopen basis given by the *cylinders* $[a]_g = \{x \in \mathcal{A}^G \mid x_g = a \in \mathcal{A}\}$. A *support* is

a finite subset $F \subset G$. Given a support F , a *pattern with support F* is an element p of \mathcal{A}^F , i.e. a finite configuration and we write $\text{supp}(p) = F$. We also denote the cylinder generated by p centered in g as $[p]_g = \bigcap_{h \in F} [p_h]_{gh}$. One says that a pattern $p \in \mathcal{A}^F$ *appears* in a configuration $x \in \mathcal{A}^G$ if there exists $g \in G$ such that for any $h \in F$, $x_{gh} = p_h$, said otherwise, if there exists g such that $x \in [p]_g$. In this case we write $p \sqsubset x$. We denote the set of finite patterns over G as $\mathcal{A}_G^* := \bigcup_{F \subset G, |F| < \infty} \mathcal{A}^F$.

Definition 1.1. A subset X of \mathcal{A}^G is a G -subshift if it is σ -invariant – $\sigma(X) \subset X$ – and closed for the cylinder topology. Equivalently, X is a G -subshift if and only if there exists a set of forbidden patterns $\mathcal{F} \subset \mathcal{A}_G^*$ that defines it.

$$X = X_{\mathcal{F}} := \{ x \in \mathcal{A}^G \mid \forall p \in \mathcal{F}, p \not\sqsubset x \} = \bigcap_{p \in \mathcal{F}, g \in G} \mathcal{A}^G \setminus [p]_g.$$

That is, a G -subshift is a shift-invariant subset of \mathcal{A}^G which can be written as the complement of a union of cylinders. If it is clear from the context, we will drop the G and simply refer to a subshift. A subshift $X \subseteq \mathcal{A}^G$ is of *finite type* – G -SFT for short – if there exists a finite set of forbidden patterns \mathcal{F} such that $X = X_{\mathcal{F}}$.

Consider a group which is generated by a finite set S . A *pattern coding* c is a finite set of tuples $c = (w_i, a_i)_{1 \leq i \leq n}$ where $w_i \in (S \cup S^{-1})^*$ and $a_i \in \mathcal{A}$. We say that a pattern coding is *consistent* if for every pair of tuples such that $w_i =_G w_j$ (w_i and w_j represent the same element under G) then $a_i = a_j$. We say a consistent pattern coding c *codifies* a pattern P if every w_i represents an element of $\text{supp}(P)$ and for every $g \in \text{supp}(P)$ there exists a tuple $(w_i, a_i) \in c$ such that $g =_G w_i$ and $P_g = a_i$.

Definition 1.2. For a finitely generated group G we say a subshift $X \subseteq \mathcal{A}^G$ is G -effectively closed if there exists a Turing machine with oracle $\text{WP}(G)$ which recognizes a set of pattern codings such that the consistent ones codify a set of patterns \mathcal{F} such that $X = X_{\mathcal{F}}$. If the same property is valid without the oracle we say X is effectively closed.

Being G -effectively closed is a generalization of the same concept for \mathbb{Z} -subshifts where the set of forbidden patterns is a recursively enumerable set of words.

Let $x \in \mathcal{A}^G$ be a configuration. The *orbit* of x is the set of configurations $\text{orb}_{\sigma}(x) = \{\sigma_g(x) \mid g \in G\}$, and the *stabilizer* of x is the set of group elements $\text{stab}_{\sigma}(x) = \{g \in G \mid \sigma_g(x) = x\}$.

Definition 1.3. A G -subshift $X \subseteq \mathcal{A}^G$ is weakly aperiodic if for every configuration $x \in X$, $|\text{orb}_{\sigma}(x)| = \infty$. A G -subshift $X \subseteq \mathcal{A}^G$ is strongly aperiodic if for every configuration $x \in X$, $\text{stab}_{\sigma}(x) = \{1_G\}$.

For infinite groups the weak concept of aperiodicity is relevant and implied by strong aperiodicity.

2. NON-EMPTY STRONGLY APERIODIC SUBSHIFTS

2.1. A non-empty strongly aperiodic subshift over $\{0, 1\}$ in any countable group.

2.2. A graph-oriented construction and some computational properties. we still don't know if this kind of construction is always possible. To our knowledge the following question remains open:

Question. *dssad?*

Acknowledgements: This work was partially supported by .

REFERENCES

- [1] Nathalie Aubrun, Sebastián Barbieri, and Mathieu Sablik. A notion of effectiveness for subshifts on finitely generated groups. *Theoretical Computer Science*, 661:35–55, 2017.
- [2] Noga Alon, Jarosław Grytczuk, Mariusz Haluszczak, and Oliver Riordan. Nonrepetitive colorings of graphs. *Random Structures & Algorithms*, 21(3-4):336–346, 2002.
- [3] Noga Alon and Joel H. Spencer. *The probabilistic method*. Wiley, 2008.
- [4] Alexis Ballier and Maya Stein. The domino problem on groups of polynomial growth. *arXiv preprint arXiv:1311.4222*, 2013.
- [5] Valérie Berthé and Laurent Vuillon. Tilings and rotations on the torus: a two-dimensional generalization of sturmian sequences. *Discrete Mathematics*, 223(1-3):27–53, 2000.
- [6] David B. Cohen. The large scale geometry of strongly aperiodic subshifts of finite type. *Advances in Mathematics*, 308:599–626, 2017.
- [7] David Carroll and Andrew Penland. Periodic points on shifts of finite type and commensurability invariants of groups. *New York Journal of Mathematics*, 21:811–822, 2015.
- [8] Tullio Ceccherini-Silberstein and Michel Coornaert. *Cellular Automata and Groups*. Springer, 2009.
- [9] Thomas Fernique. Multidimensional Sturmian Sequences and Generalized Substitutions. *International Journal of Foundations of Computer Science*, 17:575–600, 2006.
- [10] Su Gao, Steve Jackson, and Brandon Seward. A coloring property for countable groups. *Mathematical Proceedings of the Cambridge Philosophical Society*, 147:579–592, 11 2009.
- [11] Eli Glasner and Vladimir V. Uspenskij. Effective minimal subflows of bernoulli flows. *Proc. Am. Math. Soc.*, 137(9):3147–3154, 2009.
- [12] Gustav Hedlund and Marston Morse. Symbolic dynamics. *American Journal of Mathematics*, 60(4):815–866, 1938.
- [13] Mike Hochman. On the dynamics and recursive properties of multidimensional symbolic systems. *Inventiones Mathematicae*, 176(1):131–167, 2009.
- [14] Emmanuel Jeandel. Some notes about subshifts on groups. *arXiv preprint arXiv:1501.06831*, 2015.
- [15] Douglas A. Lind and Brian Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, 1995.
- [16] M. Lothaire. *Algebraic combinatorics on words*. Cambridge: Cambridge University Press, 2002.
- [17] Wilhelm Magnus, Abraham Karrass, and Donald Solitar. *Combinatorial Group Theory: Presentations of Groups in Terms of Generators and Relations*. Dover Books on Mathematics Series. Dover Publications, 2004.
- [18] N. Pytheas Fogg. *Substitutions in Dynamics, Arithmetics and Combinatorics*, volume 1794 of *Lecture Notes in Mathematics*. Springer, 2002.
- [19] Steven T. Piantadosi. Symbolic dynamics on free groups. Master’s thesis, University of North Carolina, Chapel Hill, 2006.
- [20] Steven T. Piantadosi. Symbolic dynamics on free groups. *Discrete and Continuous Dynamical Systems*, 20(3):725–738, 2008.
- [21] Ronnie Pavlov and Michael Schraudner. Classification of sofic projective subdynamics of multidimensional shifts of finite type. *Transactions of the American Mathematical Society*, 367(5), 2014.
- [22] Mark Pollicott and Michiko Yuri. *Dynamical Systems and Ergodic Theory*. London Mathematical Society Student Texts. Cambridge University Press, 1998.