

# MATH 110 Lecture 1.5

## The Limit of a Function

---

Edward Doolittle

Thursday, January 15, 2026

Department of Indigenous Knowledge and Science  
First Nations University of Canada

## The Limit of a Function

Functions with Holes

Limits

One-sided Limits

Infinite Limits

The Definition of a Limit

Examples and Exercises


# **The Limit of a Function**

---

# Functions with Holes

Consider the graph of the function

$$f(x) = 1$$



$$f(x) = 1$$

Every time I give an input to  $f$ , the function answers back with the number 1. So


- $f(7) = 1$
- $f(0.5) = 1$
- $f(1) = 1$
- $f(-\sqrt{2}) = 1$

and so on. Simple.

# Undefined values

Now consider the similar graph of the function

$$g(x) = \begin{cases} 1, & x \neq 2 \\ \text{undefined} & x = 2 \end{cases}$$



Most of the time, when I give an input to  $g$ , the function replies with the output 1. So,


- $g(7) = 1$
- $g(0.5) = 1$
- $g(1) = 1$
- $g(-\sqrt{2}) = 1$

as with  $f$ . However, if I give the input 2 to  $g$ , it doesn't respond at all.

# Domain of a Function

We say that the domain of  $g$  doesn't include 2; more precisely, the domain of  $g$  is the set

$$\{x \in \mathbb{R} : x \neq 2\}$$





## Defining an Undefined Value


If someone were to insist on us assigning a value to  $g(2)$ , we can say two things:

- Strictly speaking,  $g(2)$  is undefined!
- However, if you **insist**, the only value that makes sense for  $g(2)$  is  $g(2) = 1$ .

# The Concept of a Limit

The two statements on the previous slide are inconsistent. We can't have that in mathematics, so we work around it by saying

- $g(2)$  is undefined, but
- $\lim_{x \rightarrow 2} g(x) = 1$



## Another Example of Limits

Consider the function

$$h(x) = \frac{x(x-3)}{x-3}$$


The domain of the function is  $\{x \in \mathbb{R} : x \neq 3\}$ . So the answer to the question “What is  $h(3)$ ?” is, “It’s undefined”. But now, we can say more.

## Another Example of Limits

Consider the function

$$h(x) = \frac{x(x-3)}{x-3}$$

The domain of the function is  $\{x \in \mathbb{R} : x \neq 3\}$ . So the answer to the question “What is  $h(3)$ ?” is, “It’s undefined”. But now, we can say more. From the graph,




## Another Example of Limits

Consider the function

$$h(x) = \frac{x(x-3)}{x-3}$$

The domain of the function is  $\{x \in \mathbb{R} : x \neq 3\}$ . So the answer to the question “What is  $h(3)$ ?” is, “It’s undefined”. But now, we can say more. From the graph,


$$\lim_{x \rightarrow 3} h(x) = 3$$



## Three More Examples of Limits

Consider the function  $f(x)$  represented by the graph on the right.

For this function, we say that




## Three More Examples of Limits

Consider the function  $f(x)$  represented by the graph on the right.


For this function, we say that

$$\lim_{x \rightarrow 1} f(x) = 2$$



## Three More Examples of Limits

But what about this function?






## Three More Examples of Limits

But what about this function?

We can say that  $f(1) = 2$ , but can we say anything about limits?




## Three More Examples of Limits

But what about this function?

We can say that  $f(1) = 2$ , but can we say anything about limits?


Yes, in this case, we can still say

$$\lim_{x \rightarrow 1} f(x) = 2$$



## Three More Examples of Limits


And what about this function?



## Three More Examples of Limits

And what about this function?

In this case, we say that  $f(1) = 1$ , but can we say anything about limits?




## Three More Examples of Limits

And what about this function?

In this case, we say that  $f(1) = 1$ , but can we say anything about limits?




Yes, in this case, we still say

$$\lim_{x \rightarrow 1} f(x) = 2$$



# The Three Situations in which Limits Exist

The following three graphs illustrate the three situations in which we can talk about limits.



# Applications of Limits

- You may be wondering whether this idea of limits is really very useful.

# Applications of Limits

- You may be wondering whether this idea of limits is really very useful.
- But we have already seen an example where limits can help us:



# Applications of Limits


- You may be wondering whether this idea of limits is really very useful.
- But we have already seen an example where limits can help us:
- the tangent problem.

# Applications of Limits

- You may be wondering whether this idea of limits is really very useful.
- But we have already seen an example where limits can help us:
- the tangent problem.
- Limits can also help us solve the area problem, as we will see in chapter 5.


# Limits and the Tangent Problem

- To see how limits can help us solve the tangent problem, consider the graph of  $f(x) = x^2$  at the right.




# Limits and the Tangent Problem

- To see how limits can help us solve the tangent problem, consider the graph of  $f(x) = x^2$  at the right.
- We want to find the slope of the tangent line at the base point  $Q = (1, 1)$ .




# Limits and the Tangent Problem

- To see how limits can help us solve the tangent problem, consider the graph of  $f(x) = x^2$  at the right.
- We want to find the slope of the tangent line at the base point  $Q = (1, 1)$ .
- To find it, we find the slopes of the secant lines  $PQ$ , where  $P \neq Q$ .





# Limits and the Tangent Problem



- To see how limits can help us solve the tangent problem, consider the graph of  $f(x) = x^2$  at the right.
- We want to find the slope of the tangent line at the base point  $Q = (1, 1)$ .
- To find it, we find the slopes of the secant lines  $PQ$ , where  $P \neq Q$ .
- Let's graph the slope of the secant line as a function of the  $x$ -value of the second point  $P$ .



# The Tangent Problem for $f(x) = x^2$





# The Tangent Problem for $f(x) = x^2$









# The Tangent Problem for $f(x) = x^2$





# The Tangent Problem for $f(x) = x^2$





# The Tangent Problem for $f(x) = x^2$





# The Tangent Problem for $f(x) = x^2$





# The Tangent Problem for $f(x) = x^2$





# The Tangent Problem for $f(x) = x^2$



# The Tangent Problem for $f(x) = x^2$





# The Tangent Problem for $f(x) = x^2$









# The Tangent Problem for $f(x) = x^2$



# The Tangent Problem for $f(x) = x^2$




# The Tangent Problem for $f(x) = x^2$



## The Tangent Problem for $f(x) = x^2$

- The slope of the secant line appears to be a nice linear function of the  $x$ -value of  $P$ .
- However, there is one  $x$ -value of  $P$  which is not allowed:  $x = 1$ , where  $P = Q$ .
- But that is the most interesting  $x$ -value!
- We need to use limits to find slopes of tangents.



## Guessing Limits from Tabular Data

- We can see that calculating limits is a useful thing.

## Guessing Limits from Tabular Data

- We can see that calculating limits is a useful thing.
- However, for the time being, we will just guess limits based on the graphs of functions and/or tabular data.

## Guessing Limits from Tabular Data

- We can see that calculating limits is a useful thing.
- However, for the time being, we will just guess limits based on the graphs of functions and/or tabular data.
- See examples 1–5 in the textbook.

## Guessing Limits from Tabular Data

- We can see that calculating limits is a useful thing.
- However, for the time being, we will just guess limits based on the graphs of functions and/or tabular data.
- See examples 1–5 in the textbook.
- Note, however, that those methods are not 100% reliable. We need reliable ways of calculating limits.



## Guessing Limits from Tabular Data

- We can see that calculating limits is a useful thing.
- However, for the time being, we will just guess limits based on the graphs of functions and/or tabular data.
- See examples 1–5 in the textbook.
- Note, however, that those methods are not 100% reliable. We need reliable ways of calculating limits.
- We will learn more reliable methods for calculating limits in section 1.6.

# The Heaviside Step Function

Consider the function

$$H(x) = \begin{cases} 0, & x < 0 \\ 1, & x \geq 0 \end{cases}$$


This function was studied by electrical engineer Oliver Heaviside as a model for throwing a switch in an electric circuit.

# The Heaviside Step Function

Consider the function


$$H(x) = \begin{cases} 0, & x < 0 \\ 1, & x \geq 0 \end{cases}$$

This function was studied by electrical engineer Oliver Heaviside as a model for throwing a switch in an electric circuit.




# Limits for the Heaviside Step Function

- It is easy to see that  $\lim_{x \rightarrow 3} H(x) = 1$ ,  $\lim_{x \rightarrow -2} H(x) = 0$ , etc.




# Limits for the Heaviside Step Function

- It is easy to see that  $\lim_{x \rightarrow 3} H(x) = 1$ ,  $\lim_{x \rightarrow -2} H(x) = 0$ , etc.
- In general, we can say that  $\lim_{x \rightarrow a} H(x) = 1$  for  $a > 0$ , and  $\lim_{x \rightarrow a} H(x) = 0$  for  $a < 0$ .




# Limits for the Heaviside Step Function

- It is easy to see that  $\lim_{x \rightarrow 3} H(x) = 1$ ,  $\lim_{x \rightarrow -2} H(x) = 0$ , etc.
- In general, we can say that  $\lim_{x \rightarrow a} H(x) = 1$  for  $a > 0$ , and  $\lim_{x \rightarrow a} H(x) = 0$  for  $a < 0$ .
- But what about  $\lim_{x \rightarrow 0} H(x)$ ?




# Limits for the Heaviside Step Function

- It is easy to see that  $\lim_{x \rightarrow 3} H(x) = 1$ ,  $\lim_{x \rightarrow -2} H(x) = 0$ , etc.
- In general, we can say that  $\lim_{x \rightarrow a} H(x) = 1$  for  $a > 0$ , and  $\lim_{x \rightarrow a} H(x) = 0$  for  $a < 0$ .
- But what about  $\lim_{x \rightarrow 0} H(x)$ ?
- Since there is no sensible value we could give  $H(0)$ , we must say  $\lim_{x \rightarrow 0} H(x)$  is **undefined**.



# One-sided Limits for $H(x)$


- However, we can do better than that.






## One-sided Limits for $H(x)$

- However, we can do better than that.
- If we ignore everything to the left of 0 on the  $x$ -axis, there is a sensible value for  $H(0)$ , namely 1.




## One-sided Limits for $H(x)$

- However, we can do better than that.
- If we ignore everything to the left of 0 on the  $x$ -axis, there is a sensible value for  $H(0)$ , namely 1.
- We say that  
the limit of  $H(x)$  as  $x$  approaches 0 **from the right** is 1




# One-sided Limits for $H(x)$

- However, we can do better than that.
- If we ignore everything to the left of 0 on the  $x$ -axis, there is a sensible value for  $H(0)$ , namely 1.
- We say that  
the limit of  $H(x)$  as  $x$  approaches 0 **from the right** is 1
- In symbols, we write  $\lim_{x \rightarrow 0^+} H(x) = 1$ .




## One-sided Limits for $H(x)$

- We can do the same thing on the other side.



## One-sided Limits for $H(x)$


- We can do the same thing on the other side.
- If we ignore everything to the right of 0 on the  $x$ -axis, there is a sensible value for  $H(0)$ , namely 0.



## One-sided Limits for $H(x)$

- We can do the same thing on the other side.
- If we ignore everything to the right of 0 on the  $x$ -axis, there is a sensible value for  $H(0)$ , namely 0.
- We say that

the limit of  $H(x)$  as  $x$  approaches 0 **from the left** is 0




## One-sided Limits for $H(x)$

- We can do the same thing on the other side.
- If we ignore everything to the right of 0 on the  $x$ -axis, there is a sensible value for  $H(0)$ , namely 0.
- We say that

the limit of  $H(x)$  as  $x$  approaches 0 **from the left** is 0

- In symbols, we write  $\lim_{x \rightarrow 0^-} H(x) = 0$ .



## One-sided Limits and Two-sided Limits

- If a function has an ordinary limit at  $a$  (sometimes called a **two-sided** limit at  $a$ ), then it has both one-sided limits at  $a$ .



## One-sided Limits and Two-sided Limits

- If a function has an ordinary limit at  $a$  (sometimes called a **two-sided** limit at  $a$ ), then it has both one-sided limits at  $a$ .
- Conversely, if a function has both one-sided limits at a point  $a$ , **and those one-sided limits are equal**, it has a two-sided limit at  $a$ .

## One-sided Limits and Two-sided Limits

- If a function has an ordinary limit at  $a$  (sometimes called a **two-sided** limit at  $a$ ), then it has both one-sided limits at  $a$ .
- Conversely, if a function has both one-sided limits at a point  $a$ , **and those one-sided limits are equal**, it has a two-sided limit at  $a$ .
- To be precise, we have

$$\lim_{x \rightarrow a} f(x) = L$$

if and only if

$$\lim_{x \rightarrow a^-} f(x) = L \text{ and } \lim_{x \rightarrow a^+} f(x) = L.$$

## One-sided Limits and Two-sided Limits

- If a function has an ordinary limit at  $a$  (sometimes called a **two-sided** limit at  $a$ ), then it has both one-sided limits at  $a$ .
- Conversely, if a function has both one-sided limits at a point  $a$ , **and those one-sided limits are equal**, it has a two-sided limit at  $a$ .
- To be precise, we have

$$\lim_{x \rightarrow a} f(x) = L$$

if and only if

$$\lim_{x \rightarrow a^-} f(x) = L \text{ and } \lim_{x \rightarrow a^+} f(x) = L.$$


- So one way to show that a (two-sided) limit does not exist is to show that the corresponding one-sided limits are not equal.

## Ways in which Limits Fail to Exist

- A limit can fail to exist for several reasons.


# Ways in which Limits Fail to Exist

- A limit can fail to exist for several reasons.
- One way, which we've seen already, is for a function to behave differently on the left and on the right of our point of interest.




## Ways in which Limits Fail to Exist

- A limit can fail to exist for several reasons.
- One way, which we've seen already, is for a function to behave differently on the left and on the right of our point of interest.
- In other words, the left-hand limit differs from the right-hand limit.



# Ways in which Limits Fail to Exist

- A limit can fail to exist for several reasons.
- One way, which we've seen already, is for a function to behave differently on the left and on the right of our point of interest.
- In other words, the left-hand limit differs from the right-hand limit.
- Another way for a limit to fail to exist is where  $f$  gets really large as  $x$  approaches some value.




# Infinite Limits

- In this kind of a situation, we write

$$\lim_{x \rightarrow 0} f(x) = \infty$$

and say “the limit as  $x$  approaches 0 of  $f(x)$  is infinity”.





- In this kind of a situation, we write

$$\lim_{x \rightarrow 0} f(x) = \infty$$

and say “the limit as  $x$  approaches 0 of  $f(x)$  is infinity”.

- That expression is just an idiom;  $\infty$  is not a number.

- In this kind of a situation, we write

$$\lim_{x \rightarrow 0} f(x) = \infty$$

and say “the limit as  $x$  approaches 0 of  $f(x)$  is infinity”.

- That expression is just an idiom;  $\infty$  is not a number.
- Despite the notation saying that the limit equals something, the limit does not exist.

- In this kind of a situation, we write

$$\lim_{x \rightarrow 0} f(x) = \infty$$

and say “the limit as  $x$  approaches 0 of  $f(x)$  is infinity”.

- That expression is just an idiom;  $\infty$  is not a number.
- Despite the notation saying that the limit equals something, the limit does not exist.

- In this kind of a situation, we write


$$\lim_{x \rightarrow 0} f(x) = \infty$$

and say “the limit as  $x$  approaches 0 of  $f(x)$  is infinity”.

- That expression is just an idiom;  $\infty$  is not a number.
- Despite the notation saying that the limit equals something, the limit does not exist.
- Writing this as a limit expression just means that the limit fails to exist in a particular way.

# Negative and One-sided Infinite Limits


There are six variations on infinite limits.



$$\lim_{x \rightarrow 0} f(x) = \infty$$

# Negative and One-sided Infinite Limits


There are six variations on infinite limits.



$$\lim_{x \rightarrow 0^-} f(x) = \infty$$

# Negative and One-sided Infinite Limits


There are six variations on infinite limits.



$$\lim_{x \rightarrow 0^+} f(x) = \infty$$

# Negative and One-sided Infinite Limits

There are six variations on infinite limits.




$$\lim_{x \rightarrow 0} f(x) = -\infty$$



# Negative and One-sided Infinite Limits


There are six variations on infinite limits.



$$\lim_{x \rightarrow 0^-} f(x) = -\infty$$

# Negative and One-sided Infinite Limits

There are six variations on infinite limits.



$$\lim_{x \rightarrow 0^+} f(x) = -\infty$$

## Identifying Infinite Limits

- It will generally be easy to identify locations where a limit could be infinite.

## Identifying Infinite Limits

- It will generally be easy to identify locations where a limit could be infinite.
- Infinite limits typically occur for functions  $f$  which are of the form

$$f(x) = \frac{g(x)}{h(x)},$$

in other words, quotients of two other functions.

## Identifying Infinite Limits

- It will generally be easy to identify locations where a limit could be infinite.
- Infinite limits typically occur for functions  $f$  which are of the form

$$f(x) = \frac{g(x)}{h(x)},$$

in other words, quotients of two other functions.

- Infinite limits may occur at the values  $a$  where  $h(a) = 0$ . Notice that these values  $a$  are not in the domain of  $f$ .

# Identifying Infinite Limits

- It will generally be easy to identify locations where a limit could be infinite.
- Infinite limits typically occur for functions  $f$  which are of the form

$$f(x) = \frac{g(x)}{h(x)},$$

in other words, quotients of two other functions.

- Infinite limits may occur at the values  $a$  where  $h(a) = 0$ . Notice that these values  $a$  are not in the domain of  $f$ .
- We determine which type of limit occurs at  $a$  by investigating the size and sign of  $f$  on each side of  $a$ .

# Vertical Asymptotes

The line  $x = a$  is called a **vertical asymptote** of the curve  $y = f(x)$  if at least one of the following statements is true:

$$\lim_{x \rightarrow a} f(x) = \infty \quad \lim_{x \rightarrow a^-} f(x) = \infty \quad \lim_{x \rightarrow a^+} f(x) = \infty$$

$$\lim_{x \rightarrow a} f(x) = -\infty \quad \lim_{x \rightarrow a^-} f(x) = -\infty \quad \lim_{x \rightarrow a^+} f(x) = -\infty$$

In other words, if any of the limits as  $x$  approaches  $a$  from the left or the right is  $\pm\infty$ , we say that  $y = f(x)$  has a vertical asymptote at  $a$ .

# The Definition of a Limit

- We have been using an intuitive concept of what limits mean.



# The Definition of a Limit

- We have been using an intuitive concept of what limits mean.
- However, the definition is not good enough for serious mathematics.

# The Definition of a Limit

- We have been using an intuitive concept of what limits mean.
- However, the definition is not good enough for serious mathematics.
- A more rigorous approach to limits was formulated by Cauchy.

# The Definition of a Limit

- We have been using an intuitive concept of what limits mean.
- However, the definition is not good enough for serious mathematics.
- A more rigorous approach to limits was formulated by Cauchy.
- Cauchy's definition is based on the idea of controlling the range of values a function can take by limiting the domain.

# The Definition of a Limit

- We have been using an intuitive concept of what limits mean.
- However, the definition is not good enough for serious mathematics.
- A more rigorous approach to limits was formulated by Cauchy.
- Cauchy's definition is based on the idea of controlling the range of values a function can take by limiting the domain.
- Cauchy's definition is still somewhat vague. A completely satisfactory definition is Weierstrass's epsilon-delta definition, which can be found in section 1.7. We will not be studying the epsilon-delta definition in this course.

# The Definition of an Ordinary Limit

- We write

$$\lim_{x \rightarrow a} f(x) = L$$

and say

the limit as  $x$  approaches  $a$  of  $f(x)$  is  $L$

if we can make the values of  $f(x)$  as close to  $L$  as we like by taking  $x$  to be sufficiently close to **(but not equal to)**  $a$ .

# The Definition of a One-sided Limit

- We write

$$\lim_{x \rightarrow a^-} f(x) = L$$

and say

the limit as  $x$  approaches  $a$  from the left of  $f(x)$  is  $L$   
if we can make the values of  $f(x)$  as close to  $L$  as we like by  
taking  $x$  to be sufficiently close to **(but less than)**  $a$ .

# The Definition of a One-sided Limit

- We write

$$\lim_{x \rightarrow a^-} f(x) = L$$

and say

the limit as  $x$  approaches  $a$  from the left of  $f(x)$  is  $L$

if we can make the values of  $f(x)$  as close to  $L$  as we like by taking  $x$  to be sufficiently close to **(but less than)**  $a$ .

- There is a similar definition for the right-hand limit

$$\lim_{x \rightarrow a^+} f(x) = L. \text{ (Try writing out the definition yourself.)}$$

## The Definition of an Infinite Limit

- Let  $f$  be a function defined on both sides of  $a$ , except possibly at  $a$  itself. Then we can write

$$\lim_{x \rightarrow a} f(x) = \infty$$

and say

the limit as  $x$  approaches  $a$  of  $f(x)$  is infinity

or

$f(x)$  increases without bound as  $x$  approaches  $a$

if  $f(x)$  can be made as large as we like by taking  $x$  sufficiently close to **(but not equal to)**  $a$ .



## The Definition of an Infinite Limit

- Let  $f$  be a function defined on both sides of  $a$ , except possibly at  $a$  itself. Then we can write

$$\lim_{x \rightarrow a} f(x) = \infty$$

and say

the limit as  $x$  approaches  $a$  of  $f(x)$  is infinity

or

$f(x)$  increases without bound as  $x$  approaches  $a$

if  $f(x)$  can be made as large as we like by taking  $x$  sufficiently close to **(but not equal to)**  $a$ .

- There are similar definitions for  $\lim_{x \rightarrow a} f(x) = -\infty$  and for the corresponding one-sided limits. (Try writing out the definitions yourself.)

## Examples

1. Sketch the graph of the following function and use it to determine the values of  $a$  for which  $\lim_{x \rightarrow a} f(x)$  exists:

$$f(x) = \begin{cases} 2 - x & \text{if } x < -1 \\ x & \text{if } -1 \leq x < 1 \\ (x - 1)^2 & \text{if } 1 \leq x \end{cases}$$

2. Use a table of values to estimate the value of the following limits.

$$2.1 \quad \lim_{x \rightarrow 0} \frac{\sqrt{x+4} - 2}{x}$$

$$2.2 \quad \lim_{x \rightarrow 0} \frac{\tan 3x}{\tan 5x}$$

## Examples

3. Let

$$f(x) = \begin{cases} \sqrt{-x} & \text{if } x < 0 \\ 3 - x & \text{if } 0 \leq x < 3 \\ (x - 3)^2 & \text{if } 3 \leq x \end{cases}$$

Determine whether the limits  $\lim_{x \rightarrow 0} f(x)$  and  $\lim_{x \rightarrow 3} f(x)$  exist. If so, find the values of those limits.

4. Find the vertical asymptotes of the function

$$y = \frac{x}{x^2 + x - 2}$$

## Exercises

Now you should work on Problem Set 1.5a and Problem Set 1.15b. After you have finished them, you should try the following additional exercises from Section 1.5 for two-sided limits:

1.5 C-level: 1, 19–22, 23, 26, 35;

B-level: 32–44;

A-level: 49

and the following additional exercises from Section 1.5 for one-sided and infinite limits:

1.5 C-level: 2–3, 4–5, 7, 8–9, 10, 11–12, 15–18, 24–25, 29–34, 38–39, 40;

B-level: 6, 11, 13–14, 35–37, 41, 46;

A-level: 47–48