

MATH 110 Lecture 2.8

Related Rates

Edward Doolittle

Thursday, February 26, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Related Rates

Relations in Implicit Form

Relations with 3 non-Time Variables

Examples and Exercises

Related Rates

Related Rates

- Related rates problems are word problems which can be solved by taking first derivatives.

Related Rates

- Related rates problems are word problems which can be solved by taking first derivatives.
- Related rates problems have a reputation of being difficult.

Related Rates

- Related rates problems are word problems which can be solved by taking first derivatives.
- Related rates problems have a reputation of being difficult.
- Until now, the skills you have been learning are mostly technical.

Related Rates

- Related rates problems are word problems which can be solved by taking first derivatives.
- Related rates problems have a reputation of being difficult.
- Until now, the skills you have been learning are mostly technical.
- Related rates problems require more conceptual understanding, including the ability to develop your own formulas, but are technically easy.

Related Rates

- Related rates problems are word problems which can be solved by taking first derivatives.
- Related rates problems have a reputation of being difficult.
- Until now, the skills you have been learning are mostly technical.
- Related rates problems require more conceptual understanding, including the ability to develop your own formulas, but are technically easy.
- With practice, related rates problems will become easier.

Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.

Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.

Colony of Bacteria Problem


- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”

Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”
- It's not true that the area is increasing at a constant rate!


Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”
- It's not true that the area is increasing at a constant rate!




Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”
- It's not true that the area is increasing at a constant rate!




Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”
- It's not true that the area is increasing at a constant rate!




Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”
- It's not true that the area is increasing at a constant rate!




Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”
- It's not true that the area is increasing at a constant rate!



Colony of Bacteria Problem

- As a simple example, consider a roughly circular shaped colony of bacteria growing in a petri dish. The colony has initial radius close to 0 cm.
- We make periodic measurements of the diameter of the colony and find that the radius is increasing at a constant rate of 0.2 cm per hour.
- The real question we have, though, is “How fast is the area of the colony is increasing after 5 hours?”
- It's not true that the area is increasing at a constant rate!



Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:
 1. The variables are time t , radius $r(t)$, and area $A(t)$.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:
 1. The variables are time t , radius $r(t)$, and area $A(t)$.
 2. The relation between r and A is $A(t) = \pi (r(t))^2$.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:
 1. The variables are time t , radius $r(t)$, and area $A(t)$.
 2. The relation between r and A is $A(t) = \pi (r(t))^2$.
 3. $dA/dt = 2\pi r dr/dt$

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:
 1. The variables are time t , radius $r(t)$, and area $A(t)$.
 2. The relation between r and A is $A(t) = \pi (r(t))^2$.
 3. $dA/dt = 2\pi r dr/dt$
 4. At $t = 5$, $dr/dt = 0.2$,

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:
 1. The variables are time t , radius $r(t)$, and area $A(t)$.
 2. The relation between r and A is $A(t) = \pi (r(t))^2$.
 3. $dA/dt = 2\pi r dr/dt$
 4. At $t = 5$, $dr/dt = 0.2$, and $r = 0.2 \times 5 = 1$.

Colony of Bacteria Solution

- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:
 1. The variables are time t , radius $r(t)$, and area $A(t)$.
 2. The relation between r and A is $A(t) = \pi (r(t))^2$.
 3. $dA/dt = 2\pi r dr/dt$
 4. At $t = 5$, $dr/dt = 0.2$, and $r = 0.2 \times 5 = 1$.
 5. $dA/dt = 2\pi \cdot 1 \cdot 0.2 \approx 1.26 \text{ cm}^2/\text{hr}$.

Colony of Bacteria Solution


- In order to solve this problem, we must take each of the following steps.
 1. Identify the variables in the problem, including time.
 2. Find a **relation** between the non-time variables.
 3. Differentiate the relation.
 4. Fill in all the numbers we know.
 5. Solve for the quantity we don't know.
- We must take these steps in the correct order, or we will get the wrong answer.
- In the colony of bacteria problem, the steps are as follows:
 1. The variables are time t , radius $r(t)$, and area $A(t)$.
 2. The relation between r and A is $A(t) = \pi (r(t))^2$.
 3. $dA/dt = 2\pi r \, dr/dt$
 4. At $t = 5$, $dr/dt = 0.2$, and $r = 0.2 \times 5 = 1$.
 5. $dA/dt = 2\pi \cdot 1 \cdot 0.2 \approx 1.26 \text{ cm}^2/\text{hr}$.
- Note that we differentiated *before* we substituted any numbers.

Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?


Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?
- The variables are time, x , and y .




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?
- The variables are time, x , and y .
- The relation is $x^2 + y^2 = 10^2$.




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?
- The variables are time, x , and y .
- The relation is $x^2 + y^2 = 10^2$.
- Differentiating, $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$.




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?
- The variables are time, x , and y .
- The relation is $x^2 + y^2 = 10^2$.
- Differentiating, $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$.
- Filling in what we know, $\frac{dy}{dt} = -2$, $x = 8$.




Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?
- The variables are time, x , and y .
- The relation is $x^2 + y^2 = 10^2$.
- Differentiating, $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$.
- Filling in what we know, $\frac{dy}{dt} = -2$, $x = 8$.
- We are not given y but we can figure it out: $x^2 + y^2 = 10^2$ and $x = 8$ give $y = 6$.



Relations in Implicit Form

- Consider the following problem: A 10 ft ladder, propped against a wall, slides down at a rate of 2 ft/s. How fast is the bottom of the ladder moving when the bottom is 8 ft from the wall?
- The variables are time, x , and y .
- The relation is $x^2 + y^2 = 10^2$.
- Differentiating, $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$.
- Filling in what we know, $\frac{dy}{dt} = -2$, $x = 8$.
- We are not given y but we can figure it out: $x^2 + y^2 = 10^2$ and $x = 8$ give $y = 6$.
- So $16 \frac{dx}{dt} + 12 \cdot -2 = 0$, i.e., $\frac{dx}{dt} = 3/2$ ft/s.



Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?

Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- The variables are time, the height h of the water, the radius r of the surface of the water, and V the volume of the water.

Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- The variables are time, the height h of the water, the radius r of the surface of the water, and V the volume of the water.
- The relation is $V = \frac{1}{3}\pi r^2 h$.

Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- The variables are time, the height h of the water, the radius r of the surface of the water, and V the volume of the water.
- The relation is $V = \frac{1}{3}\pi r^2 h$.
- Note that $V(t)$, $r(t)$, and $h(t)$ are all dependent variables.

Eliminating a non-Time Variable

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?

Eliminating a non-Time Variable

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- We need to reduce the number of variables. We can either try to reduce the number of variables before or after we differentiate.

Eliminating a non-Time Variable

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- We need to reduce the number of variables. We can either try to reduce the number of variables before or after we differentiate.
- Let's try to reduce the number of variables before we differentiate.

Eliminating a non-Time Variable

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- We need to reduce the number of variables. We can either try to reduce the number of variables before or after we differentiate.
- Let's try to reduce the number of variables before we differentiate.
- We need *another relation*.

Eliminating a non-Time Variable

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- We need to reduce the number of variables. We can either try to reduce the number of variables before or after we differentiate.
- Let's try to reduce the number of variables before we differentiate.
- We need *another relation*.
- Similar triangles give $r/3 = h/10$.

Eliminating a non-Time Variable

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- We need to reduce the number of variables. We can either try to reduce the number of variables before or after we differentiate.
- Let's try to reduce the number of variables before we differentiate.
- We need *another relation*.
- Similar triangles give $r/3 = h/10$.
- Then $r^2 = 9h/10$ so $V = \frac{3\pi}{100}h^3$.

Solving the Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?

Solving the Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- Now $\frac{dV}{dt} = \frac{9\pi}{100}h^2 \frac{dh}{dt}$.

Solving the Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?
- Now $\frac{dV}{dt} = \frac{9\pi}{100}h^2 \frac{dh}{dt}$.
- We know $\frac{dV}{dt} = 2$ and $h = 5$.

Solving the Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?

- Now $\frac{dV}{dt} = \frac{9\pi}{100}h^2 \frac{dh}{dt}$.

- We know $\frac{dV}{dt} = 2$ and $h = 5$.

- Substituting the values we know, we have

$$2 = \frac{9\pi}{100}5^2 \frac{dh}{dt}.$$

Solving the Water Level Problem

- A paper cup has the shape of a cone with height 10 cm and radius 3 cm (at the top). If water is poured into the cup at a rate of $2 \text{ cm}^3/\text{s}$, how fast is the water level rising when the water is 5 cm deep?

- Now $\frac{dV}{dt} = \frac{9\pi}{100}h^2 \frac{dh}{dt}$.

- We know $\frac{dV}{dt} = 2$ and $h = 5$.

- Substituting the values we know, we have

$$2 = \frac{9\pi}{100}5^2 \frac{dh}{dt}.$$

- Solving, $\frac{dh}{dt} = \frac{8}{9\pi}$.

Two Moving Objects

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?

Two Moving Objects

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- Let O be the point on the road directly under the balloon.

Two Moving Objects

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- Let O be the point on the road directly under the balloon.
- The variables are t , $x(t)$, $y(t)$, and $z(t)$.

Two Moving Objects

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- Let O be the point on the road directly under the balloon.
- The variables are t , $x(t)$, $y(t)$, and $z(t)$.
- The relation is $x^2 + y^2 = z^2$.

Two Moving Objects

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- Let O be the point on the road directly under the balloon.
- The variables are t , $x(t)$, $y(t)$, and $z(t)$.
- The relation is $x^2 + y^2 = z^2$.
- We could try to eliminate variables using our knowledge of the motion, but let's try another approach.

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s.
A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- The relation is $x^2 + y^2 = z^2$.

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- The relation is $x^2 + y^2 = z^2$.
- Differentiate: $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 2z \frac{dz}{dt}$.

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- The relation is $x^2 + y^2 = z^2$.
- Differentiate: $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 2z \frac{dz}{dt}$.
- One equation, six variables; we need values for five variables before we can find $\frac{dz}{dt}$.

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- The relation is $x^2 + y^2 = z^2$.
- Differentiate: $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 2z \frac{dz}{dt}$.
- One equation, six variables; we need values for five variables before we can find $\frac{dz}{dt}$.
- We are given $\frac{dx}{dt} = 15$ and $\frac{dy}{dt} = 5$. We still need x , y , and z .

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s. A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s.

A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?

- $x \frac{dx}{dt} + y \frac{dy}{dt} = z \frac{dz}{dt}$, $\frac{dx}{dt} = 15$, and $\frac{dy}{dt} = 5$

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s.
A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?
- $x \frac{dx}{dt} + y \frac{dy}{dt} = z \frac{dz}{dt}$, $\frac{dx}{dt} = 15$, and $\frac{dy}{dt} = 5$
- After 3 s the boy has traveled $x = 3 \cdot 15 = 45$ ft.

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s.

A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?

- $x \frac{dx}{dt} + y \frac{dy}{dt} = z \frac{dz}{dt}$, $\frac{dx}{dt} = 15$, and $\frac{dy}{dt} = 5$
- After 3 s the boy has traveled $x = 3 \cdot 15 = 45$ ft.
- After 3 s the balloon has risen $3 \times 5 = 15$ ft, total height $y = 45 + 15 = 60$ ft.

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s.

A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?

- $x \frac{dx}{dt} + y \frac{dy}{dt} = z \frac{dz}{dt}$, $\frac{dx}{dt} = 15$, and $\frac{dy}{dt} = 5$
- After 3 s the boy has traveled $x = 3 \cdot 15 = 45$ ft.
- After 3 s the balloon has risen $3 \times 5 = 15$ ft, total height $y = 45 + 15 = 60$ ft.
- z satisfies $z^2 = 45^2 + 60^2 = 5625$ so $z = 75$.

Retaining All non-Time Variables

- A balloon is rising at a constant speed of 5 ft/s.

A boy is cycling along a straight road at a speed of 15 ft/s. When he passes under the balloon, it is 45 ft above him. How fast is the distance between the boy and the balloon increasing 3 s later?

- $x \frac{dx}{dt} + y \frac{dy}{dt} = z \frac{dz}{dt}$, $\frac{dx}{dt} = 15$, and $\frac{dy}{dt} = 5$
- After 3 s the boy has traveled $x = 3 \cdot 15 = 45$ ft.
- After 3 s the balloon has risen $3 \times 5 = 15$ ft, total height $y = 45 + 15 = 60$ ft.
- z satisfies $z^2 = 45^2 + 60^2 = 5625$ so $z = 75$.
- Solving, $\frac{dz}{dt} = \frac{1}{75} (45 \cdot 15 + 60 \cdot 5) = 13$ ft/s.

Examples and Exercises

Examples

1. The volume of a cube is increasing at a rate of $10 \text{ cm}^3/\text{min}$. How fast is the surface area increasing when the length of an edge is 30 cm ?
2. A waterskier skis over a ramp in the shape of a right-angled wedge 15 ft long, 4 ft above the water at the end. How fast is she rising when she leaves the ramp?
3. The angle of elevation of the sun is decreasing at a rate of 0.25 rad/h . How fast is the shadow cast by a 400 ft tall building increasing when the angle of elevation of the sun is $\pi/6$?

Now you should work on Problem Set 2.8. After you have finished it, you should try the following additional exercises from Section 2.8:

2.8 C-level: 1–35;

B-level: 36–40;

A-level: 41–50