

MATH 110 Lecture 2.1

Derivatives and Rates of Change

Edward Doolittle

Tuesday, January 27, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Derivatives and Rates of Change

Tangents

Velocities


Derivatives

Examples and Exercises

Derivatives and Rates of Change


Tangents

- The problem that we were originally considering was that of constructing tangents through a point $P(a, f(a))$ on the graph of f .




Tangents

- The problem that we were originally considering was that of constructing tangents through a point $P(a, f(a))$ on the graph of f .
- We did that by finding secant lines through P and $Q(x, f(x))$ and letting x approach a .




Tangents

- The problem that we were originally considering was that of constructing tangents through a point $P(a, f(a))$ on the graph of f .
- We did that by finding secant lines through P and $Q(x, f(x))$ and letting x approach a .




Tangents

- The problem that we were originally considering was that of constructing tangents through a point $P(a, f(a))$ on the graph of f .
- We did that by finding secant lines through P and $Q(x, f(x))$ and letting x approach a .




Tangents

- The problem that we were originally considering was that of constructing tangents through a point $P(a, f(a))$ on the graph of f .
- We did that by finding secant lines through P and $Q(x, f(x))$ and letting x approach a .




Tangents

- The problem that we were originally considering was that of constructing tangents through a point $P(a, f(a))$ on the graph of f .
- We did that by finding secant lines through P and $Q(x, f(x))$ and letting x approach a .



Tangents

- The problem that we were originally considering was that of constructing tangents through a point $P(a, f(a))$ on the graph of f .
- We did that by finding secant lines through P and $Q(x, f(x))$ and letting x approach a .
- We found the slope of the tangent line as a limit. The tangent line could then be described in point-slope form
$$y - y_0 = m(x - x_0).$$



The Definition of the Tangent Line

With the limit technology we developed in the previous chapter, we are now ready to give a formal definition of the tangent to a graph at a point.

The Definition of the Tangent Line

With the limit technology we developed in the previous chapter, we are now ready to give a formal definition of the tangent to a graph at a point.

Definition: The **tangent line** to the curve $y = f(x)$ at the point $P(a, f(a))$ is the line through P with the slope

$$m = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

provided that the limit exists.

The Definition of the Tangent Line

With the limit technology we developed in the previous chapter, we are now ready to give a formal definition of the tangent to a graph at a point.

Definition: The **tangent line** to the curve $y = f(x)$ at the point $P(a, f(a))$ is the line through P with the slope

$$m = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

provided that the limit exists.

If the above limit does not exist, the tangent line doesn't exist (the case where the limit is infinite is an exception that we'll study later).

Alternative Expressions for the Tangent

- In words, we can say

The tangent line at $P(a, f(a))$ is the line through P with slope equal to the limit of the slopes of the secant lines through $P(a, f(a))$ and $Q(x, f(x))$ as x approaches a .

Alternative Expressions for the Tangent

- In words, we can say
The tangent line at $P(a, f(a))$ is the line through P with slope equal to the limit of the slopes of the secant lines through $P(a, f(a))$ and $Q(x, f(x))$ as x approaches a .
- Note that an equation of the tangent line in point-slope form is

$$y - f(a) = m(x - a)$$

$$\text{where } m = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}.$$

Alternative Expressions for the Tangent

- In words, we can say
The tangent line at $P(a, f(a))$ is the line through P with slope equal to the limit of the slopes of the secant lines through $P(a, f(a))$ and $Q(x, f(x))$ as x approaches a .
- Note that an equation of the tangent line in point-slope form is

$$y - f(a) = m(x - a)$$

$$\text{where } m = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}.$$

- An alternate expression for the slope of the tangent line may be found by changing variables. Let $h = x - a$. Then we can write

Alternative Expressions for the Tangent

- In words, we can say

The tangent line at $P(a, f(a))$ is the line through P with slope equal to the limit of the slopes of the secant lines through $P(a, f(a))$ and $Q(x, f(x))$ as x approaches a .

- Note that an equation of the tangent line in point-slope form is

$$y - f(a) = m(x - a)$$

where $m = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$.

- An alternate expression for the slope of the tangent line may be found by changing variables. Let $h = x - a$. Then we can write

$$m = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

The Solution to the Tangent Problem

- We find tangent lines at a by finding a point on the line and the slope of the line.

The Solution to the Tangent Problem

- We find tangent lines at a by finding a point on the line and the slope of the line.
- The point on the line is easy: finding $P(a, f(a))$ just requires that we evaluate f at a .

The Solution to the Tangent Problem

- We find tangent lines at a by finding a point on the line and the slope of the line.
- The point on the line is easy: finding $P(a, f(a))$ just requires that we evaluate f at a .
- The slope m is harder: it is calculated as a limit

$$m = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

The Solution to the Tangent Problem

- We find tangent lines at a by finding a point on the line and the slope of the line.
- The point on the line is easy: finding $P(a, f(a))$ just requires that we evaluate f at a .
- The slope m is harder: it is calculated as a limit

$$m = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

- However, using the limit laws from the previous chapter, we will be able to calculate all such limits.

The Solution to the Tangent Problem

- We find tangent lines at a by finding a point on the line and the slope of the line.
- The point on the line is easy: finding $P(a, f(a))$ just requires that we evaluate f at a .
- The slope m is harder: it is calculated as a limit

$$m = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

- However, using the limit laws from the previous chapter, we will be able to calculate all such limits.
- In summary, we have built up a system for completely solving the tangent problem!

Comparison of Tangent and Velocity Problems

concept	tangent problem	velocity problem
independent variable	x	t

Comparison of Tangent and Velocity Problems

concept	tangent problem	velocity problem
independent variable	x	t
dependent variable	$f(x)$	$s(t)$

Comparison of Tangent and Velocity Problems

concept	tangent problem	velocity problem
independent variable	x	t
dependent variable	$f(x)$	$s(t)$
run	$\Delta x = x - a = h$	$\Delta t = t - a = h$

Comparison of Tangent and Velocity Problems

concept	tangent problem	velocity problem
independent variable	x	t
dependent variable	$f(x)$	$s(t)$
run	$\Delta x = x - a = h$	$\Delta t = t - a = h$
rise	$\Delta f = f(x) - f(a)$ $= f(a + h) - f(a)$	$\Delta s = s(t) - s(a)$ $= s(a + h) - s(a)$

Comparison of Tangent and Velocity Problems

concept	tangent problem	velocity problem
independent variable	x	t
dependent variable	$f(x)$	$s(t)$
run	$\Delta x = x - a = h$	$\Delta t = t - a = h$
rise	$\Delta f = f(x) - f(a)$ $= f(a + h) - f(a)$	$\Delta s = s(t) - s(a)$ $= s(a + h) - s(a)$
quotient	$\frac{\Delta f}{\Delta x} = \frac{f(a + h) - f(a)}{h}$ slope of secant	$\frac{\Delta s}{\Delta t} = \frac{s(a + h) - s(a)}{h}$ average velocity

Comparison of Tangent and Velocity Problems

concept	tangent problem	velocity problem
independent variable	x	t
dependent variable	$f(x)$	$s(t)$
run	$\Delta x = x - a = h$	$\Delta t = t - a = h$
rise	$\Delta f = f(x) - f(a)$ $= f(a + h) - f(a)$	$\Delta s = s(t) - s(a)$ $= s(a + h) - s(a)$
quotient	$\frac{\Delta f}{\Delta x} = \frac{f(a + h) - f(a)}{h}$ slope of secant	$\frac{\Delta s}{\Delta t} = \frac{s(a + h) - s(a)}{h}$ average velocity
limit as $h \rightarrow 0$	$\lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$ slope of tangent	$\lim_{h \rightarrow 0} \frac{s(a + h) - s(a)}{h}$ instantaneous velocity

The Formal Definition of Instantaneous Velocity

The above discussion leads to another new definition.

The Formal Definition of Instantaneous Velocity

The above discussion leads to another new definition.

Definition: The **instantaneous velocity** of a particle with position given by $s(t)$ at time $t = a$ is

$$v(a) = \lim_{h \rightarrow 0} \frac{s(a+h) - s(a)}{h}$$

provided that the limit exists.

The Formal Definition of Instantaneous Velocity

The above discussion leads to another new definition.

Definition: The **instantaneous velocity** of a particle with position given by $s(t)$ at time $t = a$ is

$$v(a) = \lim_{h \rightarrow 0} \frac{s(a+h) - s(a)}{h}$$

provided that the limit exists.

The instantaneous velocity is the limit of the average velocities as the length of the time interval $\Delta t = h = t - a$ on which the average velocities are taken approaches 0.

Derivatives

- We have seen a certain type of expression occurring in two different kinds of problems, the tangent problem and the velocity problem.

Derivatives

- We have seen a certain type of expression occurring in two different kinds of problems, the tangent problem and the velocity problem.
- The same kind of expression occurs in other problems involving rates of change (rates of chemical reactions, marginal cost in economics, etc.).

Derivatives

- We have seen a certain type of expression occurring in two different kinds of problems, the tangent problem and the velocity problem.
- The same kind of expression occurs in other problems involving rates of change (rates of chemical reactions, marginal cost in economics, etc.).
- Since it is so common, we give it a name:

Derivatives

- We have seen a certain type of expression occurring in two different kinds of problems, the tangent problem and the velocity problem.
- The same kind of expression occurs in other problems involving rates of change (rates of chemical reactions, marginal cost in economics, etc.).
- Since it is so common, we give it a name:

Definition: The **derivative of a function f at a number a** is denoted by $f'(a)$ and given by the formula

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

if this limit exists.

Expressions for the Derivative and Tangent Problem

- Alternatively, we can write

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

if this limit exists.

Expressions for the Derivative and Tangent Problem

- Alternatively, we can write

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

if this limit exists. However, I prefer the expression involving h for reasons we will see in section 3.2.

Expressions for the Derivative and Tangent Problem

- Alternatively, we can write

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

if this limit exists. However, I prefer the expression involving h for reasons we will see in section 3.2.

- Our new concept gives us an easy way of expressing the solution to the tangent problem:

The tangent line to $y = f(x)$ at the point $P(a, f(a))$ is the line through P with slope $f'(a)$.

Expressions for the Derivative and Tangent Problem

- Alternatively, we can write

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

if this limit exists. However, I prefer the expression involving h for reasons we will see in section 3.2.

- Our new concept gives us an easy way of expressing the solution to the tangent problem:

The tangent line to $y = f(x)$ at the point $P(a, f(a))$ is the line through P with slope $f'(a)$.

- We can write the equation of the tangent line in a single expression:

$$y - f(a) = f'(a)(x - a)$$

Expressions for the Velocity Problem

- We also have an alternate way to express the solution of the velocity problem

Expressions for the Velocity Problem

- We also have an alternate way to express the solution of the velocity problem
- We often write Δt for h . In that notation, the average velocity of a particle at time a over a time interval of length Δt is $\Delta s / \Delta t$, and we have

The instantaneous velocity of a particle at time a is

$$v(a) = \lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = s'(a)$$

Expressions for the Velocity Problem

- We also have an alternate way to express the solution of the velocity problem
- We often write Δt for h . In that notation, the average velocity of a particle at time a over a time interval of length Δt is $\Delta s / \Delta t$, and we have

The instantaneous velocity of a particle at time a is

$$v(a) = \lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = s'(a)$$

- We also have the concept of average **speed** and **instantaneous speed** which are given by the absolute values of the velocities: $|\Delta s / \Delta t|$ and $|s'(a)|$ respectively.

Rates of Change

- In general, for any process represented by a function $y = f(x)$, we can talk about the average rate of change and the instantaneous rate of change,

$$\frac{\Delta y}{\Delta x} \quad \text{and} \quad \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$$

respectively.

Rates of Change

- In general, for any process represented by a function $y = f(x)$, we can talk about the average rate of change and the instantaneous rate of change,

$$\frac{\Delta y}{\Delta x} \quad \text{and} \quad \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$$

respectively.

- Remember that both those concepts depend on our base number a , although a doesn't appear in the above formulas.

Rates of Change

- In general, for any process represented by a function $y = f(x)$, we can talk about the average rate of change and the instantaneous rate of change,

$$\frac{\Delta y}{\Delta x} \quad \text{and} \quad \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$$

respectively.

- Remember that both those concepts depend on our base number a , although a doesn't appear in the above formulas.
- The concept of instantaneous rate of change is more useful than average rate of change; the former just depends on a while the latter depends on a and Δx .

Examples

1. Let $f(x) = x^3 + 5x + 4$. Find $f'(2)$ from first principles (that is, from the definition of the derivative).
2. Let $g(x) = \frac{4-x}{3+x}$. Find the equation of the tangent line to the curve $y = g(x)$ at the point $P(1, g(1))$ on the curve.
3. If a ball is thrown upward starting at the ground with a velocity of 40 ft/s, its height (in feet) after t seconds is (approximately) given by $s(t) = 40t - 16t^2$.
 - 3.1 Find its (instantaneous) velocity after 1 second.
 - 3.2 Find its velocity at a seconds.
 - 3.3 When will the ball hit the ground?
 - 3.4 What will its velocity be at that time?

Solution to Example 3

- 1 Try this on your own; the answer is similar to that of the previous two questions. The answer will follow from part 2 with $a = 1$, so this part is actually redundant.
- 2 This is an example of calculating a derivative from first principles. We need to calculate the limit

$$v'(a) = \lim_{h \rightarrow 0} \frac{s(a+h) - s(a)}{h}.$$

It is best to split the calculation into parts.

$$s(a) = 40a - 16a^2$$

$$\begin{aligned} s(a+h) &= 40(a+h) - 16(a+h)^2 = 40a + 40h - 16(a^2 + 2ah + h^2) \\ &= 40a + 40h - 16a^2 - 32ah - 16h^2 \end{aligned}$$

Solution to Example 3, continued

2 Subtracting the previous two gives

$$\begin{aligned}s(a+h) - s(a) &= 40a + 40h - 16a^2 - 32ah - 16h^2 - (40a - 16a^2) \\&= 40a + 40h - 16a^2 - 32ah - 16h^2 - 40a + 16a^2 \\&= 40h - 32ah - 16h^2\end{aligned}$$

Note that all the terms without an h have cancelled from the above expression, which is a useful check. Now we can take the limit:

$$\begin{aligned}s'(a) &= \lim_{h \rightarrow 0} \frac{s(a+h) - s(a)}{h} = \lim_{h \rightarrow 0} \frac{40h - 32ah - 16h^2}{h} \\&= \lim_{h \rightarrow 0} (40 - 32a - 16h) = 40 - 32a\end{aligned}$$

Solution to Example 3, continued

- 2 In summary, the instantaneous velocity at a seconds is $40 - 32a$ ft/s. The answer to question 1 is $s'(1) = 40 - 32(1) = 8$ ft/s.
- 3 The ball is at ground level when $s = 0$, so we must have $0 = s(t) = 40t - 16t^2$. Solving for t we get $t = 0$ (rejected in this case; why?) and $40 - 16t = 0$ which implies $t = 2.5$.
- 4 The instantaneous velocity at 2.5 seconds is $s'(2.5) = 40 - 32(2.5) = -40$ ft/s, which agrees with what we would expect from conservation of energy or some other physical reasoning: it hits the ground with the same speed with which it left the ground, but in the opposite direction.

Exercises

Now you should work on Problem Set 2.1. After you have finished it, you should try the following additional exercises from Section 2.1:

2.1 C-level: 1–4, 5–8, 9–10, 15–16, 17–18, 20–23, 27–30, 31–36, 37–42, 43–44, 47–50;

B-level: 11–12, 13–14, 19, 24–26, 45–46, 51–52, 53–56, 57–58, 61;

A-level: 59–60

You should skip example 7 of this section, and don't do example 6 until you understand everything else.