

MATH 110 Lecture 2.7

Rates of Change in the Natural and Social Sciences

Edward Doolittle

Tuesday, February 24, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Rates of Change in the Natural and Social Sciences

Physics

Chemistry

Biology

Economics

Examples and Exercises

Rates of Change in the Natural and Social Sciences

Rates of Change

- Recall the concept of average rate of change.

Rates of Change

- Recall the concept of average rate of change.
- If a variable y depends on a variable x , then as x goes from x_1 to x_2 , y will go from y_1 to y_2 .

Rates of Change

- Recall the concept of average rate of change.
- If a variable y depends on a variable x , then as x goes from x_1 to x_2 , y will go from y_1 to y_2 .
- The change in x is $\Delta x = x_2 - x_1$. The corresponding change in y is $\Delta y = y_2 - y_1$.

Rates of Change

- Recall the concept of average rate of change.
- If a variable y depends on a variable x , then as x goes from x_1 to x_2 , y will go from y_1 to y_2 .
- The change in x is $\Delta x = x_2 - x_1$. The corresponding change in y is $\Delta y = y_2 - y_1$.
- The average rate of change is $\frac{\Delta y}{\Delta x}$.

Rates of Change

- Recall the concept of average rate of change.
- If a variable y depends on a variable x , then as x goes from x_1 to x_2 , y will go from y_1 to y_2 .
- The change in x is $\Delta x = x_2 - x_1$. The corresponding change in y is $\Delta y = y_2 - y_1$.
- The average rate of change is $\frac{\Delta y}{\Delta x}$.
- However, it turns out that in many applications an easier concept to work with is the instantaneous rate of change $\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$.

Rates of Change

- Recall the concept of average rate of change.
- If a variable y depends on a variable x , then as x goes from x_1 to x_2 , y will go from y_1 to y_2 .
- The change in x is $\Delta x = x_2 - x_1$. The corresponding change in y is $\Delta y = y_2 - y_1$.
- The average rate of change is $\frac{\Delta y}{\Delta x}$.
- However, it turns out that in many applications an easier concept to work with is the instantaneous rate of change $\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$.
- We will explore applications of these ideas in physics, chemistry, biology, and (micro-)economics.

Rates of Change in Physics

- The derivative concept is common in many physical applications.

Rates of Change in Physics

- The derivative concept is common in many physical applications.
- There are often different names associated with derivatives of different quantities in physics.

Rates of Change in Physics

- The derivative concept is common in many physical applications.
- There are often different names associated with derivatives of different quantities in physics.
- For example, the derivative of position with respect to time is velocity; the derivative of velocity with respect to time is acceleration.

Rates of Change in Physics

- The derivative concept is common in many physical applications.
- There are often different names associated with derivatives of different quantities in physics.
- For example, the derivative of position with respect to time is velocity; the derivative of velocity with respect to time is acceleration.
- The derivative of charge with respect to time is called **current**.

Rates of Change in Physics

- The derivative concept is common in many physical applications.
- There are often different names associated with derivatives of different quantities in physics.
- For example, the derivative of position with respect to time is velocity; the derivative of velocity with respect to time is acceleration.
- The derivative of charge with respect to time is called **current**.
- The derivative of charge with respect to position is **charge gradient**.

Rates of Change in Physics

- The derivative concept is common in many physical applications.
- There are often different names associated with derivatives of different quantities in physics.
- For example, the derivative of position with respect to time is velocity; the derivative of velocity with respect to time is acceleration.
- The derivative of charge with respect to time is called **current**.
- The derivative of charge with respect to position is **charge gradient**.
- The derivative of work with respect to time is **power**.

Rates of Change in Physics

- The derivative concept is common in many physical applications.
- There are often different names associated with derivatives of different quantities in physics.
- For example, the derivative of position with respect to time is velocity; the derivative of velocity with respect to time is acceleration.
- The derivative of charge with respect to time is called **current**.
- The derivative of charge with respect to position is **charge gradient**.
- The derivative of work with respect to time is **power**.
- The derivative of the amount of radioactive substance is called item the **rate of decay**.

Velocity and Acceleration

- We have already encountered a common application of the rate-of-change concept to physics: velocity and acceleration.

Velocity and Acceleration

- We have already encountered a common application of the rate-of-change concept to physics: velocity and acceleration.
- Time is the independent variable, usually represented by t . Position along some line is the dependent variable, usually represented by $s(t)$ or $x(t)$.

Velocity and Acceleration

- We have already encountered a common application of the rate-of-change concept to physics: velocity and acceleration.
- Time is the independent variable, usually represented by t . Position along some line is the dependent variable, usually represented by $s(t)$ or $x(t)$.
- Velocity means instantaneous velocity, the derivative of the position function. Acceleration means instantaneous acceleration, the derivative of the velocity function.

Velocity and Acceleration

- We have already encountered a common application of the rate-of-change concept to physics: velocity and acceleration.
- Time is the independent variable, usually represented by t . Position along some line is the dependent variable, usually represented by $s(t)$ or $x(t)$.
- Velocity means instantaneous velocity, the derivative of the position function. Acceleration means instantaneous acceleration, the derivative of the velocity function.
- We can interpret the velocity and make conclusions about the motion. When the velocity is positive, the position is increasing; when the velocity is negative, the position is decreasing, and when the velocity is zero, the position is stationary.

Velocity and Acceleration

- We have already encountered a common application of the rate-of-change concept to physics: velocity and acceleration.
- Time is the independent variable, usually represented by t . Position along some line is the dependent variable, usually represented by $s(t)$ or $x(t)$.
- Velocity means instantaneous velocity, the derivative of the position function. Acceleration means instantaneous acceleration, the derivative of the velocity function.
- We can interpret the velocity and make conclusions about the motion. When the velocity is positive, the position is increasing; when the velocity is negative, the position is decreasing, and when the velocity is zero, the position is stationary.
- We can make similar interpretations for the acceleration.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.
- The velocity of the particle at time t is $s'(t) = 3t^2 - 24t + 36$.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.
- The velocity of the particle at time t is $s'(t) = 3t^2 - 24t + 36$.
- The acceleration of the particle at time t is $s''(t) = 6t - 24$.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.
- The velocity of the particle at time t is $s'(t) = 3t^2 - 24t + 36$.
- The acceleration of the particle at time t is $s''(t) = 6t - 24$.
- The particle is at rest when $s'(t) = 0$, i.e., when $3t^2 - 24t + 36 = 0$.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.
- The velocity of the particle at time t is $s'(t) = 3t^2 - 24t + 36$.
- The acceleration of the particle at time t is $s''(t) = 6t - 24$.
- The particle is at rest when $s'(t) = 0$, i.e., when $3t^2 - 24t + 36 = 0$. Solving the quadratic equation we have $t^2 - 8t + 12 = 0$ which implies $t = 2$ and $t = 6$.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.
- The velocity of the particle at time t is $s'(t) = 3t^2 - 24t + 36$.
- The acceleration of the particle at time t is $s''(t) = 6t - 24$.
- The particle is at rest when $s'(t) = 0$, i.e., when $3t^2 - 24t + 36 = 0$. Solving the quadratic equation we have $t^2 - 8t + 12 = 0$ which implies $t = 2$ and $t = 6$.
- The particle is moving in the positive direction when $s'(t) > 0$, i.e., when $3t^2 - 24t + 36 > 0$.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.
- The velocity of the particle at time t is $s'(t) = 3t^2 - 24t + 36$.
- The acceleration of the particle at time t is $s''(t) = 6t - 24$.
- The particle is at rest when $s'(t) = 0$, i.e., when $3t^2 - 24t + 36 = 0$. Solving the quadratic equation we have $t^2 - 8t + 12 = 0$ which implies $t = 2$ and $t = 6$.
- The particle is moving in the positive direction when $s'(t) > 0$, i.e., when $3t^2 - 24t + 36 > 0$. Solving that inequality, we have $3(t - 2)(t - 6) > 0$, which is the case when both factors have the same sign, i.e., when $t < 2$ or $t > 6$.

Velocity and Acceleration

- Suppose, for example, a particle moves on a line according to a law of motion given by $s(t) = t^3 - 12t^2 + 36t$.
- The velocity of the particle at time t is $s'(t) = 3t^2 - 24t + 36$.
- The acceleration of the particle at time t is $s''(t) = 6t - 24$.
- The particle is at rest when $s'(t) = 0$, i.e., when $3t^2 - 24t + 36 = 0$. Solving the quadratic equation we have $t^2 - 8t + 12 = 0$ which implies $t = 2$ and $t = 6$.
- The particle is moving in the positive direction when $s'(t) > 0$, i.e., when $3t^2 - 24t + 36 > 0$. Solving that inequality, we have $3(t - 2)(t - 6) > 0$, which is the case when both factors have the same sign, i.e., when $t < 2$ or $t > 6$.
- Similarly, we can figure out when the particle is speeding up by solving the inequality $s''(t) > 0$.

Linear Density

- In the previous example, we differentiated with respect to time to get a *rate*.

Linear Density

- In the previous example, we differentiated with respect to time to get a *rate*.
- We can also differentiate with respect to a space variable.

Linear Density

- In the previous example, we differentiated with respect to time to get a *rate*.
- We can also differentiate with respect to a space variable.
- Consider a metal rod manufactured in such a way that the cumulative mass at a given location x on the rod is $m(x) = \sqrt{x}$, where m is measured in kg and x is measured in meters from the end of the rod.

Linear Density

- In the previous example, we differentiated with respect to time to get a *rate*.
- We can also differentiate with respect to a space variable.
- Consider a metal rod manufactured in such a way that the cumulative mass at a given location x on the rod is $m(x) = \sqrt{x}$, where m is measured in kg and x is measured in meters from the end of the rod.
- The mass of material between locations x_1 and x_2 is $\Delta m = m(x_2) - m(x_1)$.

Linear Density

- In the previous example, we differentiated with respect to time to get a *rate*.
- We can also differentiate with respect to a space variable.
- Consider a metal rod manufactured in such a way that the cumulative mass at a given location x on the rod is $m(x) = \sqrt{x}$, where m is measured in kg and x is measured in meters from the end of the rod.
- The mass of material between locations x_1 and x_2 is $\Delta m = m(x_2) - m(x_1)$.
- The density of material between locations x_1 and x_2 is $[m(x_2) - m(x_1)] / (x_2 - x_1) = \Delta m / \Delta x$.

Linear Density

- In the previous example, we differentiated with respect to time to get a *rate*.
- We can also differentiate with respect to a space variable.
- Consider a metal rod manufactured in such a way that the cumulative mass at a given location x on the rod is $m(x) = \sqrt{x}$, where m is measured in kg and x is measured in meters from the end of the rod.
- The mass of material between locations x_1 and x_2 is $\Delta m = m(x_2) - m(x_1)$.
- The density of material between locations x_1 and x_2 is $[m(x_2) - m(x_1)] / (x_2 - x_1) = \Delta m / \Delta x$.
- The *linear density* of material is the limit as $\Delta x \rightarrow 0$, namely dm/dx .

Linear Density

- In the previous example, we differentiated with respect to time to get a *rate*.
- We can also differentiate with respect to a space variable.
- Consider a metal rod manufactured in such a way that the cumulative mass at a given location x on the rod is $m(x) = \sqrt{x}$, where m is measured in kg and x is measured in meters from the end of the rod.
- The mass of material between locations x_1 and x_2 is $\Delta m = m(x_2) - m(x_1)$.
- The density of material between locations x_1 and x_2 is $[m(x_2) - m(x_1)] / (x_2 - x_1) = \Delta m / \Delta x$.
- The *linear density* of material is the limit as $\Delta x \rightarrow 0$, namely dm/dx .
- In the above example, the linear density is $d\sqrt{x}/dx = 1/(2\sqrt{x})$.

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.
- Chemists are interested in the rate of change of concentration.

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.
- Chemists are interested in the rate of change of concentration.
- Mass conservation tells us that

$$\frac{d[\text{NaCl}]}{dt}$$

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.
- Chemists are interested in the rate of change of concentration.
- Mass conservation tells us that

$$\frac{d[\text{NaCl}]}{dt} = \frac{d[\text{H}_2\text{O}]}{dt}$$

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.
- Chemists are interested in the rate of change of concentration.
- Mass conservation tells us that

$$\frac{d[\text{NaCl}]}{dt} = \frac{d[\text{H}_2\text{O}]}{dt} = -\frac{d[\text{NaOH}]}{dt}$$

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.
- Chemists are interested in the rate of change of concentration.
- Mass conservation tells us that

$$\frac{d[\text{NaCl}]}{dt} = \frac{d[\text{H}_2\text{O}]}{dt} = -\frac{d[\text{NaOH}]}{dt} = -\frac{d[\text{HCl}]}{dt}$$

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.
- Chemists are interested in the rate of change of concentration.
- Mass conservation tells us that

$$\frac{d[\text{NaCl}]}{dt} = \frac{d[\text{H}_2\text{O}]}{dt} = -\frac{d[\text{NaOH}]}{dt} = -\frac{d[\text{HCl}]}{dt}$$

- That derivative is called the *rate of the reaction*.

The Rate of a Reaction

- Consider a chemical reaction like $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$.
- NaOH and HCl are *reactants*, and NaCl and water H_2O are *products*.
- We measure the amount of NaCl , say, by its *concentration* in moles per liter, represented by $[\text{NaCl}]$.
- Chemists are interested in the rate of change of concentration.
- Mass conservation tells us that

$$\frac{d[\text{NaCl}]}{dt} = \frac{d[\text{H}_2\text{O}]}{dt} = -\frac{d[\text{NaOH}]}{dt} = -\frac{d[\text{HCl}]}{dt}$$

- That derivative is called the *rate of the reaction*.
- In a reaction like $2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}$, we write

$$\frac{1}{2} \frac{d[\text{H}_2\text{O}]}{dt} = -\frac{d[\text{O}_2]}{dt} = -\frac{1}{2} \frac{d[\text{H}_2]}{dt}$$

Population Dynamics

- A standard application of calculus to biology is in the study of population changes over time, i.e., population dynamics.

Population Dynamics

- A standard application of calculus to biology is in the study of population changes over time, i.e., population dynamics.
- We measure the size of a population by counting individuals, so the number is really *discrete*. However, it is convenient to use a continuous variable for the size of large populations.

Population Dynamics

- A standard application of calculus to biology is in the study of population changes over time, i.e., population dynamics.
- We measure the size of a population by counting individuals, so the number is really *discrete*. However, it is convenient to use a continuous variable for the size of large populations.
- The simplest model of population growth is exponential, $P(t) = P_0 2^t$. You will learn about exponential functions in MATH 111.

Population Dynamics

- A standard application of calculus to biology is in the study of population changes over time, i.e., population dynamics.
- We measure the size of a population by counting individuals, so the number is really *discrete*. However, it is convenient to use a continuous variable for the size of large populations.
- The simplest model of population growth is exponential, $P(t) = P_0 2^t$. You will learn about exponential functions in MATH 111.
- Typically, we have some relationship between dP/dt and P , e.g.,

$$\frac{dP}{dt} = r_0 \left(1 - \frac{P(t)}{P_c} \right) P(t) - \beta P(t)$$

and we have to determine something about $P(t)$, e.g., when is $P(t)$ constant.

Marginal Cost

- Suppose we need to produce a widget.

Marginal Cost

- Suppose we need to produce a widget.
- Producing one widget is very expensive because we have to retool our factory.

Marginal Cost

- Suppose we need to produce a widget.
- Producing one widget is very expensive because we have to retool our factory.
- However, producing **one more** widget after producing 100 widgets is probably much less expensive.

Marginal Cost

- Suppose we need to produce a widget.
- Producing one widget is very expensive because we have to retool our factory.
- However, producing **one more** widget after producing 100 widgets is probably much less expensive.
- The cost of producing one more widget after producing x units is called **the marginal cost at production level x** .

Marginal Cost

- Suppose we need to produce a widget.
- Producing one widget is very expensive because we have to retool our factory.
- However, producing **one more** widget after producing 100 widgets is probably much less expensive.
- The cost of producing one more widget after producing x units is called **the marginal cost at production level x** .
- The formula for marginal cost is $C(x+1) - C(x)$, where x is the production level and $C(x)$ is the cost for an entire production run of x units. For example, the marginal cost of producing one more unit after you have produced 100 units is $C(101) - C(100)$.

Marginal Cost

- Suppose we need to produce a widget.
- Producing one widget is very expensive because we have to retool our factory.
- However, producing **one more** widget after producing 100 widgets is probably much less expensive.
- The cost of producing one more widget after producing x units is called **the marginal cost at production level x** .
- The formula for marginal cost is $C(x+1) - C(x)$, where x is the production level and $C(x)$ is the cost for an entire production run of x units. For example, the marginal cost of producing one more unit after you have produced 100 units is $C(101) - C(100)$.
- The word *marginal* comes from the practice of keeping track of the differences $C(x+1) - C(x)$ in the margin of an account.

Marginal Cost and Derivatives

- Marginal cost can be approximated well using derivatives.

Marginal Cost and Derivatives

- Marginal cost can be approximated well using derivatives.
- Recall that $C'(x) = \lim_{h \rightarrow 0} \frac{C(x+h) - C(x)}{h}$.

Marginal Cost and Derivatives

- Marginal cost can be approximated well using derivatives.
- Recall that $C'(x) = \lim_{h \rightarrow 0} \frac{C(x+h) - C(x)}{h}$.
- Since $h = 1$ is close to 0 in a typical economics application, we have $C'(x) \approx \frac{C(x+1) - C(x)}{1} = C(x+1) - C(x)$.

Marginal Cost and Derivatives

- Marginal cost can be approximated well using derivatives.
- Recall that $C'(x) = \lim_{h \rightarrow 0} \frac{C(x+h) - C(x)}{h}$.
- Since $h = 1$ is close to 0 in a typical economics application, we have $C'(x) \approx \frac{C(x+1) - C(x)}{1} = C(x+1) - C(x)$.
- In other words, the marginal cost at production level x is usually close enough to $C'(x)$ for the latter to be a useful approximation.

Marginal Cost and Derivatives

- Marginal cost can be approximated well using derivatives.
- Recall that $C'(x) = \lim_{h \rightarrow 0} \frac{C(x+h) - C(x)}{h}$.
- Since $h = 1$ is close to 0 in a typical economics application, we have $C'(x) \approx \frac{C(x+1) - C(x)}{1} = C(x+1) - C(x)$.
- In other words, the marginal cost at production level x is usually close enough to $C'(x)$ for the latter to be a useful approximation.
- Suppose, for example, that the cost of producing x cases of cigars is $C(x) = 1000 + 30x - 0.1x^2$. The 1000 is the fixed start-up cost; the 30 in $20x$ is the cost of materials per unit, and the term $-0.1x^2$ is a correction factor for economy of scale.

Marginal Cost and Derivatives

- Marginal cost can be approximated well using derivatives.
- Recall that $C'(x) = \lim_{h \rightarrow 0} \frac{C(x+h) - C(x)}{h}$.
- Since $h = 1$ is close to 0 in a typical economics application, we have $C'(x) \approx \frac{C(x+1) - C(x)}{1} = C(x+1) - C(x)$.
- In other words, the marginal cost at production level x is usually close enough to $C'(x)$ for the latter to be a useful approximation.
- Suppose, for example, that the cost of producing x cases of cigars is $C(x) = 1000 + 30x - 0.1x^2$. The 1000 is the fixed start-up cost; the 30 in $20x$ is the cost of materials per unit, and the term $-0.1x^2$ is a correction factor for economy of scale.
- The marginal cost at production level 50 is approximately $C'(50) = 30 - 0.2(50) = 20$.

Examples and Exercises

Examples

1. A particle moves along a horizontal line so that its coordinate at time t is $x = \sqrt{b^2 + c^2 t^2}$, $t \geq 0$, where b and c are positive constants.
 - 1.1 Find the velocity and acceleration functions.
 - 1.2 Show that the particle always moves in a positive direction.
2. The mass of part of a wire is $x(1 + \sqrt{x})$ kilograms, where x is measured in meters from one end of the wire. Find the linear density of the wire when $x = 4$ m.
3. The cost, in dollars, of producing x units of a certain commodity is $C(x) = 920 + 2x - 0.02x^2 + 0.00007x^3$.
 - 3.1 Find the marginal cost function.
 - 3.2 Find $C'(100)$ and explain its meaning.
 - 3.3 Compare $C'(100)$ with the cost of producing the 101st item.

Now you should work on Problem Set 2.7. After you have finished it, you should try the following additional exercises from Section 2.7:

2.7 C-level: 1–4, 6–10, 17–18, 22, 27–30;

B-level: 11–16, 19–21, 23–24, 25–26;

A-level: 31–36