

# MATH 110 Lecture 2.9

## Linear Approximations and Differentials

---

Edward Doolittle

Not part of the course

Department of Indigenous Knowledge and Science  
First Nations University of Canada

## Linear Approximations and Differentials

Linear Approximations

Differentials

Examples and Exercises

# **Linear Approximations and Differentials**

---

# Linear Approximations

- Recall that the big idea of differential calculus is that the tangent line is the “best linear approximation” to a curve.

# Linear Approximations

- Recall that the big idea of differential calculus is that the tangent line is the “best linear approximation” to a curve.
- The equation of the tangent line in point-slope form is  $y - f(a) = f'(a)(x - a)$ .

# Linear Approximations

- Recall that the big idea of differential calculus is that the tangent line is the “best linear approximation” to a curve.
- The equation of the tangent line in point-slope form is  $y - f(a) = f'(a)(x - a)$ .
- Solving for  $y$ , we have  $y = f(a) + f'(a)(x - a)$ .

# Linear Approximations

- Recall that the big idea of differential calculus is that the tangent line is the “best linear approximation” to a curve.
- The equation of the tangent line in point-slope form is  $y - f(a) = f'(a)(x - a)$ .
- Solving for  $y$ , we have  $y = f(a) + f'(a)(x - a)$ .
- To say that the tangent line is an approximation to the curve means  $f(x) \approx f(a) + f'(a)(x - a)$ .

# Linear Approximations

- Recall that the big idea of differential calculus is that the tangent line is the “best linear approximation” to a curve.
- The equation of the tangent line in point-slope form is  $y - f(a) = f'(a)(x - a)$ .
- Solving for  $y$ , we have  $y = f(a) + f'(a)(x - a)$ .
- To say that the tangent line is an approximation to the curve means  $f(x) \approx f(a) + f'(a)(x - a)$ .
- The function on the right side of the above expressions is called the **linearization** of  $f$  at  $a$  and is sometimes denoted  $L(x)$ .



**Example:**  $f(x) = \sqrt{25 - x^2}$  near 3

- Let's find the best linear approximation to  $f(x) = \sqrt{25 - x^2}$  near  $x = 3$ .

**Example:**  $f(x) = \sqrt{25 - x^2}$  near 3

- Let's find the best linear approximation to  $f(x) = \sqrt{25 - x^2}$  near  $x = 3$ .
- We need to compute  $f(3)$  and  $f'(3)$ .

**Example:**  $f(x) = \sqrt{25 - x^2}$  near 3

- Let's find the best linear approximation to  $f(x) = \sqrt{25 - x^2}$  near  $x = 3$ .
- We need to compute  $f(3)$  and  $f'(3)$ .
- $f(3) = \sqrt{25 - 3^2} = \sqrt{16} = 4$ .

### Example: $f(x) = \sqrt{25 - x^2}$ near 3

- Let's find the best linear approximation to  $f(x) = \sqrt{25 - x^2}$  near  $x = 3$ .
- We need to compute  $f(3)$  and  $f'(3)$ .
- $f(3) = \sqrt{25 - 3^2} = \sqrt{16} = 4$ .
- $f'(x) = \frac{1}{2}(25 - x^2)^{-1/2} \cdot -2x$  by the chain rule.

### Example: $f(x) = \sqrt{25 - x^2}$ near 3

- Let's find the best linear approximation to  $f(x) = \sqrt{25 - x^2}$  near  $x = 3$ .
- We need to compute  $f(3)$  and  $f'(3)$ .
- $f(3) = \sqrt{25 - 3^2} = \sqrt{16} = 4$ .
- $f'(x) = \frac{1}{2}(25 - x^2)^{-1/2} \cdot -2x$  by the chain rule.
- $f'(3) = \frac{1}{2} \frac{1}{4} \cdot -6 = -\frac{3}{4}$ .

### Example: $f(x) = \sqrt{25 - x^2}$ near 3

- Let's find the best linear approximation to  $f(x) = \sqrt{25 - x^2}$  near  $x = 3$ .
- We need to compute  $f(3)$  and  $f'(3)$ .
- $f(3) = \sqrt{25 - 3^2} = \sqrt{16} = 4$ .
- $f'(x) = \frac{1}{2}(25 - x^2)^{-1/2} \cdot -2x$  by the chain rule.
- $f'(3) = \frac{1}{2} \frac{1}{4} \cdot -6 = -\frac{3}{4}$ .
- So the linearization of  $f(x)$  near  $x = 3$  is

$$L(x) = f(3) + f'(3)(x - 3) = 4 - \frac{3}{4}(x - 3)$$

### Example: $f(x) = \sqrt{25 - x^2}$ near 3

- Let's find the best linear approximation to  $f(x) = \sqrt{25 - x^2}$  near  $x = 3$ .
- We need to compute  $f(3)$  and  $f'(3)$ .
- $f(3) = \sqrt{25 - 3^2} = \sqrt{16} = 4$ .
- $f'(x) = \frac{1}{2}(25 - x^2)^{-1/2} \cdot -2x$  by the chain rule.
- $f'(3) = \frac{1}{2} \frac{1}{4} \cdot -6 = -\frac{3}{4}$ .
- So the linearization of  $f(x)$  near  $x = 3$  is

$$L(x) = f(3) + f'(3)(x - 3) = 4 - \frac{3}{4}(x - 3)$$

- So we can say  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$ .

## The Error in a Linear Approximation

- The approximation  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$  is exact at  $x = 3$ .



## The Error in a Linear Approximation

- The approximation  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$  is exact at  $x = 3$ .
- The further from  $x = 3$  we get, the worse the approximation.

## The Error in a Linear Approximation

- The approximation  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$  is exact at  $x = 3$ .
- The further from  $x = 3$  we get, the worse the approximation.
- The error is exactly  $4 - \frac{3}{4}(x - 3) - \sqrt{25 - x^2}$ .

## The Error in a Linear Approximation

- The approximation  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$  is exact at  $x = 3$ .
- The further from  $x = 3$  we get, the worse the approximation.
- The error is exactly  $4 - \frac{3}{4}(x - 3) - \sqrt{25 - x^2}$ .
- If we knew the error exactly we could figure out  $\sqrt{25 - x^2}$  exactly and we wouldn't need an approximation!

# The Error in a Linear Approximation

- The approximation  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$  is exact at  $x = 3$ .
- The further from  $x = 3$  we get, the worse the approximation.
- The error is exactly  $4 - \frac{3}{4}(x - 3) - \sqrt{25 - x^2}$ .
- If we knew the error exactly we could figure out  $\sqrt{25 - x^2}$  exactly and we wouldn't need an approximation!
- So the game now is to estimate the error.

# The Error in a Linear Approximation

- The approximation  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$  is exact at  $x = 3$ .
- The further from  $x = 3$  we get, the worse the approximation.
- The error is exactly  $4 - \frac{3}{4}(x - 3) - \sqrt{25 - x^2}$ .
- If we knew the error exactly we could figure out  $\sqrt{25 - x^2}$  exactly and we wouldn't need an approximation!
- So the game now is to estimate the error.
- You will study the problem of estimating the error when you study Taylor series, beginning in MATH 111.

# The Error in a Linear Approximation

- The approximation  $\sqrt{25 - x^2} \approx 4 - \frac{3}{4}(x - 3)$  is exact at  $x = 3$ .
- The further from  $x = 3$  we get, the worse the approximation.
- The error is exactly  $4 - \frac{3}{4}(x - 3) - \sqrt{25 - x^2}$ .
- If we knew the error exactly we could figure out  $\sqrt{25 - x^2}$  exactly and we wouldn't need an approximation!
- So the game now is to estimate the error.
- You will study the problem of estimating the error when you study Taylor series, beginning in MATH 111.
- For now, we will just make some educated guesses.

## Range of Applicability of a Linear Approximation

- We will actually turn the problem around. We will not ask for the error or a bound on the error.

## Range of Applicability of a Linear Approximation

- We will actually turn the problem around. We will not ask for the error or a bound on the error.
- Instead we will ask, “For what  $x$  values is the approximation  $f(x) \approx f(a) + f'(a)(x - a)$  within an error of 0.1?”



## Range of Applicability of a Linear Approximation

- We will actually turn the problem around. We will not ask for the error or a bound on the error.
- Instead we will ask, “For what  $x$  values is the approximation  $f(x) \approx f(a) + f'(a)(x - a)$  within an error of 0.1?”
- We can translate that statement to an inequality to be solved for  $x$ :

$$(f(a) + f'(a)(x - a)) - 0.1 < f(x) < (f(a) + f'(a)(x - a)) + 0.1$$

## Range of Applicability of a Linear Approximation

- We will actually turn the problem around. We will not ask for the error or a bound on the error.
- Instead we will ask, “For what  $x$  values is the approximation  $f(x) \approx f(a) + f'(a)(x - a)$  within an error of 0.1?”
- We can translate that statement to an inequality to be solved for  $x$ :

$$(f(a) + f'(a)(x - a)) - 0.1 < f(x) < (f(a) + f'(a)(x - a)) + 0.1$$

- For now, the best way for us to solve that inequality is graphically.

# Applications of Linear Approximations

- Let's find the best linear approximations to  $\sin x$  and  $\cos x$  near  $x = 0$ .

# Applications of Linear Approximations

- Let's find the best linear approximations to  $\sin x$  and  $\cos x$  near  $x = 0$ .
- We have  $\sin(0) = 0$ ,  $\sin'(0) = 1$ ,  $\cos(0) = 1$ ,  $\cos'(0) = 0$ .

# Applications of Linear Approximations

- Let's find the best linear approximations to  $\sin x$  and  $\cos x$  near  $x = 0$ .
- We have  $\sin(0) = 0$ ,  $\sin'(0) = 1$ ,  $\cos(0) = 1$ ,  $\cos'(0) = 0$ .
- The best linear approximations are then

$$\sin x \approx \sin(0) + \sin'(0)(x - 0) = 0 + 1(x - 0) = x$$

$$\cos x \approx \cos(0) + \cos'(0)(x - 0) = 1 + 0(x - 0) = 1$$

# Applications of Linear Approximations

- Let's find the best linear approximations to  $\sin x$  and  $\cos x$  near  $x = 0$ .
- We have  $\sin(0) = 0$ ,  $\sin'(0) = 1$ ,  $\cos(0) = 1$ ,  $\cos'(0) = 0$ .
- The best linear approximations are then

$$\sin x \approx \sin(0) + \sin'(0)(x - 0) = 0 + 1(x - 0) = x$$

$$\cos x \approx \cos(0) + \cos'(0)(x - 0) = 1 + 0(x - 0) = 1$$

- Those approximations are commonly used in physics.

# Applications of Linear Approximations

- Let's find the best linear approximations to  $\sin x$  and  $\cos x$  near  $x = 0$ .
- We have  $\sin(0) = 0$ ,  $\sin'(0) = 1$ ,  $\cos(0) = 1$ ,  $\cos'(0) = 0$ .
- The best linear approximations are then

$$\sin x \approx \sin(0) + \sin'(0)(x - 0) = 0 + 1(x - 0) = x$$

$$\cos x \approx \cos(0) + \cos'(0)(x - 0) = 1 + 0(x - 0) = 1$$

- Those approximations are commonly used in physics.
- For example, gravitational force on a pendulum is  $-mg \sin x$ , but for small  $x$  the approximation  $-mgx$  is much easier to work with.

# Applications of Linear Approximations

- Let's find the best linear approximations to  $\sin x$  and  $\cos x$  near  $x = 0$ .
- We have  $\sin(0) = 0$ ,  $\sin'(0) = 1$ ,  $\cos(0) = 1$ ,  $\cos'(0) = 0$ .
- The best linear approximations are then

$$\sin x \approx \sin(0) + \sin'(0)(x - 0) = 0 + 1(x - 0) = x$$

$$\cos x \approx \cos(0) + \cos'(0)(x - 0) = 1 + 0(x - 0) = 1$$

- Those approximations are commonly used in physics.
- For example, gravitational force on a pendulum is  $-mg \sin x$ , but for small  $x$  the approximation  $-mgx$  is much easier to work with.
- In optics,  $\sin x \approx x$  and  $\cos x \approx 1$  are used to make lens grinding much easier. Those approximations lead to lenses with spherical surfaces, which are very easy to manufacture and work reasonably well near the center of the field of vision.



- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .

# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .

# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .
- Writing  $\Delta x = x - a$  and  $\Delta y = f(x) - f(a)$  we have  $\Delta y \approx f'(a)\Delta x$ .

# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .
- Writing  $\Delta x = x - a$  and  $\Delta y = f(x) - f(a)$  we have  $\Delta y \approx f'(a)\Delta x$ .
- Because that approximation applies at any value  $a$ , we often write  $\Delta y \approx f'(x)\Delta x$  instead.

# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .
- Writing  $\Delta x = x - a$  and  $\Delta y = f(x) - f(a)$  we have  $\Delta y \approx f'(a)\Delta x$ .
- Because that approximation applies at any value  $a$ , we often write  $\Delta y \approx f'(x)\Delta x$  instead.
- For small values of  $\Delta x$  the approximation is as good as we like.

# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .
- Writing  $\Delta x = x - a$  and  $\Delta y = f(x) - f(a)$  we have  $\Delta y \approx f'(a)\Delta x$ .
- Because that approximation applies at any value  $a$ , we often write  $\Delta y \approx f'(x)\Delta x$  instead.
- For small values of  $\Delta x$  the approximation is as good as we like.
- We imagine that there is a value of  $\Delta x$  so small that the approximation is perfect, or at least good enough that the error is of no consequence.

# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .
- Writing  $\Delta x = x - a$  and  $\Delta y = f(x) - f(a)$  we have  $\Delta y \approx f'(a)\Delta x$ .
- Because that approximation applies at any value  $a$ , we often write  $\Delta y \approx f'(x)\Delta x$  instead.
- For small values of  $\Delta x$  the approximation is as good as we like.
- We imagine that there is a value of  $\Delta x$  so small that the approximation is perfect, or at least good enough that the error is of no consequence.
- We write  $dx$  instead of  $\Delta x$  for such small value of the change in  $x$ ; we also write  $dy$  instead of  $\Delta y$ .

# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .
- Writing  $\Delta x = x - a$  and  $\Delta y = f(x) - f(a)$  we have  $\Delta y \approx f'(a)\Delta x$ .
- Because that approximation applies at any value  $a$ , we often write  $\Delta y \approx f'(x)\Delta x$  instead.
- For small values of  $\Delta x$  the approximation is as good as we like.
- We imagine that there is a value of  $\Delta x$  so small that the approximation is perfect, or at least good enough that the error is of no consequence.
- We write  $dx$  instead of  $\Delta x$  for such small value of the change in  $x$ ; we also write  $dy$  instead of  $\Delta y$ .
- In that case, we write  $dy = f'(x) dx$  exactly.



# Differentials

- We have the approximation  $f(x) \approx f(a) + f'(a)(x - a)$ .
- We can re-write that as  $f(x) - f(a) \approx f'(a)(x - a)$ .
- Writing  $\Delta x = x - a$  and  $\Delta y = f(x) - f(a)$  we have  $\Delta y \approx f'(a)\Delta x$ .
- Because that approximation applies at any value  $a$ , we often write  $\Delta y \approx f'(x)\Delta x$  instead.
- For small values of  $\Delta x$  the approximation is as good as we like.
- We imagine that there is a value of  $\Delta x$  so small that the approximation is perfect, or at least good enough that the error is of no consequence.
- We write  $dx$  instead of  $\Delta x$  for such small value of the change in  $x$ ; we also write  $dy$  instead of  $\Delta y$ .
- In that case, we write  $dy = f'(x) dx$  exactly.
- Instead of “differences”  $\Delta x$  and  $\Delta y$  we talk of “differentials”  $dx$  and  $dy$ .

# Measurement Error

- A common application of differentials is to estimating the consequences of measurement error.

# Measurement Error

- A common application of differentials is to estimating the consequences of measurement error.
- If we have a function  $y = f(x)$  and we get  $x$  from a measurement, there will be a measurement error in  $x$  which we can call  $dx$ .

# Measurement Error

- A common application of differentials is to estimating the consequences of measurement error.
- If we have a function  $y = f(x)$  and we get  $x$  from a measurement, there will be a measurement error in  $x$  which we can call  $dx$ .
- Then the error in the calculated value  $y$  will be approximately  $dy = f'(x) dx$ .

# Measurement Error

- A common application of differentials is to estimating the consequences of measurement error.
- If we have a function  $y = f(x)$  and we get  $x$  from a measurement, there will be a measurement error in  $x$  which we can call  $dx$ .
- Then the error in the calculated value  $y$  will be approximately  $dy = f'(x) dx$ .
- For example, consider a circle. We measure its radius  $r$  and then calculate its area  $A$  using the formula  $A = \pi r^2$ .

# Measurement Error

- A common application of differentials is to estimating the consequences of measurement error.
- If we have a function  $y = f(x)$  and we get  $x$  from a measurement, there will be a measurement error in  $x$  which we can call  $dx$ .
- Then the error in the calculated value  $y$  will be approximately  $dy = f'(x) dx$ .
- For example, consider a circle. We measure its radius  $r$  and then calculate its area  $A$  using the formula  $A = \pi r^2$ .
- We have the error in the calculated value  $A$  is approximately  $dA = A'(r) dr = 2\pi r dr$ .

# Measurement Error

- A common application of differentials is to estimating the consequences of measurement error.
- If we have a function  $y = f(x)$  and we get  $x$  from a measurement, there will be a measurement error in  $x$  which we can call  $dx$ .
- Then the error in the calculated value  $y$  will be approximately  $dy = f'(x) dx$ .
- For example, consider a circle. We measure its radius  $r$  and then calculate its area  $A$  using the formula  $A = \pi r^2$ .
- We have the error in the calculated value  $A$  is approximately  $dA = A'(r) dr = 2\pi r dr$ .
- Suppose we have  $r = 3\text{cm}$  with an error of 0.1 cm or so.

# Measurement Error

- A common application of differentials is to estimating the consequences of measurement error.
- If we have a function  $y = f(x)$  and we get  $x$  from a measurement, there will be a measurement error in  $x$  which we can call  $dx$ .
- Then the error in the calculated value  $y$  will be approximately  $dy = f'(x) dx$ .
- For example, consider a circle. We measure its radius  $r$  and then calculate its area  $A$  using the formula  $A = \pi r^2$ .
- We have the error in the calculated value  $A$  is approximately  $dA = A'(r) dr = 2\pi r dr$ .
- Suppose we have  $r = 3\text{cm}$  with an error of 0.1 cm or so.
- Then we have  $A = \pi r^2 = 9\pi$  with an error of  $dA = 2\pi r dr = 0.6\pi$  or so.



# Examples and Exercises

---

## Examples

1. Find the linear approximation of  $f(x) = \sqrt{1-x}$  near  $a = 0$  and use it to approximate  $\sqrt{0.9}$  and  $\sqrt{1.01}$ .
2. For the function  $y = 2x - x^2$ , compute  $\Delta y$  and  $dy$  for  $x = 2$  and  $dx = \Delta x = -0.4$ . Sketch a diagram illustrating  $dx$ ,  $dy$ , and  $\Delta y$ .
3. The radius of a circular disk is given as 24 cm with a maximum error in measurement of 0.2 cm
  - 3.1 Use differentials to estimate the maximum error in the calculated area of the disk.
  - 3.2 What is the relative error? What is the percentage error?
4. Use differentials to estimate the amount of paint needed to apply a coat of paint 0.05 cm thick to a hemispherical dome with diameter 50 m.

Now you should work on Problem Set 2.9. After you have finished it, you should try the following additional exercises from Section 2.9:

2.9 C-level: 1–30;

B-level: 31–34;

A-level: 35–42