

# MATH 110 Lecture 1.8

## Continuity

---

Edward Doolittle

Thursday, January 22, 2026

Department of Indigenous Knowledge and Science  
First Nations University of Canada

## Continuity (I)

The Definition of Continuity

Continuity Theorems

## Continuity (II)

Composition of Functions

Applications of Continuity




Examples

# Continuity (I)

---




# Examples of Continuous and Discontinuous Functions

Recall the following three graphs from Lecture 2.






# Examples of Continuous and Discontinuous Functions

Recall the following three graphs from Lecture 2. One of these graphs is not like the others.



# Examples of Continuous and Discontinuous Functions


Recall the following three graphs from Lecture 2. One of these graphs is not like the others.



The middle graph is special.


# The Idea of Continuity

- In the middle graph on the previous slide, there were ‘no surprises.’



# The Idea of Continuity


- In the middle graph on the previous slide, there were ‘no surprises.’
- The limits of the function and its values are consistent.






# The Idea of Continuity

- In the middle graph on the previous slide, there were ‘no surprises.’
- The limits of the function and its values are consistent.
- In the other two graphs, the limits of the function are inconsistent with its values.




# The Idea of Continuity

- In the middle graph on the previous slide, there were ‘no surprises.’
- The limits of the function and its values are consistent.
- In the other two graphs, the limits of the function are inconsistent with its values.
- In the first case,  $\lim_{x \rightarrow 1} f(x) = 2$  but  $f(1)$  is undefined.



# The Idea of Continuity

- In the middle graph on the previous slide, there were ‘no surprises.’
- The limits of the function and its values are consistent.
- In the other two graphs, the limits of the function are inconsistent with its values.
- In the first case,  $\lim_{x \rightarrow 1} f(x) = 2$  but  $f(1)$  is undefined.
- In the second case,  $\lim_{x \rightarrow 1} f(x) = 2$  and  $f(1)$  is defined, but  $f(1) \neq 2$ .



# The Definition of Continuity

We can make the special property of the middle graph precise by introducing a definition:

# The Definition of Continuity

We can make the special property of the middle graph precise by introducing a definition:

**Definition:** We say a function  $f$  is **continuous at**  $a$  if the following statement is true:

$$\lim_{x \rightarrow a} f(x) = f(a)$$

# The Definition of Continuity

We can make the special property of the middle graph precise by introducing a definition:

**Definition:** We say a function  $f$  is **continuous at**  $a$  if the following statement is true:

$$\lim_{x \rightarrow a} f(x) = f(a)$$

The definition really says three things:

1.  $\lim_{x \rightarrow a} f(x)$  exists.
2.  $f(a)$  exists.
3. The above two values are equal.

# The Definition of Continuity

We can make the special property of the middle graph precise by introducing a definition:

**Definition:** We say a function  $f$  is **continuous at**  $a$  if the following statement is true:

$$\lim_{x \rightarrow a} f(x) = f(a)$$

The definition really says three things:


1.  $\lim_{x \rightarrow a} f(x)$  exists.
2.  $f(a)$  exists.
3. The above two values are equal.

If any of those conditions is violated, we say  $f$  is **discontinuous at**  $a$ .

# Types of Discontinuities

The discontinuities of a function can be classified into four categories.

1. **Removable discontinuities:** where the limit  $\lim_{x \rightarrow a}$  exists but is not equal to  $f(a)$ .






# Types of Discontinuities

The discontinuities of a function can be classified into four categories.


1. **Removable discontinuities:** where the limit  $\lim_{x \rightarrow a}$  exists but is not equal to  $f(a)$ .



# Types of Discontinuities

The discontinuities of a function can be classified into four categories.


1. **Removable discontinuities:** where the limit  $\lim_{x \rightarrow a}$  exists but is not equal to  $f(a)$ .
2. **Jump discontinuities:** where the one-sided limits exist but are not equal.



# Types of Discontinuities

The discontinuities of a function can be classified into four categories.


1. **Removable discontinuities:** where the limit  $\lim_{x \rightarrow a}$  exists but is not equal to  $f(a)$ .
2. **Jump discontinuities:** where the one-sided limits exist but are not equal.
3. **Infinite discontinuities:** where either or both of the one-sided limits are infinite.



# Types of Discontinuities


The discontinuities of a function can be classified into four categories.

1. **Removable discontinuities:** where the limit  $\lim_{x \rightarrow a}$  exists but is not equal to  $f(a)$ .
2. **Jump discontinuities:** where the one-sided limits exist but are not equal.
3. **Infinite discontinuities:** where either or both of the one-sided limits are infinite.
4. All other cases: where either of the one-sided limits doesn't exist (i.e., the function oscillates rapidly).



# One-sided Continuity

We can define the concept of one-sided continuity using one-sided limits.




# One-sided Continuity

We can define the concept of one-sided continuity using one-sided limits.

**Definition:** We say a function  $f$  is **continuous from the right at  $a$**  if the following statement is true:

$$\lim_{x \rightarrow a^+} f(x) = f(a)$$



# One-sided Continuity


We can define the concept of one-sided continuity using one-sided limits.

**Definition:** We say a function  $f$  is **continuous from the right at  $a$**  if the following statement is true:

$$\lim_{x \rightarrow a^+} f(x) = f(a)$$

**Definition:** We say a function  $f$  is **continuous from the left at  $a$**  if the following statement is true:

$$\lim_{x \rightarrow a^-} f(x) = f(a)$$



## Continuity on an Interval

We are often interested in functions which are nice **everywhere** in some domain. So we introduce a new definition:



# Continuity on an Interval

We are often interested in functions which are nice **everywhere** in some domain. So we introduce a new definition:


**Definition:** A function is called **continuous on an interval** if it is continuous at each point  $a$  in the interval.

# Continuity on an Interval

We are often interested in functions which are nice **everywhere** in some domain. So we introduce a new definition:

**Definition:** A function is called **continuous on an interval** if it is continuous at each point  $a$  in the interval.

E.g., continuous on  $(-\infty, \infty)$ ,




# Continuity on an Interval

We are often interested in functions which are nice **everywhere** in some domain. So we introduce a new definition:

**Definition:** A function is called **continuous on an interval** if it is continuous at each point  $a$  in the interval.

E.g., continuous on  $(-\infty, \infty)$ , continuous on  $(0, \infty)$ .




# Continuity on an Interval

We are often interested in functions which are nice **everywhere** in some domain. So we introduce a new definition:

**Definition:** A function is called **continuous on an interval** if it is continuous at each point  $a$  in the interval.

E.g., continuous on  $(-\infty, \infty)$ , continuous on  $(0, \infty)$ .

If the interval has an endpoint or endpoints, we understand *continuous* at the endpoint to mean *continuous from the right* or *continuous from the left* as appropriate.




# Continuity on an Interval

We are often interested in functions which are nice **everywhere** in some domain. So we introduce a new definition:

**Definition:** A function is called **continuous on an interval** if it is continuous at each point  $a$  in the interval.

E.g., continuous on  $(-\infty, \infty)$ , continuous on  $(0, \infty)$ .

If the interval has an endpoint or endpoints, we understand *continuous* at the endpoint to mean *continuous from the right* or *continuous from the left* as appropriate. E.g., continuous on  $[-1, 1]$ ,




# Continuity on an Interval

We are often interested in functions which are nice **everywhere** in some domain. So we introduce a new definition:

**Definition:** A function is called **continuous on an interval** if it is continuous at each point  $a$  in the interval.

E.g., continuous on  $(-\infty, \infty)$ , continuous on  $(0, \infty)$ .

If the interval has an endpoint or endpoints, we understand *continuous* at the endpoint to mean *continuous from the right* or *continuous from the left* as appropriate. E.g., continuous on  $[-1, 1]$ , not continuous on  $[0, \infty)$ .



# Continuity and Graphs

1. The discontinuities of a function can be easily located by visual inspection of its graph.

# Continuity and Graphs

1. The discontinuities of a function can be easily located by visual inspection of its graph.
2. The discontinuities are where the graph of the function is broken.



# Continuity and Graphs

1. The discontinuities of a function can be easily located by visual inspection of its graph.
2. The discontinuities are where the graph of the function is broken.
3. A function which is continuous on an interval can be drawn without lifting the pencil from the paper.

## Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions.

## Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions. Suppose  $f$  and  $g$  are continuous at  $a$ .

# Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions. Suppose  $f$  and  $g$  are continuous at  $a$ .

1. The sum  $f + g$  is continuous at  $a$ .

# Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions. Suppose  $f$  and  $g$  are continuous at  $a$ .

1. The sum  $f + g$  is continuous at  $a$ .
2. The difference  $f - g$  is continuous at  $a$ .

# Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions. Suppose  $f$  and  $g$  are continuous at  $a$ .

1. The sum  $f + g$  is continuous at  $a$ .
2. The difference  $f - g$  is continuous at  $a$ .
3. The constant multiple  $cf$  is continuous at  $a$ .

# Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions. Suppose  $f$  and  $g$  are continuous at  $a$ .

1. The sum  $f + g$  is continuous at  $a$ .
2. The difference  $f - g$  is continuous at  $a$ .
3. The constant multiple  $cf$  is continuous at  $a$ .
4. The product  $fg$  is continuous at  $a$ .

# Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions. Suppose  $f$  and  $g$  are continuous at  $a$ .

1. The sum  $f + g$  is continuous at  $a$ .
2. The difference  $f - g$  is continuous at  $a$ .
3. The constant multiple  $cf$  is continuous at  $a$ .
4. The product  $fg$  is continuous at  $a$ .
5. The quotient  $f/g$  is continuous at  $a$  **under the condition that**  
 $g(a) \neq 0$ .



# Combining Continuous Functions

The limit laws for combining functions give rules for combining continuous functions. Suppose  $f$  and  $g$  are continuous at  $a$ .

1. The sum  $f + g$  is continuous at  $a$ .
2. The difference  $f - g$  is continuous at  $a$ .
3. The constant multiple  $cf$  is continuous at  $a$ .
4. The product  $fg$  is continuous at  $a$ .
5. The quotient  $f/g$  is continuous at  $a$  **under the condition that**  
 $g(a) \neq 0$ .

In summary, any algebraic combination of continuous functions is continuous, provided we don't divide by zero.

# Continuity of Polynomial and Rational Functions

- Again, from the properties of limits, the constant function  $f(x) = c$  is continuous.

# Continuity of Polynomial and Rational Functions

- Again, from the properties of limits, the constant function  $f(x) = c$  is continuous.
- Also, the ‘identity’ function  $g(x) = x$  is continuous.

# Continuity of Polynomial and Rational Functions

- Again, from the properties of limits, the constant function  $f(x) = c$  is continuous.
- Also, the ‘identity’ function  $g(x) = x$  is continuous.
- From the above results, it follows that polynomials like  $h(x) = x^3 - 5x^2 + 2x - 1$  are continuous on the whole real line.

# Continuity of Polynomial and Rational Functions

- Again, from the properties of limits, the constant function  $f(x) = c$  is continuous.
- Also, the ‘identity’ function  $g(x) = x$  is continuous.
- From the above results, it follows that polynomials like  $h(x) = x^3 - 5x^2 + 2x - 1$  are continuous on the whole real line.
- It also follows that rational functions like  $q(x) = \frac{x^4 - 3x^2 + 5}{x^2 - 1}$  are continuous everywhere except where their denominators are 0.

# Continuity of Polynomial and Rational Functions

- Again, from the properties of limits, the constant function  $f(x) = c$  is continuous.
- Also, the ‘identity’ function  $g(x) = x$  is continuous.
- From the above results, it follows that polynomials like  $h(x) = x^3 - 5x^2 + 2x - 1$  are continuous on the whole real line.
- It also follows that rational functions like  $q(x) = \frac{x^4 - 3x^2 + 5}{x^2 - 1}$  are continuous everywhere except where their denominators are 0.
- We can summarize those statements by saying that polynomials and rational functions are continuous on their domains.

**Example:**  $f(x) = 1/x$

Consider the function  $f(x) = 1/x$ .

**Example:**  $f(x) = 1/x$

Consider the function  $f(x) = 1/x$ .


- The theorem on the previous slide tells us that  $f(x) = 1/x$  is continuous on its domain.



## Example: $f(x) = 1/x$

Consider the function  $f(x) = 1/x$ .


- The theorem on the previous slide tells us that  $f(x) = 1/x$  is continuous on its domain.
- But the graph of  $f(x)$  doesn't look continuous.



## Example: $f(x) = 1/x$

Consider the function  $f(x) = 1/x$ .


- The theorem on the previous slide tells us that  $f(x) = 1/x$  is continuous on its domain.
- But the graph of  $f(x)$  doesn't look continuous.
- The domain of  $f$  is



## Example: $f(x) = 1/x$

Consider the function  $f(x) = 1/x$ .


- The theorem on the previous slide tells us that  $f(x) = 1/x$  is continuous on its domain.
- But the graph of  $f(x)$  doesn't look continuous.
- The domain of  $f$  is  $\{x \in \mathbb{R} : x \neq 0\}$ .



## Example: $f(x) = 1/x$

Consider the function  $f(x) = 1/x$ .


- The theorem on the previous slide tells us that  $f(x) = 1/x$  is continuous on its domain.
- But the graph of  $f(x)$  doesn't look continuous.
- The domain of  $f$  is  $\{x \in \mathbb{R} : x \neq 0\}$ .
- The function is continuous on its domain because there is no point in the domain at which it is discontinuous.



## Example: $f(x) = 1/x$

Consider the function  $f(x) = 1/x$ .

- The theorem on the previous slide tells us that  $f(x) = 1/x$  is continuous on its domain.
- But the graph of  $f(x)$  doesn't look continuous.
- The domain of  $f$  is  $\{x \in \mathbb{R} : x \neq 0\}$ .
- The function is continuous on its domain because there is no point in the domain at which it is discontinuous.
- But the function is not continuous on  $(-\infty, \infty)$ .



# Continuity of Root Functions

Recall the theorem for limits of root functions:

# Continuity of Root Functions

Recall the theorem for limits of root functions:

$$\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$$

for  $a > 0$ .

# Continuity of Root Functions

Recall the theorem for limits of root functions:

$$\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$$

for  $a > 0$ . We also have the result

$$\lim_{x \rightarrow 0^+} \sqrt[n]{x} = 0 = \sqrt[n]{0}$$



# Continuity of Root Functions

Recall the theorem for limits of root functions:

$$\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$$

for  $a > 0$ . We also have the result

$$\lim_{x \rightarrow 0^+} \sqrt[n]{x} = 0 = \sqrt[n]{0}$$

From those results, it follows that any root function is continuous on the interval  $[0, \infty)$ .

# Continuity of Root Functions

Recall the theorem for limits of root functions:

$$\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$$

for  $a > 0$ . We also have the result

$$\lim_{x \rightarrow 0^+} \sqrt[n]{x} = 0 = \sqrt[n]{0}$$

From those results, it follows that any root function is continuous on the interval  $[0, \infty)$ . When  $n$  is odd, we can also take  $n$ th roots of negative numbers, and we get the result

# Continuity of Root Functions

Recall the theorem for limits of root functions:

$$\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$$

for  $a > 0$ . We also have the result

$$\lim_{x \rightarrow 0^+} \sqrt[n]{x} = 0 = \sqrt[n]{0}$$

From those results, it follows that any root function is continuous on the interval  $[0, \infty)$ . When  $n$  is odd, we can also take  $n$ th roots of negative numbers, and we get the result any odd root function is continuous on the interval  $(-\infty, \infty)$ .

# Continuity of Trig Functions

We have the basic trig limits

$$\lim_{x \rightarrow 0} \sin x = 0$$

$$\lim_{x \rightarrow 0} \cos x = 1$$

# Continuity of Trig Functions

We have the basic trig limits



$$\lim_{x \rightarrow 0} \sin x = 0$$

$$\lim_{x \rightarrow 0} \cos x = 1$$

which are obvious from the graphs of sin and cos or from basic trigonometry.

# Continuity of Trig Functions

We have the basic trig limits



$$\lim_{x \rightarrow 0} \sin x = 0$$

$$\lim_{x \rightarrow 0} \cos x = 1$$

which are obvious from the graphs of sin and cos or from basic trigonometry. From those limits and the identities

$$\sin(x+h) = \sin x \cos h + \cos x \sin h$$

$$\cos(x+h) = \cos x \cos h - \sin x \sin h$$

it follows that sin and cos are continuous on the whole real line.

## Use of Continuity to Evaluate Limits

If we know that a function is continuous at  $a$ , it requires no effort to evaluate its limit at  $a$ :

## Use of Continuity to Evaluate Limits

If we know that a function is continuous at  $a$ , it requires no effort to evaluate its limit at  $a$ :

$$\lim_{x \rightarrow a} f(x) = f(a)$$



## Use of Continuity to Evaluate Limits

If we know that a function is continuous at  $a$ , it requires no effort to evaluate its limit at  $a$ :

$$\lim_{x \rightarrow a} f(x) = f(a)$$

For example, from the previous results we know that  $f(x) = \frac{\sin x + 1}{x^2 - 4}$  is continuous at  $x = 3$  (because  $\sin x + 1$  and  $x^2 - 4$  are continuous and  $x^2 - 4 \neq 0$  at  $x = 3$ ), so

## Use of Continuity to Evaluate Limits

If we know that a function is continuous at  $a$ , it requires no effort to evaluate its limit at  $a$ :

$$\lim_{x \rightarrow a} f(x) = f(a)$$

For example, from the previous results we know that  $f(x) = \frac{\sin x + 1}{x^2 - 4}$  is continuous at  $x = 3$  (because  $\sin x + 1$  and  $x^2 - 4$  are continuous and  $x^2 - 4 \neq 0$  at  $x = 3$ ), so

$$\lim_{x \rightarrow 3} \frac{\sin x + 1}{x^2 - 4} = \frac{\sin 3 + 1}{3^2 - 4} \approx 0.2282$$

## Continuity (II)

---

# Composition of Functions

There is one way of combining functions we haven't talked about yet. We can chain together functions by using the output from one function as the input to the next, like this

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

# Composition of Functions

There is one way of combining functions we haven't talked about yet. We can chain together functions by using the output from one function as the input to the next, like this

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

Note that in our poor system of notation,  $g$  acts first but is written after  $f$ .

# Composition of Functions

There is one way of combining functions we haven't talked about yet. We can chain together functions by using the output from one function as the input to the next, like this

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

Note that in our poor system of notation,  $g$  acts first but is written after  $f$ .

E.g., if  $f(x) = \sqrt{x}$  and  $g(x) = x^2 + 1$ , we can construct a new function

$$h(x) = f(g(x)) = \sqrt{g(x)} = \sqrt{x^2 + 1}$$

# Composition of Functions

There is one way of combining functions we haven't talked about yet. We can chain together functions by using the output from one function as the input to the next, like this

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

Note that in our poor system of notation,  $g$  acts first but is written after  $f$ .

E.g., if  $f(x) = \sqrt{x}$  and  $g(x) = x^2 + 1$ , we can construct a new function

$$h(x) = f(g(x)) = \sqrt{g(x)} = \sqrt{x^2 + 1}$$

The resulting function  $h(x)$  is called the **composition of  $f$  and  $g$**  and is sometimes written as  $h = f \circ g$  when we don't want to put the  $x$ s in.

## Composition of Functions: Order Matters

Note that  $f \circ g \neq g \circ f$  in general. In the above case we have


$$(f \circ g)(x) = \sqrt{x^2 + 1} \quad \text{and} \quad (g \circ f)(x) = (f(x))^2 + 1 = (\sqrt{x})^2 + 1$$



# Composition of Functions: Order Matters

Note that  $f \circ g \neq g \circ f$  in general. In the above case we have

$$(f \circ g)(x) = \sqrt{x^2 + 1} \quad \text{and} \quad (g \circ f)(x) = (f(x))^2 + 1 = (\sqrt{x})^2 + 1$$




$$y = \sqrt{x^2 + 1}$$


# Composition of Functions: Order Matters

Note that  $f \circ g \neq g \circ f$  in general. In the above case we have

$$(f \circ g)(x) = \sqrt{x^2 + 1} \quad \text{and} \quad (g \circ f)(x) = (f(x))^2 + 1 = (\sqrt{x})^2 + 1$$



$$y = \sqrt{x^2 + 1}$$



$$y = (\sqrt{x})^2 + 1$$

## Aside: A Rant on Nonlinearity

Also note: we cannot replace  $\sqrt{x^2 + 1}$  with  $\sqrt{x^2} + \sqrt{1}$ !

## Aside: A Rant on Nonlinearity

Also note: we cannot replace  $\sqrt{x^2 + 1}$  with  $\sqrt{x^2} + \sqrt{1}$ ! We say that the square root function is not linear.

## Aside: A Rant on Nonlinearity

Also note: we cannot replace  $\sqrt{x^2 + 1}$  with  $\sqrt{x^2} + \sqrt{1}$ ! We say that the square root function is not linear.

Most of the functions we know are not linear.

$\sqrt{a+b}$  **cannot be replaced with**  $\sqrt{a} + \sqrt{b}$

$(a+b)^2$  **cannot be replaced with**  $a^2 + b^2$

$\sin(a+b)$  **cannot be replaced with**  $\sin(a) + \sin(b)$

$f(a+b)$  **cannot be replaced with**  $f(a) + f(b)$

## Aside: A Rant on Nonlinearity

Also note: we cannot replace  $\sqrt{x^2 + 1}$  with  $\sqrt{x^2} + \sqrt{1}$ ! We say that the square root function is not linear.

Most of the functions we know are not linear.

$\sqrt{a+b}$  **cannot be replaced with**  $\sqrt{a} + \sqrt{b}$

$(a+b)^2$  **cannot be replaced with**  $a^2 + b^2$

$\sin(a+b)$  **cannot be replaced with**  $\sin(a) + \sin(b)$

$f(a+b)$  **cannot be replaced with**  $f(a) + f(b)$

Sometimes the left sides are equal to the right sides **but not in general.**

## Aside: A Rant on Nonlinearity

Also note: we cannot replace  $\sqrt{x^2 + 1}$  with  $\sqrt{x^2} + \sqrt{1}$ ! We say that the square root function is not linear.

Most of the functions we know are not linear.

$\sqrt{a+b}$  **cannot be replaced with**  $\sqrt{a} + \sqrt{b}$

$(a+b)^2$  **cannot be replaced with**  $a^2 + b^2$

$\sin(a+b)$  **cannot be replaced with**  $\sin(a) + \sin(b)$

$f(a+b)$  **cannot be replaced with**  $f(a) + f(b)$

Sometimes the left sides are equal to the right sides **but not in general**.

Functions for which  $f(a+b) = f(a) + f(b)$  are not typical. You study them in linear algebra, but calculus was invented because most useful functions are not like that.


## Another Aside: $x + 1$ vs. $(\sqrt{x})^2 + 1$ vs. $\sqrt{x^2} + 1$

We can't even say  $x + 1$  is the same thing as  $(\sqrt{x})^2 + 1$  if we are being completely correct; the domains are different. Also,  $\sqrt{x^2} + 1$  is different yet again:



## Another Aside: $x + 1$ vs. $(\sqrt{x})^2 + 1$ vs. $\sqrt{x^2} + 1$


We can't even say  $x + 1$  is the same thing as  $(\sqrt{x})^2 + 1$  if we are being completely correct; the domains are different. Also,  $\sqrt{x^2} + 1$  is different yet again:




$$y = x + 1$$

## Another Aside: $x + 1$ vs. $(\sqrt{x})^2 + 1$ vs. $\sqrt{x^2} + 1$

We can't even say  $x + 1$  is the same thing as  $(\sqrt{x})^2 + 1$  if we are being completely correct; the domains are different. Also,  $\sqrt{x^2} + 1$  is different yet again:




$$y = x + 1$$




$$y = (\sqrt{x})^2 + 1$$

## Another Aside: $x + 1$ vs. $(\sqrt{x})^2 + 1$ vs. $\sqrt{x^2} + 1$


We can't even say  $x + 1$  is the same thing as  $(\sqrt{x})^2 + 1$  if we are being completely correct; the domains are different. Also,  $\sqrt{x^2} + 1$  is different yet again:



$$y = x + 1$$



$$y = (\sqrt{x})^2 + 1$$



$$y = \sqrt{x^2} + 1$$

Moral of this story: keep track of not on the formulas describing functions, but also their domains.

# Limits and Compositions of Functions

- Next, we need to figure out how compositions of functions behave under limits.

# Limits and Compositions of Functions

- Next, we need to figure out how compositions of functions behave under limits.
- Unfortunately, “the limit of a composition is the composition of the limits”, but that doesn’t make any sense: limits are numbers and we can’t compose numbers, only functions.

# Limits and Compositions of Functions

- Next, we need to figure out how compositions of functions behave under limits.
- Unfortunately, “the limit of a composition is the composition of the limits”, but that doesn’t make any sense: limits are numbers and we can’t compose numbers, only functions.
- However, we can say “the limit of a function of a function is the function of the limit of the function” under certain conditions.

# Limits and Compositions of Functions

- Next, we need to figure out how compositions of functions behave under limits.
- Unfortunately, “the limit of a composition is the composition of the limits”, but that doesn’t make any sense: limits are numbers and we can’t compose numbers, only functions.
- However, we can say “the limit of a function of a function is the function of the limit of the function” under certain conditions.

**Theorem:** If  $f$  is continuous at  $b$  and  $\lim_{x \rightarrow a} g(x) = b$ , then

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x))$$

# Limits and Compositions of Functions

- Next, we need to figure out how compositions of functions behave under limits.
- Unfortunately, “the limit of a composition is the composition of the limits”, but that doesn’t make any sense: limits are numbers and we can’t compose numbers, only functions.
- However, we can say “the limit of a function of a function is the function of the limit of the function” under certain conditions.

**Theorem:** If  $f$  is continuous at  $b$  and  $\lim_{x \rightarrow a} g(x) = b$ , then

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x))$$

The idea is that we can exchange the  $\lim$  symbol with any continuous function  $f$ .



## Example: Breaking a Function into a Composition

Consider the rather complicated looking function

$$h(x) = \frac{(x-1)^2 \sin(1/(x-1)) + 3}{(x-1)^2 \sin(1/(x-1)) - 3}$$

## Example: Breaking a Function into a Composition

Consider the rather complicated looking function

$$h(x) = \frac{(x-1)^2 \sin(1/(x-1)) + 3}{(x-1)^2 \sin(1/(x-1)) - 3}$$

We try to express  $h$  as a composition by identifying common components.

## Example: Breaking a Function into a Composition

Consider the rather complicated looking function

$$h(x) = \frac{(x-1)^2 \sin(1/(x-1)) + 3}{(x-1)^2 \sin(1/(x-1)) - 3}$$

We try to express  $h$  as a composition by identifying common components.

## Example: Breaking a Function into a Composition

Consider the rather complicated looking function

$$h(x) = \frac{(x-1)^2 \sin(1/(x-1)) + 3}{(x-1)^2 \sin(1/(x-1)) - 3}$$

We try to express  $h$  as a composition by identifying common components. We now have a good candidate for an ‘inner function’:

$$g(x) = (x-1)^2 \sin(1/(x-1))$$

## Example: Breaking a Function into a Composition

Consider the rather complicated looking function

$$h(x) = \frac{(x-1)^2 \sin(1/(x-1)) + 3}{(x-1)^2 \sin(1/(x-1)) - 3}$$

We try to express  $h$  as a composition by identifying common components. We now have a good candidate for an ‘inner function’:

$$g(x) = (x-1)^2 \sin(1/(x-1))$$

Now we can write

$$h(x) = \frac{g(x) + 3}{g(x) - 3}$$

## Example: Breaking a Function into a Composition

Consider the rather complicated looking function

$$h(x) = \frac{(x-1)^2 \sin(1/(x-1)) + 3}{(x-1)^2 \sin(1/(x-1)) - 3}$$

We try to express  $h$  as a composition by identifying common components. We now have a good candidate for an ‘inner function’:

$$g(x) = (x-1)^2 \sin(1/(x-1))$$

Now we can write

$$h(x) = \frac{g(x) + 3}{g(x) - 3}$$

which tells us that  $f(u)$  should be as follows.

$$f(u) = \frac{u + 3}{u - 3}.$$

## Example: Breaking a Function into a Composition

Consider the rather complicated looking function

$$h(x) = \frac{(x-1)^2 \sin(1/(x-1)) + 3}{(x-1)^2 \sin(1/(x-1)) - 3}$$

We try to express  $h$  as a composition by identifying common components. We now have a good candidate for an ‘inner function’:

$$g(x) = (x-1)^2 \sin(1/(x-1))$$

Now we can write

$$h(x) = \frac{g(x) + 3}{g(x) - 3}$$

which tells us that  $f(u)$  should be as follows. Check  $h(x) = f(g(x))$ .

$$f(u) = \frac{u + 3}{u - 3}.$$

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .



## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) \stackrel{?}{=} f(\lim_{x \rightarrow 1} g(x))$$

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) \stackrel{?}{=} f(\lim_{x \rightarrow 1} g(x))$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold.

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) \stackrel{?}{=} f(\lim_{x \rightarrow 1} g(x))$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x-1)^2 \sin(1/(x-1))$$

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) \stackrel{?}{=} f(\lim_{x \rightarrow 1} g(x))$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x-1)^2 \sin(1/(x-1))$$

which we can do using the squeeze theorem.

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) \stackrel{?}{=} f(\lim_{x \rightarrow 1} g(x))$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x - 1)^2 \sin(1/(x - 1)) = 0$$

which we can do using the squeeze theorem.

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) \stackrel{?}{=} f(\lim_{x \rightarrow 1} g(x))$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x - 1)^2 \sin(1/(x - 1)) = 0$$

which we can do using the squeeze theorem. Then  $b = 0$ , and now we need to check whether  $f(x) = (x + 3)/(x - 3)$  is continuous at  $b = 0$ . Since 0 is in the domain of the function, we know that  $f$  is continuous at  $b$ .

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) \stackrel{?}{=} f(\lim_{x \rightarrow 1} g(x))$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x - 1)^2 \sin(1/(x - 1)) = 0$$

which we can do using the squeeze theorem. Then  $b = 0$ , and now we need to check whether  $f(x) = (x + 3)/(x - 3)$  is continuous at  $b = 0$ . Since 0 is in the domain of the function, we know that  $f$  is continuous at  $b$ . Therefore the hypotheses are satisfied and we can complete the calculation.

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) = f(\lim_{x \rightarrow 1} g(x))$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x - 1)^2 \sin(1/(x - 1)) = 0$$

which we can do using the squeeze theorem. Then  $b = 0$ , and now we need to check whether  $f(x) = (x + 3)/(x - 3)$  is continuous at  $b = 0$ . Since 0 is in the domain of the function, we know that  $f$  is continuous at  $b$ . Therefore the hypotheses are satisfied and we can complete the calculation.



## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) = f(\lim_{x \rightarrow 1} g(x)) = f(0)$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x - 1)^2 \sin(1/(x - 1)) = 0$$

which we can do using the squeeze theorem. Then  $b = 0$ , and now we need to check whether  $f(x) = (x + 3)/(x - 3)$  is continuous at  $b = 0$ . Since 0 is in the domain of the function, we know that  $f$  is continuous at  $b$ . Therefore the hypotheses are satisfied and we can complete the calculation.

## Example: The Limit of a Composition

Now that we have broken  $h$  into a composition of functions, we can use our new limit theorem to find the limit  $\lim_{x \rightarrow 1} h(x)$ .

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} f(g(x)) = f(\lim_{x \rightarrow 1} g(x)) = f(0) = \frac{0+3}{0-3} = -1$$

In order to remove the question mark, we need to check whether the hypotheses of the theorem hold. First, we need to evaluate

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} (x-1)^2 \sin(1/(x-1)) = 0$$

which we can do using the squeeze theorem. Then  $b = 0$ , and now we need to check whether  $f(x) = (x+3)/(x-3)$  is continuous at  $b = 0$ . Since 0 is in the domain of the function, we know that  $f$  is continuous at  $b$ . Therefore the hypotheses are satisfied and we can complete the calculation.

## Composition of Continuous Functions

From the previous results and the definition of continuity, we have the following theorem.

**Theorem:** If  $g$  is continuous at  $a$  and if  $f$  is continuous at  $g(a)$ , then the function  $(f \circ g)(x) = f(g(x))$  is continuous at  $a$ .

# Composition of Continuous Functions

From the previous results and the definition of continuity, we have the following theorem.

**Theorem:** If  $g$  is continuous at  $a$  and if  $f$  is continuous at  $g(a)$ , then the function  $(f \circ g)(x) = f(g(x))$  is continuous at  $a$ .

We also have a result about the continuity of a composition of functions on an interval.

**Theorem:** If  $g$  and  $f$  are continuous on their domains, then the composition  $(f \circ g)(x) = f(g(x))$  is continuous on its domain.

# Composition of Continuous Functions

From the previous results and the definition of continuity, we have the following theorem.

**Theorem:** If  $g$  is continuous at  $a$  and if  $f$  is continuous at  $g(a)$ , then the function  $(f \circ g)(x) = f(g(x))$  is continuous at  $a$ .

We also have a result about the continuity of a composition of functions on an interval.

**Theorem:** If  $g$  and  $f$  are continuous on their domains, then the composition  $(f \circ g)(x) = f(g(x))$  is continuous on its domain.

We say that ‘the composition of continuous functions is continuous’.

# Using Continuity to Evaluate Limits of Compositions

Consider the function

$$h(x) = \sqrt{\frac{x^2 - 1}{x + 3}}$$

## Using Continuity to Evaluate Limits of Compositions

Consider the function

$$h(x) = \sqrt{\frac{x^2 - 1}{x + 3}}$$

We can write  $h$  as the composition of two functions

$$f(u) = \sqrt{u} \quad \text{and} \quad g(x) = \frac{x^2 - 1}{x + 3}$$

## Using Continuity to Evaluate Limits of Compositions

Consider the function

$$h(x) = \sqrt{\frac{x^2 - 1}{x + 3}}$$

We can write  $h$  as the composition of two functions

$$f(u) = \sqrt{u} \quad \text{and} \quad g(x) = \frac{x^2 - 1}{x + 3}$$

Both those functions are continuous on their domains, so  $h(x)$  is continuous on its domain.



## Using Continuity to Evaluate Limits of Compositions

Consider the function

$$h(x) = \sqrt{\frac{x^2 - 1}{x + 3}}$$

We can write  $h$  as the composition of two functions

$$f(u) = \sqrt{u} \quad \text{and} \quad g(x) = \frac{x^2 - 1}{x + 3}$$

Both those functions are continuous on their domains, so  $h(x)$  is continuous on its domain. Therefore limits of  $h$  can be simply evaluated, for example

$$\lim_{x \rightarrow 4} \sqrt{\frac{x^2 - 1}{x + 3}} = \sqrt{\frac{4^2 - 1}{4 + 3}}$$


provided the latter expression makes sense.

## The Intermediate Value Theorem (IVT)

- Another major application of continuity is the **Intermediate Value Theorem**.


# The Intermediate Value Theorem (IVT)

- Another major application of continuity is the **Intermediate Value Theorem**.
- The idea: no matter how I connect the points  $(-1, -1)$  and  $(1, 1)$ , the graph of a continuous function must intersect the  $x$ -axis at some number  $c$ ; at that point  $f(c) = 0$  and we say  $c$  is a root of  $f$ .




# The Intermediate Value Theorem (IVT)

- Another major application of continuity is the **Intermediate Value Theorem**.
- The idea: no matter how I connect the points  $(-1, -1)$  and  $(1, 1)$ , the graph of a continuous function must intersect the  $x$ -axis at some number  $c$ ; at that point  $f(c) = 0$  and we say  $c$  is a root of  $f$ .




# The Intermediate Value Theorem (IVT)

- Another major application of continuity is the **Intermediate Value Theorem**.
- The idea: no matter how I connect the points  $(-1, -1)$  and  $(1, 1)$ , the graph of a continuous function must intersect the  $x$ -axis at some number  $c$ ; at that point  $f(c) = 0$  and we say  $c$  is a root of  $f$ .




# The Intermediate Value Theorem (IVT)

- Another major application of continuity is the **Intermediate Value Theorem**.
- The idea: no matter how I connect the points  $(-1, -1)$  and  $(1, 1)$ , the graph of a continuous function must intersect the  $x$ -axis at some number  $c$ ; at that point  $f(c) = 0$  and we say  $c$  is a root of  $f$ .
- $f$  might have multiple roots, but the IVT guarantees there is at least one.




# The Intermediate Value Theorem (IVT)

- Another major application of continuity is the **Intermediate Value Theorem**.
- The idea: no matter how I connect the points  $(-1, -1)$  and  $(1, 1)$ , the graph of a continuous function must intersect the  $x$ -axis at some number  $c$ ; at that point  $f(c) = 0$  and we say  $c$  is a root of  $f$ .
- $f$  might have multiple roots, but the IVT guarantees there is at least one.
- Functions which are not continuous on  $[-1, 1]$  may have a root,



# The Intermediate Value Theorem (IVT)

- Another major application of continuity is the **Intermediate Value Theorem**.
- The idea: no matter how I connect the points  $(-1, -1)$  and  $(1, 1)$ , the graph of a continuous function must intersect the  $x$ -axis at some number  $c$ ; at that point  $f(c) = 0$  and we say  $c$  is a root of  $f$ .
- $f$  might have multiple roots, but the IVT guarantees there is at least one.
- Functions which are not continuous on  $[-1, 1]$  may have a root, but there is no guarantee.





## Formal Statement of the IVT

- The Intermediate Value Theorem may be stated as follows:

**Theorem:** Suppose that  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number satisfying  $f(a) \leq N \leq f(b)$ . Then there exists a number  $c$  in the interval  $(a, b)$  such that  $f(c) = N$ .

## Formal Statement of the IVT

- The Intermediate Value Theorem may be stated as follows:

**Theorem:** Suppose that  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number satisfying  $f(a) \leq N \leq f(b)$ . Then there exists a number  $c$  in the interval  $(a, b)$  such that  $f(c) = N$ .

- In practice, we usually have  $f(a)$  any negative number,  $f(b)$  any positive number, and  $N = 0$ ;

## Formal Statement of the IVT

- The Intermediate Value Theorem may be stated as follows:

**Theorem:** Suppose that  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number satisfying  $f(a) \leq N \leq f(b)$ . Then there exists a number  $c$  in the interval  $(a, b)$  such that  $f(c) = N$ .

- In practice, we usually have  $f(a)$  any negative number,  $f(b)$  any positive number, and  $N = 0$ ;
- or  $f(a)$  any positive number,  $f(b)$  any negative number, and  $N = 0$ .

# Formal Statement of the IVT

- The Intermediate Value Theorem may be stated as follows:

**Theorem:** Suppose that  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number satisfying  $f(a) \leq N \leq f(b)$ . Then there exists a number  $c$  in the interval  $(a, b)$  such that  $f(c) = N$ .

- In practice, we usually have  $f(a)$  any negative number,  $f(b)$  any positive number, and  $N = 0$ ;
- or  $f(a)$  any positive number,  $f(b)$  any negative number, and  $N = 0$ .
- In simpler words, we can say:

If a continuous function changes sign from  $x = a$  to  $x = b$ , it must have a root between  $a$  and  $b$ .

## Existence of Roots Using the IVT

The intermediate value theorem can be used to establish the existence of roots of equations that we cannot solve exactly.

## Existence of Roots Using the IVT

The intermediate value theorem can be used to establish the existence of roots of equations that we cannot solve exactly. For example, let  $f(x) = x^3 - 3x + 1$  and consider the equation  $f(x) = 0$ , i.e.,

$$x^3 - 3x + 1 = 0$$

## Existence of Roots Using the IVT

The intermediate value theorem can be used to establish the existence of roots of equations that we cannot solve exactly. For example, let  $f(x) = x^3 - 3x + 1$  and consider the equation  $f(x) = 0$ , i.e.,

$$x^3 - 3x + 1 = 0$$

You probably don't know a method for solving this equation exactly. However, we can use the IVT to establish the existence of a solution.

## Existence of Roots Using the IVT

The intermediate value theorem can be used to establish the existence of roots of equations that we cannot solve exactly. For example, let  $f(x) = x^3 - 3x + 1$  and consider the equation  $f(x) = 0$ , i.e.,

$$x^3 - 3x + 1 = 0$$

You probably don't know a method for solving this equation exactly. However, we can use the IVT to establish the existence of a solution. Note that  $f(1) = (1)^3 - 3(1) + 1 = -1$  and  $f(2) = 2^3 - 3(2) + 1 = 1$ . Since  $f$  is continuous and we have

$$f(1) < 0 < f(2)$$

it follows from the IVT that there is some number  $c$  in the interval  $(1, 2)$  such that  $f(c) = 0$ .



## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

Suppose, for example, that we know  $f$  is continuous,  $f(0) < 0$  and  $f(1) > 0$ .

## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

Suppose, for example, that we know  $f$  is continuous,  $f(0) < 0$  and  $f(1) > 0$ . Then the IVT tells us that  $f$  has a root somewhere in the interval  $[0, 1]$ .

## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

Suppose, for example, that we know  $f$  is continuous,  $f(0) < 0$  and  $f(1) > 0$ . Then the IVT tells us that  $f$  has a root somewhere in the interval  $[0, 1]$ . We subdivide the interval at the midway point 0.5. There are three cases:

## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

Suppose, for example, that we know  $f$  is continuous,  $f(0) < 0$  and  $f(1) > 0$ . Then the IVT tells us that  $f$  has a root somewhere in the interval  $[0, 1]$ . We subdivide the interval at the midway point 0.5.

There are three cases:

1.  $f(0.5) = 0$ : we have found the root, and we can stop.

## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

Suppose, for example, that we know  $f$  is continuous,  $f(0) < 0$  and  $f(1) > 0$ . Then the IVT tells us that  $f$  has a root somewhere in the interval  $[0, 1]$ . We subdivide the interval at the midway point 0.5.

There are three cases:

1.  $f(0.5) = 0$ : we have found the root, and we can stop.
2.  $f(0.5) < 0$ : the IVT tells us that there is a root in the interval  $(0.5, 1)$ .

## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

Suppose, for example, that we know  $f$  is continuous,  $f(0) < 0$  and  $f(1) > 0$ . Then the IVT tells us that  $f$  has a root somewhere in the interval  $[0, 1]$ . We subdivide the interval at the midway point 0.5.

There are three cases:

1.  $f(0.5) = 0$ : we have found the root, and we can stop.
2.  $f(0.5) < 0$ : the IVT tells us that there is a root in the interval  $(0.5, 1)$ .
3.  $f(0.5) > 0$ : the IVT tells us that there is a root in the interval  $(0, 0.5)$ .

## Location of Roots Using the IVT

Once we know a root exists in some interval, we can use the IVT to estimate the root by **successive approximations**.

Suppose, for example, that we know  $f$  is continuous,  $f(0) < 0$  and  $f(1) > 0$ . Then the IVT tells us that  $f$  has a root somewhere in the interval  $[0, 1]$ . We subdivide the interval at the midway point 0.5.

There are three cases:

1.  $f(0.5) = 0$ : we have found the root, and we can stop.
2.  $f(0.5) < 0$ : the IVT tells us that there is a root in the interval  $(0.5, 1)$ .
3.  $f(0.5) > 0$ : the IVT tells us that there is a root in the interval  $(0, 0.5)$ .

We can then continue to subdivide the interval to get better and better approximations.



# Examples

1. Consider the function

$$f(x) = \frac{\sin x}{x+1}$$

1.1 Find the domain of  $f$ .

1.2 Show that  $f$  is continuous in its domain.

1.3 Evaluate  $\lim_{x \rightarrow \pi} f(x)$ .

1.4 Evaluate  $\lim_{x \rightarrow -1^+} f(x)$ .

2. Show that the function

$$g(x) = \begin{cases} \sin x & \text{if } -\pi \leq x \leq \pi/4 \\ \cos x & \text{if } \pi/4 < x \leq \pi \end{cases}$$

is continuous on the interval  $[-\pi, \pi]$ .

## More Examples

1. Let

$$f(x) = \begin{cases} 2x - x^2 & \text{if } 0 \leq x \leq 2 \\ 2 - x & \text{if } 2 < x \leq 3 \\ x - 4 & \text{if } 3 < x < 4 \\ \pi & \text{if } 4 \leq x \end{cases}$$

Determine whether  $f$  is continuous, continuous from the left, and/or continuous from the right at each point in its domain, and sketch the graph of  $f$ .

2. Let  $g(x) = \lfloor x \rfloor$ , the greatest integer less than or equal to  $x$ , and let  $h(x) = \lfloor x \rfloor + \lfloor -x \rfloor$ .

2.1 For what values of  $a$  does  $\lim_{x \rightarrow a} g(x)$  exist?

2.2 For what values of  $a$  is  $g(x)$  continuous at  $a$ ?

2.3 For what values of  $a$  does  $\lim_{x \rightarrow a} h(x)$  exist?

2.4 For what values of  $a$  is  $h(x)$  continuous at  $a$ ?

## Yet More Examples

1. 1.1 Find the domain of  $f(x) = \frac{\sqrt{x^2 - 9}}{x^2 - 2}$ .  
1.2 Show that  $f$  is continuous on its domain.  
1.3 Find  $\lim_{x \rightarrow 5} f(x)$ .
2. Use the Intermediate Value Theorem to show that the equation  $2 \sin x = 3 - 2x$  has a solution on the interval  $(0, 1)$ .
3. 3.1 Show that there is a root of the equation  $2x^3 + x^2 + 2 = 0$  in the interval  $(-2, -1)$ .  
3.2 Find the root to an accuracy of two decimal points.

Now you should work on Problem Set 1.8. After you have finished it, you should try the following additional exercises from Section 1.8:

1.8 C-level: 1–2, 3–4, 5–6, 7–8, 9–10, 11–14, 15–16, 17–20, 22, 23–24, 35–38, 39–40, 41–43, 49, 53–56;

B-level: 21, 19, 25–32, 33–34, 44–45, 47, 48, 50–52, 57–58, 59–60, 61–62;

A-level: 46, 63–73