

MATH 110 Lecture 3.4

Limits at Infinity; Horizontal Asymptotes

Edward Doolittle

Thursday, March 12, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Limits at Infinity; Horizontal Asymptotes

Horizontal Asymptotes

Limits at Infinity

Infinite Limits at Infinity

Examples and Exercises

Limits at Infinity; Horizontal Asymptotes

Large Scale Behaviour of a Function

- We have studied how to graph a function by concentrating on its interesting features, which can all be found in some limited domain $[a, b]$.

Large Scale Behaviour of a Function

- We have studied how to graph a function by concentrating on its interesting features, which can all be found in some limited domain $[a, b]$.
- However, sometimes we are more interested in the large scale behaviour of a function.

Large Scale Behaviour of a Function

- We have studied how to graph a function by concentrating on its interesting features, which can all be found in some limited domain $[a, b]$.
- However, sometimes we are more interested in the large scale behaviour of a function.
- Consider $C(t) = \frac{30t}{200+t}$ which models the concentration of salt in a tank of water after t minutes (see 3.4.60).

Large Scale Behaviour of a Function

- We have studied how to graph a function by concentrating on its interesting features, which can all be found in some limited domain $[a, b]$.
- However, sometimes we are more interested in the large scale behaviour of a function.
- Consider $C(t) = \frac{30t}{200+t}$ which models the concentration of salt in a tank of water after t minutes (see 3.4.60).
- We are likely most interested in what happens to $C(t)$ over the long term because the short term behaviour is quickly changing and we expect the concentration to settle down to its long term behaviour after a moderate period of time.

Large Scale Behaviour of a Function

- We have studied how to graph a function by concentrating on its interesting features, which can all be found in some limited domain $[a, b]$.
- However, sometimes we are more interested in the large scale behaviour of a function.
- Consider $C(t) = \frac{30t}{200 + t}$ which models the concentration of salt in a tank of water after t minutes (see 3.4.60).
- We are likely most interested in what happens to $C(t)$ over the long term because the short term behaviour is quickly changing and we expect the concentration to settle down to its long term behaviour after a moderate period of time.
- Knowledge of the long term behaviour (also called *end behaviour*) of a function completes the picture and provides a kind of a frame within which we can get a more accurate graph.

Graph of $C(t) = 30t(200 + t)^{-1}$


- Let's graph $C(t)$.

Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$


Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.




Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.
- C'' undef at -200 .




Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.
- C'' undef at -200 .




Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.
- C'' undef at -200 . Concave up on $(-\infty, -200)$, Concave down on $(-200, \infty)$.




Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.
- C'' undef at -200 . Concave up on $(-\infty, -200)$, Concave down on $(-200, \infty)$.
- Vertical asymptote $x = -200$ useful information.




Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.
- C'' undef at -200 . Concave up on $(-\infty, -200)$, Concave down on $(-200, \infty)$.
- Vertical asymptote $x = -200$ useful information.




Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.
- C'' undef at -200 . Concave up on $(-\infty, -200)$, Concave down on $(-200, \infty)$.
- Vertical asymptote $x = -200$ useful information.



Graph of $C(t) = 30t(200 + t)^{-1}$

- Let's graph $C(t)$.
- $C'(t) = 6000(200 + t)^{-2}$,
 $C''(t) = -12000(200 + t)^{-3}$
- Increasing wherever defined.
- C'' undef at -200 . Concave up on $(-\infty, -200)$, Concave down on $(-200, \infty)$.
- Vertical asymptote $x = -200$ useful information.
- If we could predict the horizontal asymptote $y = 30$, C would be easier to graph.




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

$$\bullet f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$$


Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$;




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$;




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$;




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$; concave down on $(-\infty, -1/\sqrt{3})$ and $(1/\sqrt{3}, \infty)$; Concave up on $(-1/\sqrt{3}, 1/\sqrt{3})$.




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$; concave down on $(-\infty, -1/\sqrt{3})$ and $(1/\sqrt{3}, \infty)$; Concave up on $(-1/\sqrt{3}, 1/\sqrt{3})$.
- Concavity changes at $\pm 1/\sqrt{3}$ so they are inflection points.




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$; concave down on $(-\infty, -1/\sqrt{3})$ and $(1/\sqrt{3}, \infty)$; Concave up on $(-1/\sqrt{3}, 1/\sqrt{3})$.
- Concavity changes at $\pm 1/\sqrt{3}$ so they are inflection points.




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$; concave down on $(-\infty, -1/\sqrt{3})$ and $(1/\sqrt{3}, \infty)$; Concave up on $(-1/\sqrt{3}, 1/\sqrt{3})$.
- Concavity changes at $\pm 1/\sqrt{3}$ so they are inflection points.




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$; concave down on $(-\infty, -1/\sqrt{3})$ and $(1/\sqrt{3}, \infty)$; Concave up on $(-1/\sqrt{3}, 1/\sqrt{3})$.
- Concavity changes at $\pm 1/\sqrt{3}$ so they are inflection points.




Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$; concave down on $(-\infty, -1/\sqrt{3})$ and $(1/\sqrt{3}, \infty)$; Concave up on $(-1/\sqrt{3}, 1/\sqrt{3})$.
- Concavity changes at $\pm 1/\sqrt{3}$ so they are inflection points.



Graph of $f(x) = (x^2 - 1)(x^2 + 1)^{-1}$

- $f'(x) = \frac{4x}{(x^2 + 1)^2}, f''(x) = \frac{-12(x + 1/\sqrt{3})(x - 1/\sqrt{3})}{(x^2 + 1)^3}$
- Critical point at $x = 0$; decreasing for $x < 0$; increasing for $x > 0$; critical point at $x = 0$ is a minimum by first derivative test.
- f'' roots at $\pm 1/\sqrt{3}$; concave down on $(-\infty, -1/\sqrt{3})$ and $(1/\sqrt{3}, \infty)$; Concave up on $(-1/\sqrt{3}, 1/\sqrt{3})$.
- Concavity changes at $\pm 1/\sqrt{3}$ so they are inflection points.
- Note the large-scale “frame” on the graph provided by $y = 1$.



Limits at Infinity and Horizontal Asymptotes


- We can find horizontal asymptotes by calculating limits at infinity.

Limits at Infinity and Horizontal Asymptotes

- We can find horizontal asymptotes by calculating limits at infinity.
- **Definition:** Let f be a function defined on some interval (a, ∞) . Then $\lim_{x \rightarrow \infty} f(x) = L$ means that the values of $f(x)$ can be made as close to L as we like by taking x sufficiently large.



Limits at Infinity and Horizontal Asymptotes

- We can find horizontal asymptotes by calculating limits at infinity.
- **Definition:** Let f be a function defined on some interval (a, ∞) . Then $\lim_{x \rightarrow \infty} f(x) = L$ means that the values of $f(x)$ can be made as close to L as we like by taking x sufficiently large.






Limits at Infinity and Horizontal Asymptotes

- We can find horizontal asymptotes by calculating limits at infinity.
- **Definition:** Let f be a function defined on some interval (a, ∞) . Then $\lim_{x \rightarrow \infty} f(x) = L$ means that the values of $f(x)$ can be made as close to L as we like by taking x sufficiently large.




Limits at Infinity and Horizontal Asymptotes



- We can find horizontal asymptotes by calculating limits at infinity.
- **Definition:** Let f be a function defined on some interval (a, ∞) . Then $\lim_{x \rightarrow \infty} f(x) = L$ means that the values of $f(x)$ can be made as close to L as we like by taking x sufficiently large.






Limits at Infinity and Horizontal Asymptotes, ctd






Limits at Infinity and Horizontal Asymptotes, ctd



Limits at Infinity and Horizontal Asymptotes, ctd




Limits at Infinity and Horizontal Asymptotes, ctd



- **Definition:** Let f be a function defined on some interval $(-\infty, b)$. Then $\lim_{x \rightarrow -\infty} f(x) = L$ means that the values of $f(x)$ can be made as close to L as we like by taking x sufficiently large negative.

Limits at Infinity and Horizontal Asymptotes, ctd



- **Definition:** Let f be a function defined on some interval $(-\infty, b)$. Then $\lim_{x \rightarrow -\infty} f(x) = L$ means that the values of $f(x)$ can be made as close to L as we like by taking x sufficiently large negative.
- **Definition:** The line $y = L$ is called a **horizontal asymptote** of the curve $y = f(x)$ if either $\lim_{x \rightarrow \infty} f(x) = L$ or $\lim_{x \rightarrow -\infty} f(x) = L$.

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.
- Similarly, $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$ for any $r > 0$.

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.
- Similarly, $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- We can use those results to find $\lim_{x \rightarrow \pm\infty} \frac{p(x)}{q(x)}$ for any rational function. The process is to divide through by the highest power of x that appears in the denominator then apply limit laws.

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.
- Similarly, $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- We can use those results to find $\lim_{x \rightarrow \pm\infty} \frac{p(x)}{q(x)}$ for any rational function. The process is to divide through by the highest power of x that appears in the denominator then apply limit laws.
- For example, in $C(x) = 30x/(200 + x)$, the highest power of x in the denominator is x^1 so

$$\lim_{x \rightarrow \infty} C(x) = \lim_{x \rightarrow \infty} \frac{30x}{200 + x}$$

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.
- Similarly, $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- We can use those results to find $\lim_{x \rightarrow \pm\infty} \frac{p(x)}{q(x)}$ for any rational function. The process is to divide through by the highest power of x that appears in the denominator then apply limit laws.
- For example, in $C(x) = 30x/(200 + x)$, the highest power of x in the denominator is x^1 so

$$\lim_{x \rightarrow \infty} C(x) = \lim_{x \rightarrow \infty} \frac{30x}{200 + x} = \lim_{x \rightarrow \infty} \frac{30x \cdot \frac{1}{x}}{(200 + x) \cdot \frac{1}{x}}$$

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.
- Similarly, $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- We can use those results to find $\lim_{x \rightarrow \pm\infty} \frac{p(x)}{q(x)}$ for any rational function. The process is to divide through by the highest power of x that appears in the denominator then apply limit laws.
- For example, in $C(x) = 30x/(200 + x)$, the highest power of x in the denominator is x^1 so

$$\lim_{x \rightarrow \infty} C(x) = \lim_{x \rightarrow \infty} \frac{30x}{200 + x} = \lim_{x \rightarrow \infty} \frac{30x \cdot \frac{1}{x}}{(200 + x) \cdot \frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{30}{\frac{200}{x} + 1}$$

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.
- Similarly, $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- We can use those results to find $\lim_{x \rightarrow \pm\infty} \frac{p(x)}{q(x)}$ for any rational function. The process is to divide through by the highest power of x that appears in the denominator then apply limit laws.
- For example, in $C(x) = 30x/(200 + x)$, the highest power of x in the denominator is x^1 so

$$\begin{aligned}\lim_{x \rightarrow \infty} C(x) &= \lim_{x \rightarrow \infty} \frac{30x}{200 + x} = \lim_{x \rightarrow \infty} \frac{30x \cdot \frac{1}{x}}{(200 + x) \cdot \frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{30}{\frac{200}{x} + 1} \\ &= \frac{\lim_{x \rightarrow \infty} 30}{\lim_{x \rightarrow \infty} \frac{200}{x} + 1}\end{aligned}$$

Limits at Infinity for Rational Functions

- Note that $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- For example, $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$, $\lim_{x \rightarrow \infty} \frac{1}{x^{1/2}} = 0$, etc.
- Similarly, $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$ for any $r > 0$.
- We can use those results to find $\lim_{x \rightarrow \pm\infty} \frac{p(x)}{q(x)}$ for any rational function. The process is to divide through by the highest power of x that appears in the denominator then apply limit laws.
- For example, in $C(x) = 30x/(200 + x)$, the highest power of x in the denominator is x^1 so

$$\begin{aligned}\lim_{x \rightarrow \infty} C(x) &= \lim_{x \rightarrow \infty} \frac{30x}{200 + x} = \lim_{x \rightarrow \infty} \frac{30x \cdot \frac{1}{x}}{(200 + x) \cdot \frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{30}{\frac{200}{x} + 1} \\ &= \frac{\lim_{x \rightarrow \infty} 30}{\lim_{x \rightarrow \infty} \frac{200}{x} + 1} = \frac{30}{0 + 1} = 30\end{aligned}$$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1} / (5x^2 + 3x + 2)$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1} / (5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1} / (5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .
- **Note:** $\sqrt{x^4 + 1} \neq \sqrt{x^4} + \sqrt{1}$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1}/(5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .
- **Note:** $\sqrt{x^4 + 1} \neq \sqrt{x^4} + \sqrt{1}$
- To calculate the limit,

$$\lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1}}{5x^2 + 3x + 2} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \frac{1}{x^2}}{(5x^2 + 3x + 2) \frac{1}{x^2}}$$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1}/(5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .
- **Note:** $\sqrt{x^4 + 1} \neq \sqrt{x^4} + \sqrt{1}$
- To calculate the limit,

$$\lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1}}{5x^2 + 3x + 2} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \frac{1}{x^2}}{(5x^2 + 3x + 2) \frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \sqrt{\frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}}$$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1}/(5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .
- **Note:** $\sqrt{x^4 + 1} \neq \sqrt{x^4} + \sqrt{1}$
- To calculate the limit,

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1}}{5x^2 + 3x + 2} &= \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \frac{1}{x^2}}{(5x^2 + 3x + 2) \frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \sqrt{\frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} \\ &= \lim_{x \rightarrow \infty} \frac{\sqrt{(x^4 + 1) \frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}}\end{aligned}$$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1}/(5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .
- **Note:** $\sqrt{x^4 + 1} \neq \sqrt{x^4} + \sqrt{1}$
- To calculate the limit,

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1}}{5x^2 + 3x + 2} &= \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \frac{1}{x^2}}{(5x^2 + 3x + 2) \frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \sqrt{\frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} \\ &= \lim_{x \rightarrow \infty} \frac{\sqrt{(x^4 + 1) \frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{1 + \frac{1}{x^4}}}{1 + \frac{3}{x} + \frac{2}{x^2}}\end{aligned}$$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1}/(5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .
- **Note:** $\sqrt{x^4 + 1} \neq \sqrt{x^4} + \sqrt{1}$
- To calculate the limit,

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1}}{5x^2 + 3x + 2} &= \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \frac{1}{x^2}}{(5x^2 + 3x + 2) \frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \sqrt{\frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} \\ &= \lim_{x \rightarrow \infty} \frac{\sqrt{(x^4 + 1) \frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{1 + \frac{1}{x^4}}}{5 + \lim_{x \rightarrow \infty} \frac{3}{x} + \lim_{x \rightarrow \infty} \frac{2}{x^2}}\end{aligned}$$

Limits at Infinity for Rational/Root Functions

- The same idea works for some functions with roots.
- Consider the function $f(x) = \sqrt{x^4 + 1}/(5x^2 + 3x + 2)$
- The highest power of x that appears in the denominator is x^2 .
- **Note:** $\sqrt{x^4 + 1} \neq \sqrt{x^4} + \sqrt{1}$
- To calculate the limit,

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1}}{5x^2 + 3x + 2} &= \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \frac{1}{x^2}}{(5x^2 + 3x + 2) \frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^4 + 1} \sqrt{\frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} \\&= \lim_{x \rightarrow \infty} \frac{\sqrt{(x^4 + 1) \frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} = \lim_{x \rightarrow \infty} \frac{\sqrt{1 + \frac{1}{x^4}}}{5 + \frac{3}{x} + \frac{2}{x^2}} = \frac{\sqrt{1 + \lim_{x \rightarrow \infty} \frac{1}{x^4}}}{5 + \lim_{x \rightarrow \infty} \frac{3}{x} + \lim_{x \rightarrow \infty} \frac{2}{x^2}} \\&= \frac{\sqrt{1 + 0}}{5 + 0 + 0} = \frac{1}{5}\end{aligned}$$

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.
- Multiplying by the conjugate radical,

$$f(x) = \frac{\sqrt{4x^2 + 3x} - 2x}{1} \cdot \frac{\sqrt{4x^2 + 3x} + 2x}{\sqrt{4x^2 + 3x} + 2x}$$

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.
- Multiplying by the conjugate radical,

$$f(x) = \frac{\sqrt{4x^2 + 3x} - 2x}{1} \cdot \frac{\sqrt{4x^2 + 3x} + 2x}{\sqrt{4x^2 + 3x} + 2x} = \frac{3x}{\sqrt{4x^2 + 3x} + 2x}$$

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.
- Multiplying by the conjugate radical,

$$f(x) = \frac{\sqrt{4x^2 + 3x} - 2x}{1} \cdot \frac{\sqrt{4x^2 + 3x} + 2x}{\sqrt{4x^2 + 3x} + 2x} = \frac{3x}{\sqrt{4x^2 + 3x} + 2x}$$

- The limit has been reduced to the situation of the previous slide.

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.
- Multiplying by the conjugate radical,

$$f(x) = \frac{\sqrt{4x^2 + 3x} - 2x}{1} \cdot \frac{\sqrt{4x^2 + 3x} + 2x}{\sqrt{4x^2 + 3x} + 2x} = \frac{3x}{\sqrt{4x^2 + 3x} + 2x}$$

- The limit has been reduced to the situation of the previous slide.
- The highest power of x in the denominator is x^1 . (We count the x^2 under the square root sign as x^1 .) We have

$$\lim_{x \rightarrow \infty} \frac{3x}{\sqrt{4x^2 + 3x} + 2x}$$

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.
- Multiplying by the conjugate radical,

$$f(x) = \frac{\sqrt{4x^2 + 3x} - 2x}{1} \cdot \frac{\sqrt{4x^2 + 3x} + 2x}{\sqrt{4x^2 + 3x} + 2x} = \frac{3x}{\sqrt{4x^2 + 3x} + 2x}$$

- The limit has been reduced to the situation of the previous slide.
- The highest power of x in the denominator is x^1 . (We count the x^2 under the square root sign as x^1 .) We have

$$\lim_{x \rightarrow \infty} \frac{3x}{\sqrt{4x^2 + 3x} + 2x} = \lim_{x \rightarrow \infty} \frac{\frac{3x}{x}}{\sqrt{\frac{4x^2}{x^2} + \frac{3x}{x^2}} + \frac{2x}{x}}$$

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.
- Multiplying by the conjugate radical,

$$f(x) = \frac{\sqrt{4x^2 + 3x} - 2x}{1} \cdot \frac{\sqrt{4x^2 + 3x} + 2x}{\sqrt{4x^2 + 3x} + 2x} = \frac{3x}{\sqrt{4x^2 + 3x} + 2x}$$

- The limit has been reduced to the situation of the previous slide.
- The highest power of x in the denominator is x^1 . (We count the x^2 under the square root sign as x^1 .) We have

$$\lim_{x \rightarrow \infty} \frac{3x}{\sqrt{4x^2 + 3x} + 2x} = \lim_{x \rightarrow \infty} \frac{\frac{3x}{x}}{\sqrt{\frac{4x^2}{x^2} + \frac{3x}{x^2} + \frac{2x}{x}}} = \frac{3}{\sqrt{4 + 0} + 2}$$

Limits at Infinity for $\sqrt{ax^2 + b} - cx$

- The usual trick for handling limits of the form $\lim_{x \rightarrow \infty} \sqrt{ax^2 + b} - cx$ is to multiply by the conjugate radical.
- Consider $f(x) = \sqrt{4x^2 + 3x} - 2x$.
- Multiplying by the conjugate radical,

$$f(x) = \frac{\sqrt{4x^2 + 3x} - 2x}{1} \cdot \frac{\sqrt{4x^2 + 3x} + 2x}{\sqrt{4x^2 + 3x} + 2x} = \frac{3x}{\sqrt{4x^2 + 3x} + 2x}$$

- The limit has been reduced to the situation of the previous slide.
- The highest power of x in the denominator is x^1 . (We count the x^2 under the square root sign as x^1 .) We have


$$\lim_{x \rightarrow \infty} \frac{3x}{\sqrt{4x^2 + 3x} + 2x} = \lim_{x \rightarrow \infty} \frac{\frac{3x}{x}}{\sqrt{\frac{4x^2}{x^2} + \frac{3x}{x^2} + \frac{2x}{x}}} = \frac{3}{\sqrt{4 + 0} + 2} = \frac{3}{4}$$

The Squeeze Theorem for Limits at Infinity

- The Squeeze Theorem also applies to limits at infinity.


The Squeeze Theorem for Limits at Infinity

- The Squeeze Theorem also applies to limits at infinity.
- Consider $f(x) = \frac{\sin x}{x}$.




The Squeeze Theorem for Limits at Infinity

- The Squeeze Theorem also applies to limits at infinity.
- Consider $f(x) = \frac{\sin x}{x}$.
- We have $-1 \leq \sin x \leq 1$, so $-1/x \leq (\sin x)/x \leq 1/x$ for x positive.




The Squeeze Theorem for Limits at Infinity

- The Squeeze Theorem also applies to limits at infinity.
- Consider $f(x) = \frac{\sin x}{x}$.
- We have $-1 \leq \sin x \leq 1$, so $-1/x \leq (\sin x)/x \leq 1/x$ for x positive.
- We have $\lim_{x \rightarrow \infty} -\frac{1}{x} = 0$ and $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$.




The Squeeze Theorem for Limits at Infinity

- The Squeeze Theorem also applies to limits at infinity.
- Consider $f(x) = \frac{\sin x}{x}$.
- We have $-1 \leq \sin x \leq 1$, so $-1/x \leq (\sin x)/x \leq 1/x$ for x positive.
- We have $\lim_{x \rightarrow \infty} -\frac{1}{x} = 0$ and $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$.
- Therefore by the Squeeze Theorem,
$$\lim_{x \rightarrow \infty} \frac{\sin x}{x} = 0.$$



The Squeeze Theorem for Limits at Infinity

- The Squeeze Theorem also applies to limits at infinity.
- Consider $f(x) = \frac{\sin x}{x}$.
- We have $-1 \leq \sin x \leq 1$, so $-1/x \leq (\sin x)/x \leq 1/x$ for x positive.
- We have $\lim_{x \rightarrow \infty} -\frac{1}{x} = 0$ and $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$.
- Therefore by the Squeeze Theorem,
$$\lim_{x \rightarrow \infty} \frac{\sin x}{x} = 0.$$
- Similarly, you can show
$$\lim_{x \rightarrow -\infty} \frac{\sin x}{x} = 0. \text{ (Try it!)}$$



Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.

Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.
- We use one of the following notations:

$$\lim_{x \rightarrow \infty} f(x) = \infty$$

Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.
- We use one of the following notations:

$$\lim_{x \rightarrow \infty} f(x) = \infty \qquad \lim_{x \rightarrow \infty} f(x) = -\infty$$

Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.
- We use one of the following notations:

$$\lim_{x \rightarrow \infty} f(x) = \infty \qquad \lim_{x \rightarrow \infty} f(x) = -\infty$$

$$\lim_{x \rightarrow -\infty} f(x) = \infty$$

Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.
- We use one of the following notations:

$$\lim_{x \rightarrow \infty} f(x) = \infty \qquad \lim_{x \rightarrow \infty} f(x) = -\infty$$

$$\lim_{x \rightarrow -\infty} f(x) = \infty \qquad \lim_{x \rightarrow -\infty} f(x) = -\infty$$


Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.
- We use one of the following notations:

$$\lim_{x \rightarrow \infty} f(x) = \infty \qquad \lim_{x \rightarrow \infty} f(x) = -\infty$$

$$\lim_{x \rightarrow -\infty} f(x) = \infty \qquad \lim_{x \rightarrow -\infty} f(x) = -\infty$$

- For example, $\lim_{x \rightarrow \infty} x^3 = \infty$




Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.
- We use one of the following notations:

$$\lim_{x \rightarrow \infty} f(x) = \infty \qquad \lim_{x \rightarrow \infty} f(x) = -\infty$$

$$\lim_{x \rightarrow -\infty} f(x) = \infty \qquad \lim_{x \rightarrow -\infty} f(x) = -\infty$$

- For example, $\lim_{x \rightarrow \infty} x^3 = \infty$
- Also $\lim_{x \rightarrow -\infty} x^3 = -\infty$.




Infinite Limits at Infinity

- Instead of tending to a finite value L , some functions $f(x)$ grow without bound as x grows.
- We use one of the following notations:

$$\lim_{x \rightarrow \infty} f(x) = \infty \qquad \lim_{x \rightarrow \infty} f(x) = -\infty$$

$$\lim_{x \rightarrow -\infty} f(x) = \infty \qquad \lim_{x \rightarrow -\infty} f(x) = -\infty$$

- For example, $\lim_{x \rightarrow \infty} x^3 = \infty$
- Also $\lim_{x \rightarrow -\infty} x^3 = -\infty$.
- Also, $\lim_{x \rightarrow \infty} x^3 - x = \lim_{x \rightarrow \infty} x(x^2 - 1) = \infty$ because both multiplicands tend to ∞ .



Infinite Limits at Infinity for Rational Functions

- If the degree of the numerator of a rational function $f(x)$ exceeds the degree of the denominator, the limit at infinity of $f(x)$ is $\pm\infty$.

Infinite Limits at Infinity for Rational Functions

- If the degree of the numerator of a rational function $f(x)$ exceeds the degree of the denominator, the limit at infinity of $f(x)$ is $\pm\infty$.
- The limit can be calculated by dividing through by the highest power of x in the denominator, as usual.

Infinite Limits at Infinity for Rational Functions

- If the degree of the numerator of a rational function $f(x)$ exceeds the degree of the denominator, the limit at infinity of $f(x)$ is $\pm\infty$.
- The limit can be calculated by dividing through by the highest power of x in the denominator, as usual.
- Consider $\lim_{x \rightarrow \infty} \frac{-3x^2 + 2}{5x - 4}$.

Infinite Limits at Infinity for Rational Functions

- If the degree of the numerator of a rational function $f(x)$ exceeds the degree of the denominator, the limit at infinity of $f(x)$ is $\pm\infty$.
- The limit can be calculated by dividing through by the highest power of x in the denominator, as usual.
- Consider $\lim_{x \rightarrow \infty} \frac{-3x^2 + 2}{5x - 4}$.
- The highest power of x in the denominator is x^1 . We have

$$\lim_{x \rightarrow \infty} \frac{-3x^2 + 2}{5x - 4} = \lim_{x \rightarrow \infty} \frac{(-3x^2 + 2) \cdot \frac{1}{x}}{(5x - 4) \cdot \frac{1}{x}}$$

Infinite Limits at Infinity for Rational Functions

- If the degree of the numerator of a rational function $f(x)$ exceeds the degree of the denominator, the limit at infinity of $f(x)$ is $\pm\infty$.
- The limit can be calculated by dividing through by the highest power of x in the denominator, as usual.
- Consider $\lim_{x \rightarrow \infty} \frac{-3x^2 + 2}{5x - 4}$.
- The highest power of x in the denominator is x^1 . We have

$$\lim_{x \rightarrow \infty} \frac{-3x^2 + 2}{5x - 4} = \lim_{x \rightarrow \infty} \frac{(-3x^2 + 2) \cdot \frac{1}{x}}{(5x - 4) \cdot \frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{-3x + \frac{2}{x}}{5 - \frac{4}{x}}$$

Infinite Limits at Infinity for Rational Functions

- If the degree of the numerator of a rational function $f(x)$ exceeds the degree of the denominator, the limit at infinity of $f(x)$ is $\pm\infty$.
- The limit can be calculated by dividing through by the highest power of x in the denominator, as usual.
- Consider $\lim_{x \rightarrow \infty} \frac{-3x^2 + 2}{5x - 4}$.
- The highest power of x in the denominator is x^1 . We have

$$\lim_{x \rightarrow \infty} \frac{-3x^2 + 2}{5x - 4} = \lim_{x \rightarrow \infty} \frac{(-3x^2 + 2) \cdot \frac{1}{x}}{(5x - 4) \cdot \frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{-3x + \frac{2}{x}}{5 - \frac{4}{x}}$$

- The numerator of the expression tends to $-\infty$, the denominator to the finite value 1, so the result is $-\infty$.

Examples and Exercises

Examples

1. Find the following limits at infinity.

$$1.1 \quad \lim_{x \rightarrow \infty} \frac{3x^4 + x - 5}{6x^4 - 2x^2 + 1}$$

$$1.2 \quad \lim_{x \rightarrow -\infty} \frac{\sqrt{4x^2 + 1}}{3x - 1}$$

$$1.3 \quad \lim_{x \rightarrow -\infty} (x^2 + x^3)$$

$$1.4 \quad \lim_{x \rightarrow -\infty} \left(\sqrt{4x^2 + 3x - 2x} \right)$$

$$1.5 \quad \lim_{x \rightarrow \infty} \frac{\sin^4 x}{\sqrt{x}}$$

2. Sketch a graph of $y = \frac{1-x}{1+x}$ illustrating intervals of increase/decrease, extrema, concavity, inflection points, and horizontal and vertical asymptotes.

3. The same for $y = \frac{\sqrt{2x^2 + 1}}{3x - 5}$.

Now you should work on Problem Set 3.4. After you have finished it, you should try the following additional exercises from Section 3.4:

3.4 C-level: 1–40, 44–46, 48–51, 57–60;

B-level: 41–43, 47, 52–56, 61–64;

A-level: 65–74