

MATH 110 Lecture 2.6

Implicit Differentiation

Edward Doolittle

Thursday, February 12, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Implicit Differentiation

Implicit Functions


Implicit Differentiation

Examples and Exercises

Implicit Differentiation


Implicit and Explicit Functions

- Consider the curve defined by $x^2 + y^2 = 1$.




Implicit and Explicit Functions

- Consider the curve defined by $x^2 + y^2 = 1$.
- That expression can be solved for y to get two different continuous functions.




Implicit and Explicit Functions

- Consider the curve defined by $x^2 + y^2 = 1$.
- That expression can be solved for y to get two different continuous functions.
- $y = \sqrt{1 - x^2}$




Implicit and Explicit Functions

- Consider the curve defined by $x^2 + y^2 = 1$.
- That expression can be solved for y to get two different continuous functions.
- $y = \sqrt{1 - x^2}$
- $y = -\sqrt{1 - x^2}$




Implicit and Explicit Functions

- Consider the curve defined by $x^2 + y^2 = 1$.
- That expression can be solved for y to get two different continuous functions.
- $y = \sqrt{1 - x^2}$
- $y = -\sqrt{1 - x^2}$
- We say that $y = \sqrt{1 - x^2}$ and $y = -\sqrt{1 - x^2}$ are **explicit** functions.




Implicit and Explicit Functions

- Consider the curve defined by $x^2 + y^2 = 1$.
- That expression can be solved for y to get two different continuous functions.
- $y = \sqrt{1 - x^2}$
- $y = -\sqrt{1 - x^2}$
- We say that $y = \sqrt{1 - x^2}$ and $y = -\sqrt{1 - x^2}$ are **explicit** functions.
- The formula $x^2 + y^2 = 1$ is an **implicit** definition of those functions.




Explicit Functions are Hard to Find

- Consider the curve defined by
 $x^3 + y^3 = 6xy$.



Explicit Functions are Hard to Find


- Consider the curve defined by
 $x^3 + y^3 = 6xy$.
- That expression can be solved for y only with considerable difficulty.



Explicit Functions are Hard to Find

- Consider the curve defined by
$$x^3 + y^3 = 6xy.$$
- That expression can be solved for y only with considerable difficulty.
- For example, the simplest explicit function determine by $x^3 + y^3 = 6xy$ is

$$y = \sqrt[3]{-\frac{1}{2}x^3 + \sqrt{\frac{1}{4}x^6 - 8x^3}} + \sqrt[3]{-\frac{1}{2}x^3 - \sqrt{\frac{1}{4}x^6 - 8x^3}}$$



Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.
- We use the process of implicit differentiation.

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.
- We use the process of implicit differentiation.
- We differentiate an expression as if it were a function, where we use the chain rule every time we see a y .

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.
- We use the process of implicit differentiation.
- We differentiate an expression as if it were a function, where we use the chain rule every time we see a y .
- For example, for $x^2 + y^2 = 1$ we have

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.
- We use the process of implicit differentiation.
- We differentiate an expression as if it were a function, where we use the chain rule every time we see a y .
- For example, for $x^2 + y^2 = 1$ we have

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1$$

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.
- We use the process of implicit differentiation.
- We differentiate an expression as if it were a function, where we use the chain rule every time we see a y .
- For example, for $x^2 + y^2 = 1$ we have

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1 \implies \frac{d}{dx}x^2 + \frac{d}{dx}y^2 = 0$$

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.
- We use the process of implicit differentiation.
- We differentiate an expression as if it were a function, where we use the chain rule every time we see a y .
- For example, for $x^2 + y^2 = 1$ we have

$$\begin{aligned}\frac{d}{dx}(x^2 + y^2) &= \frac{d}{dx}1 \implies \frac{d}{dx}x^2 + \frac{d}{dx}y^2 = 0 \\ &\implies 2x + 2y\frac{dy}{dx} = 0\end{aligned}$$

Implicit Differentiation

- Some implicit functions can't be solved explicitly at all.
- But we may need to differentiate them anyway.
- We use the process of implicit differentiation.
- We differentiate an expression as if it were a function, where we use the chain rule every time we see a y .
- For example, for $x^2 + y^2 = 1$ we have

$$\begin{aligned}\frac{d}{dx}(x^2 + y^2) &= \frac{d}{dx}1 \implies \frac{d}{dx}x^2 + \frac{d}{dx}y^2 = 0 \\ &\implies 2x + 2y\frac{dy}{dx} = 0 \\ &\implies \frac{dy}{dx} = -\frac{x}{y}\end{aligned}$$

Comparing Implicit and Explicit Differentiation

- Note that in the result of implicit differentiation we have a y .

Comparing Implicit and Explicit Differentiation

- Note that in the result of implicit differentiation we have a y .
- We usually can't get rid of the y unless the function can be expressed in explicit form $y = f(x)$.

Comparing Implicit and Explicit Differentiation

- Note that in the result of implicit differentiation we have a y .
- We usually can't get rid of the y unless the function can be expressed in explicit form $y = f(x)$.
- The two ways of calculating the derivative should agree. Recall the example $x^2 + y^2 = 1$.

Comparing Implicit and Explicit Differentiation

- Note that in the result of implicit differentiation we have a y .
- We usually can't get rid of the y unless the function can be expressed in explicit form $y = f(x)$.
- The two ways of calculating the derivative should agree. Recall the example $x^2 + y^2 = 1$.
- Differentiating implicitly then substituting $y = \sqrt{1 - x^2}$, we have
$$\frac{dy}{dx} = -\frac{x}{y} = -\frac{x}{\sqrt{1 - x^2}}.$$

Comparing Implicit and Explicit Differentiation

- Note that in the result of implicit differentiation we have a y .
- We usually can't get rid of the y unless the function can be expressed in explicit form $y = f(x)$.
- The two ways of calculating the derivative should agree. Recall the example $x^2 + y^2 = 1$.
- Differentiating implicitly then substituting $y = \sqrt{1 - x^2}$, we have
$$\frac{dy}{dx} = -\frac{x}{y} = -\frac{x}{\sqrt{1 - x^2}}.$$
- On the other hand, starting with $y = \sqrt{1 - x^2}$ we have

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(1 - x^2)^{1/2} \\ &= \frac{1}{2}(1 - x^2)^{-1/2} \frac{d}{dx}(1 - x^2) = -\frac{x}{\sqrt{1 - x^2}}\end{aligned}$$

Comparing Implicit and Explicit Differentiation

- Note that in the result of implicit differentiation we have a y .
- We usually can't get rid of the y unless the function can be expressed in explicit form $y = f(x)$.
- The two ways of calculating the derivative should agree. Recall the example $x^2 + y^2 = 1$.
- Differentiating implicitly then substituting $y = \sqrt{1 - x^2}$, we have $\frac{dy}{dx} = -\frac{x}{y} = -\frac{x}{\sqrt{1 - x^2}}$.
- On the other hand, starting with $y = \sqrt{1 - x^2}$ we have

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(1 - x^2)^{1/2} \\ &= \frac{1}{2}(1 - x^2)^{-1/2} \frac{d}{dx}(1 - x^2) = -\frac{x}{\sqrt{1 - x^2}}\end{aligned}$$

- The two methods agree. (What happens if $y = -\sqrt{1 - x^2}$?)

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.
- The only good option is implicit differentiation:

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.
- The only good option is implicit differentiation:

$$\frac{d}{dx}(x^3 + y^3) = \frac{d}{dx}(6xy)$$

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.
- The only good option is implicit differentiation:

$$\frac{d}{dx}(x^3 + y^3) = \frac{d}{dx}(6xy) \implies \frac{d}{dx}x^3 + \frac{d}{dx}y^3 = 6\left(\frac{d}{dx}x\right)y + 6x\frac{d}{dx}y$$

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.
- The only good option is implicit differentiation:

$$\begin{aligned}\frac{d}{dx}(x^3 + y^3) &= \frac{d}{dx}(6xy) \implies \frac{d}{dx}x^3 + \frac{d}{dx}y^3 = 6\left(\frac{d}{dx}x\right)y + 6x\frac{d}{dx}y \\ &\implies 3x^2 + 3y^2\frac{dy}{dx} = 6y + 6x\frac{dy}{dx}\end{aligned}$$

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.
- The only good option is implicit differentiation:

$$\begin{aligned}\frac{d}{dx}(x^3 + y^3) &= \frac{d}{dx}(6xy) \implies \frac{d}{dx}x^3 + \frac{d}{dx}y^3 = 6\left(\frac{d}{dx}x\right)y + 6x\frac{d}{dx}y \\ &\implies 3x^2 + 3y^2\frac{dy}{dx} = 6y + 6x\frac{dy}{dx} \\ &\implies (3y^2 - 6x)\frac{dy}{dx} = 6y - 3x^2\end{aligned}$$

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.
- The only good option is implicit differentiation:

$$\begin{aligned}\frac{d}{dx}(x^3 + y^3) &= \frac{d}{dx}(6xy) \implies \frac{d}{dx}x^3 + \frac{d}{dx}y^3 = 6\left(\frac{d}{dx}x\right)y + 6x\frac{d}{dx}y \\ &\implies 3x^2 + 3y^2\frac{dy}{dx} = 6y + 6x\frac{dy}{dx} \\ &\implies (3y^2 - 6x)\frac{dy}{dx} = 6y - 3x^2 \\ &\implies \frac{dy}{dx} = \frac{6y - 3x^2}{3y^2 - 6x}\end{aligned}$$

Implicit Differentiation is the Better Option

- In most cases, it is difficult or impossible to put an implicit function in explicit form. Consider $x^3 + y^3 = 6xy$.
- The explicit form for y is too difficult for us to remember.
- The only good option is implicit differentiation:

$$\begin{aligned}\frac{d}{dx}(x^3 + y^3) &= \frac{d}{dx}(6xy) \implies \frac{d}{dx}x^3 + \frac{d}{dx}y^3 = 6\left(\frac{d}{dx}x\right)y + 6x\frac{d}{dx}y \\ &\implies 3x^2 + 3y^2\frac{dy}{dx} = 6y + 6x\frac{dy}{dx} \\ &\implies (3y^2 - 6x)\frac{dy}{dx} = 6y - 3x^2 \\ &\implies \frac{dy}{dx} = \frac{6y - 3x^2}{3y^2 - 6x}\end{aligned}$$

- Note that to complete the calculation I had to solve for dy/dx .

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.
- We have no choice but to use implicit differentiation.

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.
- We have no choice but to use implicit differentiation.

$$\frac{d}{dx} \sin(xy) = \frac{d}{dx} (x^2 + y)$$

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.
- We have no choice but to use implicit differentiation.

$$\frac{d}{dx} \sin(xy) = \frac{d}{dx} (x^2 + y) \implies \cos(xy) \frac{d}{dx} (xy) = \frac{d}{dx} x^2 + \frac{d}{dx} y$$

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.
- We have no choice but to use implicit differentiation.

$$\begin{aligned}\frac{d}{dx} \sin(xy) &= \frac{d}{dx} (x^2 + y) \implies \cos(xy) \frac{d}{dx} (xy) = \frac{d}{dx} x^2 + \frac{d}{dx} y \\ &\implies \cos(xy) (y + xy') = 2x + y'\end{aligned}$$

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.
- We have no choice but to use implicit differentiation.

$$\begin{aligned}\frac{d}{dx} \sin(xy) &= \frac{d}{dx} (x^2 + y) \implies \cos(xy) \frac{d}{dx} (xy) = \frac{d}{dx} x^2 + \frac{d}{dx} y \\ &\implies \cos(xy) (y + xy') = 2x + y' \\ &\implies (x \cos(xy) - 1) y' = 2x - y \cos(xy)\end{aligned}$$

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.
- We have no choice but to use implicit differentiation.

$$\begin{aligned}\frac{d}{dx} \sin(xy) &= \frac{d}{dx} (x^2 + y) \implies \cos(xy) \frac{d}{dx} (xy) = \frac{d}{dx} x^2 + \frac{d}{dx} y \\ &\implies \cos(xy) (y + xy') = 2x + y' \\ &\implies (x \cos(xy) - 1) y' = 2x - y \cos(xy) \\ &\implies y' = \frac{2x - y \cos(xy)}{x \cos(xy) - 1}\end{aligned}$$

When Implicit Differentiation is Necessary

- Sometimes it is impossible to put an implicit function in explicit form. Consider $\sin(xy) = x^2 + y$. I can't solve for y explicitly.
- We have no choice but to use implicit differentiation.

$$\begin{aligned}\frac{d}{dx} \sin(xy) &= \frac{d}{dx} (x^2 + y) \implies \cos(xy) \frac{d}{dx} (xy) = \frac{d}{dx} x^2 + \frac{d}{dx} y \\ &\implies \cos(xy) (y + xy') = 2x + y' \\ &\implies (x \cos(xy) - 1) y' = 2x - y \cos(xy) \\ &\implies y' = \frac{2x - y \cos(xy)}{x \cos(xy) - 1}\end{aligned}$$

- We can now construct tangents on the curve, *if we know a point on the curve*. Points on the curve will be given, or special x values where y can be solved explicitly will be chosen, or points on the curve can be approximated by numerical methods.

Second Derivatives by Implicit Differentiation

- We can also find second derivatives by implicit differentiation.
Again consider $x^2 + y^2 = 1$.

Second Derivatives by Implicit Differentiation

- We can also find second derivatives by implicit differentiation.

Again consider $x^2 + y^2 = 1$.

- By implicit differentiation we have the first derivative is $y' = -\frac{x}{y}$.

Second Derivatives by Implicit Differentiation

- We can also find second derivatives by implicit differentiation.
Again consider $x^2 + y^2 = 1$.
- By implicit differentiation we have the first derivative is $y' = -\frac{x}{y}$.
- Differentiating a second time by the quotient rule we have

$$y'' = \frac{-y + xy'}{y^2}$$

Second Derivatives by Implicit Differentiation

- We can also find second derivatives by implicit differentiation.

Again consider $x^2 + y^2 = 1$.

- By implicit differentiation we have the first derivative is $y' = -\frac{x}{y}$.
- Differentiating a second time by the quotient rule we have

$$y'' = \frac{-y + xy'}{y^2}$$

- Substituting $y' = -x/y$ we have

$$y'' = \frac{-y + x(-x/y)}{y^2}$$

Second Derivatives by Implicit Differentiation

- We can also find second derivatives by implicit differentiation.

Again consider $x^2 + y^2 = 1$.

- By implicit differentiation we have the first derivative is $y' = -\frac{x}{y}$.
- Differentiating a second time by the quotient rule we have

$$y'' = \frac{-y + xy'}{y^2}$$

- Substituting $y' = -x/y$ we have

$$y'' = \frac{-y + x(-x/y)}{y^2} = -\frac{y^2 + x^2}{y^3}$$

Second Derivatives by Implicit Differentiation

- We can also find second derivatives by implicit differentiation.

Again consider $x^2 + y^2 = 1$.

- By implicit differentiation we have the first derivative is $y' = -\frac{x}{y}$.
- Differentiating a second time by the quotient rule we have

$$y'' = \frac{-y + xy'}{y^2}$$

- Substituting $y' = -x/y$ we have

$$y'' = \frac{-y + x(-x/y)}{y^2} = -\frac{y^2 + x^2}{y^3} = -\frac{1}{y^3}$$

Second Derivatives by Implicit Differentiation

- We can also find second derivatives by implicit differentiation.

Again consider $x^2 + y^2 = 1$.

- By implicit differentiation we have the first derivative is $y' = -\frac{x}{y}$.
- Differentiating a second time by the quotient rule we have

$$y'' = \frac{-y + xy'}{y^2}$$

- Substituting $y' = -x/y$ we have

$$y'' = \frac{-y + x(-x/y)}{y^2} = -\frac{y^2 + x^2}{y^3} = -\frac{1}{y^3}$$

- Compare that with finding the second derivative of the explicit function.

Examples

1. Find y' where $x \tan y = y - 1$.
2. Find y'' if $x^6 + y^6 = 1$.
3. Find the equations of the tangent line and normal line to the curve $x^2 + 4xy + y^2 = 13$ at the point $(2, 1)$.
4. Find the points on the ellipse $x^2 + 2y^2 = 1$ where the tangent line has slope 1.
5. Show that the length of the portion of any tangent line to the astroid $x^{2/3} + y^{2/3} = a^{2/3}$ cut off by the coordinate axes is constant.

Now you should work on Problem Set 2.6. After you have finished it, you should try the following additional exercises from Section 2.6:

2.6 C-level: 1–4, 5–20, 25–30, 31–32, 33–34, 35–38, 39–40;

B-level: 43, 44–45, 46–47;

A-level: 48, 49–52, 53–54, 55, 57–62