

MATH 110 Lecture 3.1

Maximum and Minimum Values

Edward Doolittle

Tuesday, March 3, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Maximum and Minimum Values

Absolute and Local Maxima and Minima

The Extreme Value Theorem

Fermat's Theorem and Critical Numbers

The Closed Interval Method

Examples and Exercises

Maximum and Minimum Values

Introduction to Applications of Differentiation

- The big idea of chapter 2 (Derivatives) was to obtain information about derivatives df/dx when we are given a function $f(x)$.

Introduction to Applications of Differentiation

- The big idea of chapter 2 (Derivatives) was to obtain information about derivatives df/dx when we are given a function $f(x)$.
- The answers in chapter 2 were almost all of the form $dy/dx = \text{something}$ or $dy/dt = \text{something}$.

Introduction to Applications of Differentiation

- The big idea of chapter 2 (Derivatives) was to obtain information about derivatives df/dx when we are given a function $f(x)$.
- The answers in chapter 2 were almost all of the form $dy/dx = \text{something}$ or $dy/dt = \text{something}$.
- The big idea of chapter 3 (Application of Differentiation) is to obtain information about $f(x)$ from the derivative df/dx .

Introduction to Applications of Differentiation

- The big idea of chapter 2 (Derivatives) was to obtain information about derivatives df/dx when we are given a function $f(x)$.
- The answers in chapter 2 were almost all of the form $dy/dx = \text{something}$ or $dy/dt = \text{something}$.
- The big idea of chapter 3 (Application of Differentiation) is to obtain information about $f(x)$ from the derivative df/dx .
- The major classes of applications we will study in this chapter are optimization, curve sketching, and approximation.

Introduction to Applications of Differentiation

- The big idea of chapter 2 (Derivatives) was to obtain information about derivatives df/dx when we are given a function $f(x)$.
- The answers in chapter 2 were almost all of the form $dy/dx = \text{something}$ or $dy/dt = \text{something}$.
- The big idea of chapter 3 (Application of Differentiation) is to obtain information about $f(x)$ from the derivative df/dx .
- The major classes of applications we will study in this chapter are optimization, curve sketching, and approximation.
- In section 3.1 we will study the simplest type of optimization, finding the maximum and minimum values of a continuous function on a closed interval (an interval containing its endpoints).

Absolute Maxima and Minima

- Typically we are interested in the questions “What is the best we can do?” or “What is the most we can get from a situation?”

Absolute Maxima and Minima

- Typically we are interested in the questions “What is the best we can do?” or “What is the most we can get from a situation?”
- In the language of mathematics, if we have a function $f(x)$, we are interested in finding a number c such that $f(c)$ is the biggest number that $f(x)$ takes.

Absolute Maxima and Minima

- Typically we are interested in the questions “What is the best we can do?” or “What is the most we can get from a situation?”
- In the language of mathematics, if we have a function $f(x)$, we are interested in finding a number c such that $f(c)$ is the biggest number that $f(x)$ takes.
- We can write that as $f(c) \geq f(x)$ for all x in the domain of f .

Absolute Maxima and Minima

- Typically we are interested in the questions “What is the best we can do?” or “What is the most we can get from a situation?”
- In the language of mathematics, if we have a function $f(x)$, we are interested in finding a number c such that $f(c)$ is the biggest number that $f(x)$ takes.
- We can write that as $f(c) \geq f(x)$ for all x in the domain of f .
- **Definition:** A function f has an **absolute maximum** or **global maximum** at c if $f(c) \geq f(x)$ for all x in D , the domain of f .

Absolute Maxima and Minima

- Typically we are interested in the questions “What is the best we can do?” or “What is the most we can get from a situation?”
- In the language of mathematics, if we have a function $f(x)$, we are interested in finding a number c such that $f(c)$ is the biggest number that $f(x)$ takes.
- We can write that as $f(c) \geq f(x)$ for all x in the domain of f .
- **Definition:** A function f has an **absolute maximum** or **global maximum** at c if $f(c) \geq f(x)$ for all x in D , the domain of f .
- We can make a similar definition for Absolute minimum values by changing the direction of the inequality.

Absolute Maxima and Minima

- Typically we are interested in the questions “What is the best we can do?” or “What is the most we can get from a situation?”
- In the language of mathematics, if we have a function $f(x)$, we are interested in finding a number c such that $f(c)$ is the biggest number that $f(x)$ takes.
- We can write that as $f(c) \geq f(x)$ for all x in the domain of f .
- **Definition:** A function f has an **absolute maximum** or **global maximum** at c if $f(c) \geq f(x)$ for all x in D , the domain of f .
- We can make a similar definition for Absolute minimum values by changing the direction of the inequality.
- **Definition:** A function f has an **absolute minimum** or **global minimum** at c if $f(c) \leq f(x)$ for all x in D , the domain of f .

Absolute Maxima and Minima

- Typically we are interested in the questions “What is the best we can do?” or “What is the most we can get from a situation?”
- In the language of mathematics, if we have a function $f(x)$, we are interested in finding a number c such that $f(c)$ is the biggest number that $f(x)$ takes.
- We can write that as $f(c) \geq f(x)$ for all x in the domain of f .
- **Definition:** A function f has an **absolute maximum** or **global maximum** at c if $f(c) \geq f(x)$ for all x in D , the domain of f .
- We can make a similar definition for Absolute minimum values by changing the direction of the inequality.
- **Definition:** A function f has an **absolute minimum** or **global minimum** at c if $f(c) \leq f(x)$ for all x in D , the domain of f .
- The maximum and minimum values of f (i.e., $f(c)$ in the above, not c) are called the **extreme values** of f .

Local Maxima and Minima

- Imagine you are standing on a hill. You are standing higher than anything immediately around you. You can't go any higher by moving in any direction. But you are not standing on the highest point on earth (Mount Everest).

Local Maxima and Minima

- Imagine you are standing on a hill. You are standing higher than anything immediately around you. You can't go any higher by moving in any direction. But you are not standing on the highest point on earth (Mount Everest).
- In that example, the peak of Mount Everest is the absolute maximum, while your hill is a local maximum. Local maxima and minima are useful and interesting in their own right.

Local Maxima and Minima

- Imagine you are standing on a hill. You are standing higher than anything immediately around you. You can't go any higher by moving in any direction. But you are not standing on the highest point on earth (Mount Everest).
- In that example, the peak of Mount Everest is the absolute maximum, while your hill is a local maximum. Local maxima and minima are useful and interesting in their own right.
- **Definition:** A function f has a **local maximum** or **relative maximum** at c if $f(c) \geq f(x)$ when x is near c .

Local Maxima and Minima

- Imagine you are standing on a hill. You are standing higher than anything immediately around you. You can't go any higher by moving in any direction. But you are not standing on the highest point on earth (Mount Everest).
- In that example, the peak of Mount Everest is the absolute maximum, while your hill is a local maximum. Local maxima and minima are useful and interesting in their own right.
- **Definition:** A function f has a **local maximum** or **relative maximum** at c if $f(c) \geq f(x)$ when x is near c .
- **Definition:** A function f has a **local minimum** or **relative minimum** at c if $f(c) \leq f(x)$ when x is near c .

Local Maxima and Minima

- Imagine you are standing on a hill. You are standing higher than anything immediately around you. You can't go any higher by moving in any direction. But you are not standing on the highest point on earth (Mount Everest).
- In that example, the peak of Mount Everest is the absolute maximum, while your hill is a local maximum. Local maxima and minima are useful and interesting in their own right.
- **Definition:** A function f has a **local maximum** or **relative maximum** at c if $f(c) \geq f(x)$ when x is near c .
- **Definition:** A function f has a **local minimum** or **relative minimum** at c if $f(c) \leq f(x)$ when x is near c .
- To be a little more precise, “ x is near c ” means “in some open interval containing c ”. An open interval is an interval which does not include its endpoints.

Existence of Absolute Maxima and Minima

- One of the first questions we should ask is “when is a function guaranteed to have an absolute maximum and minimum value?”

Existence of Absolute Maxima and Minima

- One of the first questions we should ask is “when is a function guaranteed to have an absolute maximum and minimum value?”
- We could waste a lot of time looking for absolute maxima and minima if they don't exist.

Existence of Absolute Maxima and Minima

- One of the first questions we should ask is “when is a function guaranteed to have an absolute maximum and minimum value?”
- We could waste a lot of time looking for absolute maxima and minima if they don't exist.
- Fortunately, in many common situations we can guarantee the existence of maxima and minima.

Existence of Absolute Maxima and Minima

- One of the first questions we should ask is “when is a function guaranteed to have an absolute maximum and minimum value?”
- We could waste a lot of time looking for absolute maxima and minima if they don't exist.
- Fortunately, in many common situations we can guarantee the existence of maxima and minima.
- The situation we will concentrate on in this section is when f is a continuous function defined on a closed domain $D = [a, b]$.

Existence of Absolute Maxima and Minima

- One of the first questions we should ask is “when is a function guaranteed to have an absolute maximum and minimum value?”
- We could waste a lot of time looking for absolute maxima and minima if they don't exist.
- Fortunately, in many common situations we can guarantee the existence of maxima and minima.
- The situation we will concentrate on in this section is when f is a continuous function defined on a closed domain $D = [a, b]$.
- **Extreme Value Theorem:** If f is continuous on a closed interval $[a, b]$ then f attains an absolute maximum value $f(c)$ and an absolute minimum value $f(d)$ for some numbers c and d in $[a, b]$.

Existence of Absolute Maxima and Minima

- One of the first questions we should ask is “when is a function guaranteed to have an absolute maximum and minimum value?”
- We could waste a lot of time looking for absolute maxima and minima if they don't exist.
- Fortunately, in many common situations we can guarantee the existence of maxima and minima.
- The situation we will concentrate on in this section is when f is a continuous function defined on a closed domain $D = [a, b]$.
- **Extreme Value Theorem:** If f is continuous on a closed interval $[a, b]$ then f attains an absolute maximum value $f(c)$ and an absolute minimum value $f(d)$ for some numbers c and d in $[a, b]$.
- The Extreme Value Theorem is similar to the Intermediate Value Theorem in chapter 1. Both are hard to prove. The idea is to use a successive approximations process.

Location of Extreme Values

- Note that a maximum can occur at a point c between the endpoints a and b of the interval, or it can occur at an endpoint a or b .

Location of Extreme Values

- Note that a maximum can occur at a point c between the endpoints a and b of the interval, or it can occur at an endpoint a or b .
- Similarly, a minimum can occur between a and b , or at a , or at b .

Location of Extreme Values

- Note that a maximum can occur at a point c between the endpoints a and b of the interval, or it can occur at an endpoint a or b .
- Similarly, a minimum can occur between a and b , or at a , or at b .
- There may also be multiple numbers c at which the function attains its maximum. The Extreme Value Theorem guarantees one but there may be more.

Fermat's Theorem

- We want to find local extrema of a given function f .

Fermat's Theorem

- We want to find local extrema of a given function f .
- Fortunately, the derivative f' will help us find extrema.

Fermat's Theorem

- We want to find local extrema of a given function f .
- Fortunately, the derivative f' will help us find extrema.
- The idea is that if we are at a local maximum, we can't go any higher.

Fermat's Theorem

- We want to find local extrema of a given function f .
- Fortunately, the derivative f' will help us find extrema.
- The idea is that if we are at a local maximum, we can't go any higher.
- If $f'(c) > 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx > 0$; that means that if $f'(c) > 0$, f can't have a maximum at c .

Fermat's Theorem

- We want to find local extrema of a given function f .
- Fortunately, the derivative f' will help us find extrema.
- The idea is that if we are at a local maximum, we can't go any higher.
- If $f'(c) > 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx > 0$; that means that if $f'(c) > 0$, f can't have a maximum at c .
- Similarly, if $f'(c) < 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx < 0$; that means that if $f'(c) < 0$, f can't have a maximum at c either.

Fermat's Theorem

- We want to find local extrema of a given function f .
- Fortunately, the derivative f' will help us find extrema.
- The idea is that if we are at a local maximum, we can't go any higher.
- If $f'(c) > 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx > 0$; that means that if $f'(c) > 0$, f can't have a maximum at c .
- Similarly, if $f'(c) < 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx < 0$; that means that if $f'(c) < 0$, f can't have a maximum at c either.
- Similar reasoning applies for minima.

Fermat's Theorem

- We want to find local extrema of a given function f .
- Fortunately, the derivative f' will help us find extrema.
- The idea is that if we are at a local maximum, we can't go any higher.
- If $f'(c) > 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx > 0$; that means that if $f'(c) > 0$, f can't have a maximum at c .
- Similarly, if $f'(c) < 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx < 0$; that means that if $f'(c) < 0$, f can't have a maximum at c either.
- Similar reasoning applies for minima.
- **Fermat's Theorem:** If f has a local maximum or minimum at c , and $f'(c)$ exists, then $f'(c) = 0$.

Fermat's Theorem

- We want to find local extrema of a given function f .
- Fortunately, the derivative f' will help us find extrema.
- The idea is that if we are at a local maximum, we can't go any higher.
- If $f'(c) > 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx > 0$; that means that if $f'(c) > 0$, f can't have a maximum at c .
- Similarly, if $f'(c) < 0$, then $f(c + dx) \approx f(c) + f'(c)dx > f(c)$ for $dx < 0$; that means that if $f'(c) < 0$, f can't have a maximum at c either.
- Similar reasoning applies for minima.
- **Fermat's Theorem:** If f has a local maximum or minimum at c , and $f'(c)$ exists, then $f'(c) = 0$.
- For those of you who are interested, a rigorous proof of Fermat's Theorem is in the textbook.

Critical Numbers

- Unfortunately, the analysis of the local maxima and minima of a function is not so simple if f is not differentiable everywhere.

Critical Numbers

- Unfortunately, the analysis of the local maxima and minima of a function is not so simple if f is not differentiable everywhere.
- We introduce a new definition to help us single out all the points of interest.

Critical Numbers

- Unfortunately, the analysis of the local maxima and minima of a function is not so simple if f is not differentiable everywhere.
- We introduce a new definition to help us single out all the points of interest.
- **Definition:** A **critical number** of a function f is a number c such that either $f'(c) = 0$ or $f'(c)$ does not exist.

Critical Numbers

- Unfortunately, the analysis of the local maxima and minima of a function is not so simple if f is not differentiable everywhere.
- We introduce a new definition to help us single out all the points of interest.
- **Definition:** A **critical number** of a function f is a number c such that either $f'(c) = 0$ or $f'(c)$ does not exist.
- With that new definition, Fermat's Theorem can be re-phrased as follows:

Critical Numbers

- Unfortunately, the analysis of the local maxima and minima of a function is not so simple if f is not differentiable everywhere.
- We introduce a new definition to help us single out all the points of interest.
- **Definition:** A **critical number** of a function f is a number c such that either $f'(c) = 0$ or $f'(c)$ does not exist.
- With that new definition, Fermat's Theorem can be re-phrased as follows:
- **Fermat's Theorem:** If f has a local maximum or minimum at c , then c is a critical number of f .

The Closed Interval Method for Optimization

- We now turn to the problem of finding the absolute maximum and minimum values of a function f .

The Closed Interval Method for Optimization

- We now turn to the problem of finding the absolute maximum and minimum values of a function f .
- For the time being, the only situation in which we are guaranteed absolute maximum and minimum values is the context of the Extreme Value Theorem, we concentrate on continuous functions f defined on a closed interval $[a, b]$.

The Closed Interval Method for Optimization

- We now turn to the problem of finding the absolute maximum and minimum values of a function f .
- For the time being, the only situation in which we are guaranteed absolute maximum and minimum values is the context of the Extreme Value Theorem, we concentrate on continuous functions f defined on a closed interval $[a, b]$.
- A process which is guaranteed to find the absolute maximum and minimum values of such an f is as follows:

The Closed Interval Method for Optimization

- We now turn to the problem of finding the absolute maximum and minimum values of a function f .
- For the time being, the only situation in which we are guaranteed absolute maximum and minimum values is the context of the Extreme Value Theorem, we concentrate on continuous functions f defined on a closed interval $[a, b]$.
- A process which is guaranteed to find the absolute maximum and minimum values of such an f is as follows:
 1. Find the critical numbers c of f in the open interval (a, b)

The Closed Interval Method for Optimization

- We now turn to the problem of finding the absolute maximum and minimum values of a function f .
- For the time being, the only situation in which we are guaranteed absolute maximum and minimum values is the context of the Extreme Value Theorem, we concentrate on continuous functions f defined on a closed interval $[a, b]$.
- A process which is guaranteed to find the absolute maximum and minimum values of such an f is as follows:
 1. Find the critical numbers c of f in the open interval (a, b)
 2. Find the values of f at each of those critical numbers.

The Closed Interval Method for Optimization

- We now turn to the problem of finding the absolute maximum and minimum values of a function f .
- For the time being, the only situation in which we are guaranteed absolute maximum and minimum values is the context of the Extreme Value Theorem, we concentrate on continuous functions f defined on a closed interval $[a, b]$.
- A process which is guaranteed to find the absolute maximum and minimum values of such an f is as follows:
 1. Find the critical numbers c of f in the open interval (a, b)
 2. Find the values of f at each of those critical numbers.
 3. Find the values of f at each of the endpoints a and b .

The Closed Interval Method for Optimization

- We now turn to the problem of finding the absolute maximum and minimum values of a function f .
- For the time being, the only situation in which we are guaranteed absolute maximum and minimum values is the context of the Extreme Value Theorem, we concentrate on continuous functions f defined on a closed interval $[a, b]$.
- A process which is guaranteed to find the absolute maximum and minimum values of such an f is as follows:
 1. Find the critical numbers c of f in the open interval (a, b)
 2. Find the values of f at each of those critical numbers.
 3. Find the values of f at each of the endpoints a and b .
 4. The largest of the values from the previous two steps is the absolute maximum value; the smallest of the values is the absolute minimum value.

The Closed Interval Theorem in Action

- For example, let's try find the absolute extrema of the function $f(x) = x^4 - 2x^2 + 3$ on the interval $[-2, 3]$.

The Closed Interval Theorem in Action

- For example, let's try find the absolute extrema of the function $f(x) = x^4 - 2x^2 + 3$ on the interval $[-2, 3]$.
 1. We find the critical numbers of f by noting where $f'(x) = 4x^3 - 4x$ exists (everywhere) and solving the equation $f'(c) = 0$ i.e., $4c(c-1)(c+1) = 0$ which gives critical numbers $c = 0, c = -1, c = 1$.

The Closed Interval Theorem in Action

- For example, let's try find the absolute extrema of the function $f(x) = x^4 - 2x^2 + 3$ on the interval $[-2, 3]$.
 1. We find the critical numbers of f by noting where $f'(x) = 4x^3 - 4x$ exists (everywhere) and solving the equation $f'(c) = 0$ i.e., $4c(c-1)(c+1) = 0$ which gives critical numbers $c = 0, c = -1, c = 1$.
 2. We evaluate f at each of those critical numbers: $f(0) = 3$, $f(-1) = 2, f(1) = 2$.

The Closed Interval Theorem in Action

- For example, let's try find the absolute extrema of the function $f(x) = x^4 - 2x^2 + 3$ on the interval $[-2, 3]$.
 1. We find the critical numbers of f by noting where $f'(x) = 4x^3 - 4x$ exists (everywhere) and solving the equation $f'(c) = 0$ i.e., $4c(c-1)(c+1) = 0$ which gives critical numbers $c = 0, c = -1, c = 1$.
 2. We evaluate f at each of those critical numbers: $f(0) = 3$, $f(-1) = 2, f(1) = 2$.
 3. We evaluate f at the endpoints of the domain: $f(-2) = 11$, $f(3) = 66$.

The Closed Interval Theorem in Action

- For example, let's try find the absolute extrema of the function $f(x) = x^4 - 2x^2 + 3$ on the interval $[-2, 3]$.
 1. We find the critical numbers of f by noting where $f'(x) = 4x^3 - 4x$ exists (everywhere) and solving the equation $f'(c) = 0$ i.e., $4c(c - 1)(c + 1) = 0$ which gives critical numbers $c = 0, c = -1, c = 1$.
 2. We evaluate f at each of those critical numbers: $f(0) = 3$, $f(-1) = 2, f(1) = 2$.
 3. We evaluate f at the endpoints of the domain: $f(-2) = 11$, $f(3) = 66$.
 4. The largest of the above values is 66, so f has a maximum value of 66 at the endpoint $b = 3$. The smallest of the above values is 2, so f has a minimum value of 2 at both interior points $c = -1$ and $c = 1$.

Examples and Exercises

Examples

1. Find the critical numbers of the given functions.

1.1 $f(x) = x^3 + 3x^2 - 24x$

1.2 $s(t) = 3t^4 + 4t^3 - 6t^2$

1.3 $g(y) = \frac{y-1}{y^2-y+1}$

2. Find the local and absolute extreme values of the function on the given interval.

2.1 $f(x) = x\sqrt{1-x}, [-1, 1]$

2.2 $f(x) = (x^2 + 2x)^3, [-2, 1]$

2.3 $f(x) = \sin x + \cos^2 x, [0, \pi]$

3. Between 0° C and 30° C , the volume V (in cm^3) of 1 kg of water at temperature T is given approximately by

$$V = 999.87 - 0.06426T + 0.0085043T^2 - 0.0000679T^3$$

Find the temperature at which water has its maximum density.

Now you should work on Problem Set 3.1. After you have finished it, you should try the following additional exercises from Section 3.1:

3.1 C-level: 1–42, 45–56; B-level: 57–65, 67–68; A-level: 43–44, 69–70, 72