

MATH 110 Lecture 4.2

The Definite Integral

Edward Doolittle

Thursday, March 26, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

The Definite Integral

Definite Integrals

Evaluating Integrals

Properties of Integrals

Examples and Exercises

The Definite Integral

Definition of a Definite Integral

- We have seen expressions like $\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$ for areas.

Definition of a Definite Integral

- We have seen expressions like $\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$ for areas.
- Such expressions show up in other contexts, e.g., calculating work done. They are so common that they have a special notation.

Definition of a Definite Integral

- We have seen expressions like $\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$ for areas.
- Such expressions show up in other contexts, e.g., calculating work done. They are so common that they have a special notation.
- **Definition:** Let f be a function on an interval $[a, b]$. Divide the interval into equal subintervals of length $\Delta x = (b - a)/n$ and let $x_0 = a, x_1 = a + (1/n)(b - a), \dots, x_n = a + (n/n)(b - a)/n = b$ be the endpoints of those intervals. Let x_i^* be any **sample points** in those intervals. Then the **definite integral of f from a to b** is

$$\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$$

provided that the limit exists. If it does exist, we say f is integrable.

Notation

- The symbol \int is called an integral sign.

Notation

- The symbol \int is called an integral sign.
- It's an elongated S and stands for the Latin word “summa” for sum.

Notation

- The symbol \int is called an integral sign.
- It's an elongated S and stands for the Latin word “summa” for sum.
- In the notation $\int_a^b f(x) dx$, $f(x)$ is called the **integrand**.

Notation

- The symbol \int is called an integral sign.
- It's an elongated S and stands for the Latin word “summa” for sum.
- In the notation $\int_a^b f(x) dx$, $f(x)$ is called the **integrand**.
- a and b are called the **limits of integration**; a is the **lower limit** and b is the **upper limit**.

- EJD

- EJD

Examples and Exercises

Examples

1. 1.1 Evaluate the Riemann sum for $f(x) = x^2 - x$, $0 \leq x \leq 2$ with four subintervals, taking the sample points to be right endpoints. Explain, with the aid of a diagram, what the Riemann sum represents.
1.2 Use the definition of a definite integral (with right endpoints) to calculate the value of the integral $\int_0^2 (x^2 - x) dx$.
2. Evaluate $\int_0^1 (x + \sqrt{1 - x^2}) dx$ by interpreting it in terms of areas.
3. If $\int_0^6 f(x) dx = 10$ and $\int_0^4 f(x) dx = 7$, find $\int_4^6 f(x) dx$.

Now you should work on Problem Set 4.2. After you have finished it, you should try the following additional exercises from Section 4.2:

4.2 C-level: 1–8, 17–30, 33–51;

B-level: 52–64;

A-level: 65–75