

MATH 110 Lecture 2.3

Differentiation Formulas

Edward Doolittle

Tuesday, February 3, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Differentiation Formulas

Bottom Level Rules for Derivatives

Reduction Formulas for Derivatives

Application: Tangent and Normal Lines

Examples and Exercises

Differentiation Formulas

Review of the Reduction Strategy for Limits

- Recall when we evaluated limits, we applied limit laws to break a complicated limit into simpler parts.

Review of the Reduction Strategy for Limits

- Recall when we evaluated limits, we applied limit laws to break a complicated limit into simpler parts.
- For example, we can apply the sum rule to the limit

$$\lim_{x \rightarrow 2} (3x + 7) = \lim_{x \rightarrow 2} (3x) + \lim_{x \rightarrow 2} 7$$

to break a complicated limit into simpler parts.

Review of the Reduction Strategy for Limits

- Recall when we evaluated limits, we applied limit laws to break a complicated limit into simpler parts.
- For example, we can apply the sum rule to the limit

$$\lim_{x \rightarrow 2} (3x + 7) = \lim_{x \rightarrow 2} (3x) + \lim_{x \rightarrow 2} 7$$

to break a complicated limit into simpler parts.

- The first summand can be broken down further by the product rule for limits:

$$\lim_{x \rightarrow 2} (3x) = (\lim_{x \rightarrow 2} 3)(\lim_{x \rightarrow 2} x)$$

Review of the Reduction Strategy for Limits

- Recall when we evaluated limits, we applied limit laws to break a complicated limit into simpler parts.
- For example, we can apply the sum rule to the limit

$$\lim_{x \rightarrow 2} (3x + 7) = \lim_{x \rightarrow 2} (3x) + \lim_{x \rightarrow 2} 7$$

to break a complicated limit into simpler parts.

- The first summand can be broken down further by the product rule for limits:

$$\lim_{x \rightarrow 2} (3x) = (\lim_{x \rightarrow 2} 3)(\lim_{x \rightarrow 2} x)$$

- The resulting simple limits can be evaluated by applying the bottom level rules $\lim_{x \rightarrow 2} c = c$ for $c = 3$ and $c = 7$, and $\lim_{x \rightarrow 2} x = 2$.

Review of the Reduction Strategy for Limits

- Recall when we evaluated limits, we applied limit laws to break a complicated limit into simpler parts.
- For example, we can apply the sum rule to the limit

$$\lim_{x \rightarrow 2} (3x + 7) = \lim_{x \rightarrow 2} (3x) + \lim_{x \rightarrow 2} 7$$

to break a complicated limit into simpler parts.

- The first summand can be broken down further by the product rule for limits:

$$\lim_{x \rightarrow 2} (3x) = (\lim_{x \rightarrow 2} 3)(\lim_{x \rightarrow 2} x)$$

- The resulting simple limits can be evaluated by applying the bottom level rules $\lim_{x \rightarrow 2} c = c$ for $c = 3$ and $c = 7$, and $\lim_{x \rightarrow 2} x = 2$.
- We would like to develop a similar strategy for evaluating derivatives.

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.
- These rules are slightly different from the corresponding rules for limits.

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.
- These rules are slightly different from the corresponding rules for limits.
- First, suppose $f(x) = c$, a constant. Then

$$\frac{d}{dx}f(x)$$

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.
- These rules are slightly different from the corresponding rules for limits.
- First, suppose $f(x) = c$, a constant. Then

$$\frac{d}{dx}f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.
- These rules are slightly different from the corresponding rules for limits.
- First, suppose $f(x) = c$, a constant. Then

$$\frac{d}{dx}f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h}$$

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.
- These rules are slightly different from the corresponding rules for limits.
- First, suppose $f(x) = c$, a constant. Then

$$\frac{d}{dx}f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = \lim_{h \rightarrow 0} \frac{0}{h}$$

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.
- These rules are slightly different from the corresponding rules for limits.
- First, suppose $f(x) = c$, a constant. Then

$$\frac{d}{dx}f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = 0$$

Bottom Level Rules for Derivatives: Constant Functions

- Let us begin this time with some bottom-level rules for derivatives.
- These rules are slightly different from the corresponding rules for limits.
- First, suppose $f(x) = c$, a constant. Then

$$\frac{d}{dx}f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = 0$$

- In words, the derivative of a constant is 0.

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x)$$

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{x+h-x}{h}$$

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{x+h-x}{h} = \lim_{h \rightarrow 0} \frac{h}{h}$$

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{x+h-x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{x+h-x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

- In words, the derivative of the identity function is 1.

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{x+h-x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

- In words, the derivative of the identity function is 1.
- The derivatives of constant functions and of the identity function will be the building blocks for most of the derivative calculations in this section.

Bottom Level Rules for Derivatives: the Identity Function

- Now let $f(x) = x$. That function is known as the ‘identity function’ because the output of f is identical to the input.
- Now, calculating the derivative,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{x+h-x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

- In words, the derivative of the identity function is 1.
- The derivatives of constant functions and of the identity function will be the building blocks for most of the derivative calculations in this section.
- There are other bottom-level derivatives that we will learn later. Each time we introduce a new kind of function (e.g., sin, cos, exp, log, ...) there will be a new bottom-level derivative that will have to be calculated from first principles.

Sums, Differences, and Constant Multiples for Derivatives

- As with limits, derivatives behave in a simple way with respect to sums, differences, and constant multiples.

Sums, Differences, and Constant Multiples for Derivatives

- As with limits, derivatives behave in a simple way with respect to sums, differences, and constant multiples.
- The derivative of a sum is the sum of the derivatives

$$(f + g)' = f' + g' \quad \text{or} \quad \frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$

Sums, Differences, and Constant Multiples for Derivatives

- As with limits, derivatives behave in a simple way with respect to sums, differences, and constant multiples.
- The derivative of a sum is the sum of the derivatives

$$(f + g)' = f' + g' \quad \text{or} \quad \frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$

- The derivative of a difference is the difference of the derivatives.

$$(f - g)' = f' - g' \quad \text{or} \quad \frac{d}{dx}(f(x) - g(x)) = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$$

Sums, Differences, and Constant Multiples for Derivatives

- As with limits, derivatives behave in a simple way with respect to sums, differences, and constant multiples.
- The derivative of a sum is the sum of the derivatives

$$(f + g)' = f' + g' \quad \text{or} \quad \frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$

- The derivative of a difference is the difference of the derivatives.

$$(f - g)' = f' - g' \quad \text{or} \quad \frac{d}{dx}(f(x) - g(x)) = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$$

- The derivative of a constant multiple is the constant multiple of the derivative.

$$(cf)' = cf' \quad \text{or} \quad \frac{d}{dx}(cf(x)) = c \frac{d}{dx}f(x)$$

Justifying the Sum Rule for Derivatives

- The sum, difference, and constant multiple rules are direct consequences of the corresponding rules for limits.

Justifying the Sum Rule for Derivatives

- The sum, difference, and constant multiple rules are direct consequences of the corresponding rules for limits.
- For example, we have

$$(f + g)'(x)$$

Justifying the Sum Rule for Derivatives

- The sum, difference, and constant multiple rules are direct consequences of the corresponding rules for limits.
- For example, we have

$$(f + g)'(x) = \lim_{h \rightarrow 0} \frac{(f + g)(x + h) - (f + g)(x)}{h}$$

Justifying the Sum Rule for Derivatives

- The sum, difference, and constant multiple rules are direct consequences of the corresponding rules for limits.
- For example, we have

$$\begin{aligned}(f + g)'(x) &= \lim_{h \rightarrow 0} \frac{(f + g)(x + h) - (f + g)(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x + h) + g(x + h) - f(x) - g(x)}{h}\end{aligned}$$

Justifying the Sum Rule for Derivatives

- The sum, difference, and constant multiple rules are direct consequences of the corresponding rules for limits.
- For example, we have

$$\begin{aligned}(f + g)'(x) &= \lim_{h \rightarrow 0} \frac{(f + g)(x + h) - (f + g)(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{f(x + h) + g(x + h) - f(x) - g(x)}{h} \\&= \lim_{h \rightarrow 0} \left(\frac{f(x + h) - f(x)}{h} + \frac{g(x + h) - g(x)}{h} \right)\end{aligned}$$

Justifying the Sum Rule for Derivatives

- The sum, difference, and constant multiple rules are direct consequences of the corresponding rules for limits.
- For example, we have

$$\begin{aligned}(f + g)'(x) &= \lim_{h \rightarrow 0} \frac{(f + g)(x + h) - (f + g)(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{f(x + h) + g(x + h) - f(x) - g(x)}{h} \\&= \lim_{h \rightarrow 0} \left(\frac{f(x + h) - f(x)}{h} + \frac{g(x + h) - g(x)}{h} \right)\end{aligned}$$

- Now we apply the sum rule for limits to show the latter expression is the same as $f'(x) + g'(x)$.

First Attempt at a Product Rule for Derivatives

- We've had some success we've had in guessing rules for derivatives from rules from the rules for limits.

First Attempt at a Product Rule for Derivatives

- We've had some success we've had in guessing rules for derivatives from rules from the rules for limits.
- It might seem natural to guess that the derivative of a product is the product of the derivatives.

First Attempt at a Product Rule for Derivatives

- We've had some success we've had in guessing rules for derivatives from rules from the rules for limits.
- It might seem natural to guess that the derivative of a product is the product of the derivatives.
- That is incorrect, however. Consider the example $f(x) = 1$, $g(x) = x$.

First Attempt at a Product Rule for Derivatives

- We've had some success we've had in guessing rules for derivatives from rules from the rules for limits.
- It might seem natural to guess that the derivative of a product is the product of the derivatives.
- That is incorrect, however. Consider the example $f(x) = 1$, $g(x) = x$.
- The derivative of the product is the derivative of x which is 1.

First Attempt at a Product Rule for Derivatives

- We've had some success we've had in guessing rules for derivatives from rules from the rules for limits.
- It might seem natural to guess that the derivative of a product is the product of the derivatives.
- That is incorrect, however. Consider the example $f(x) = 1$, $g(x) = x$.
- The derivative of the product is the derivative of x which is 1.
- The derivative of $f(x)$ is 0 and the derivative of $g(x)$ is 1, so the product of the derivatives is 0.

First Attempt at a Product Rule for Derivatives

- We've had some success we've had in guessing rules for derivatives from rules from the rules for limits.
- It might seem natural to guess that the derivative of a product is the product of the derivatives.
- That is incorrect, however. Consider the example $f(x) = 1$, $g(x) = x$.
- The derivative of the product is the derivative of x which is 1.
- The derivative of $f(x)$ is 0 and the derivative of $g(x)$ is 1, so the product of the derivatives is 0.
- Even in this simple case, the derivative of the product (1) is different from the product of the derivatives (0).

First Attempt at a Product Rule for Derivatives

- We've had some success we've had in guessing rules for derivatives from rules from the rules for limits.
- It might seem natural to guess that the derivative of a product is the product of the derivatives.
- That is incorrect, however. Consider the example $f(x) = 1$, $g(x) = x$.
- The derivative of the product is the derivative of x which is 1.
- The derivative of $f(x)$ is 0 and the derivative of $g(x)$ is 1, so the product of the derivatives is 0.
- Even in this simple case, the derivative of the product (1) is different from the product of the derivatives (0).
- What is the correct rule?

Discovering the Product Rule for Derivatives

- The product rule for derivatives is not what we might naively expect. $(fg)'$ is **not** $f'g'$!

Discovering the Product Rule for Derivatives

- The product rule for derivatives is not what we might naively expect. $(fg)'$ is **not** $f'g'$!
- To understand what the derivative of a product should be, consider the meaning of the derivative. If I change x by a small amount dx , this is what happens to f :

$$f(x + dx) = f(x) + f'(x)dx + \text{terms in } dx^2, \text{ etc.}$$

Discovering the Product Rule for Derivatives

- The product rule for derivatives is not what we might naively expect. $(fg)'$ is **not** $f'g'$!
- To understand what the derivative of a product should be, consider the meaning of the derivative. If I change x by a small amount dx , this is what happens to f :

$$f(x + dx) = f(x) + f'(x)dx + \text{terms in } dx^2, \text{ etc.}$$

- Now consider what happens to the product $f(x)g(x)$ when we change x by a small amount dx :

$$\begin{aligned} f(x + dx)g(x + dx) &= (f(x) + f'(x)dx + \cdots)(g(x) + g'(x)dx + \cdots) \\ &= f(x)g(x) + (f'(x)g(x) + f(x)g'(x)) dx + \cdots \end{aligned}$$

Discovering the Product Rule for Derivatives

- The product rule for derivatives is not what we might naively expect. $(fg)'$ is **not** $f'g'$!
- To understand what the derivative of a product should be, consider the meaning of the derivative. If I change x by a small amount dx , this is what happens to f :

$$f(x + dx) = f(x) + f'(x)dx + \text{terms in } dx^2, \text{ etc.}$$

- Now consider what happens to the product $f(x)g(x)$ when we change x by a small amount dx :

$$\begin{aligned} f(x + dx)g(x + dx) &= (f(x) + f'(x)dx + \cdots)(g(x) + g'(x)dx + \cdots) \\ &= f(x)g(x) + (f'(x)g(x) + f(x)g'(x)) dx + \cdots \end{aligned}$$

- This tells us that $(f(x)g(x))'$ should be $f'(x)g(x) + f(x)g'(x)$.

The Product Rule for Derivatives

- In summary, we have the formula in Leibniz notation

$$\frac{d}{dx} (f(x)g(x)) = \left(\frac{d}{dx} f(x) \right) g(x) + f(x) \left(\frac{d}{dx} g(x) \right)$$

The Product Rule for Derivatives

- In summary, we have the formula in Leibniz notation

$$\frac{d}{dx}(f(x)g(x)) = \left(\frac{d}{dx}f(x)\right)g(x) + f(x)\left(\frac{d}{dx}g(x)\right)$$

- Or, in more compact Newton notation,

$$(fg)' = f'g + fg'$$

The Product Rule for Derivatives

- In summary, we have the formula in Leibniz notation

$$\frac{d}{dx}(f(x)g(x)) = \left(\frac{d}{dx}f(x)\right)g(x) + f(x)\left(\frac{d}{dx}g(x)\right)$$

- Or, in more compact Newton notation,

$$(fg)' = f'g + fg'$$

- Note that the derivative of a product **is not** the product of the derivatives!

The Product Rule for Derivatives

- In summary, we have the formula in Leibniz notation

$$\frac{d}{dx} (f(x)g(x)) = \left(\frac{d}{dx} f(x) \right) g(x) + f(x) \left(\frac{d}{dx} g(x) \right)$$

- Or, in more compact Newton notation,

$$(fg)' = f'g + fg'$$

- Note that the derivative of a product **is not** the product of the derivatives!
- Derivatives differ from limits in this regard.

Discovering the Quotient Rule for Derivatives

- In a similar way, the quotient rule for derivatives is not what we might naively expect. Consider $f(x) = x$, $g(x) = x$. Then $(f/g)' = 0$ but $f'/g' = 1$.

Discovering the Quotient Rule for Derivatives

- In a similar way, the quotient rule for derivatives is not what we might naively expect. Consider $f(x) = x$, $g(x) = x$. Then $(f/g)' = 0$ but $f'/g' = 1$.
- We can figure out what the correct rule must be by differentiating the equation $f = g(f/g)$ using the product rule.

Discovering the Quotient Rule for Derivatives

- In a similar way, the quotient rule for derivatives is not what we might naively expect. Consider $f(x) = x$, $g(x) = x$. Then $(f/g)' = 0$ but $f'/g' = 1$.
- We can figure out what the correct rule must be by differentiating the equation $f = g(f/g)$ using the product rule.
- We have $f' = g'(f/g) + g(f/g)'$.

Discovering the Quotient Rule for Derivatives

- In a similar way, the quotient rule for derivatives is not what we might naively expect. Consider $f(x) = x$, $g(x) = x$. Then $(f/g)' = 0$ but $f'/g' = 1$.
- We can figure out what the correct rule must be by differentiating the equation $f = g(f/g)$ using the product rule.
- We have $f' = g'(f/g) + g(f/g)'$.
- Solving for $(f/g)'$, we have

$$\left(\frac{f}{g}\right)' = \frac{gf' - fg'}{g^2}$$

Discovering the Quotient Rule for Derivatives

- In a similar way, the quotient rule for derivatives is not what we might naively expect. Consider $f(x) = x$, $g(x) = x$. Then $(f/g)' = 0$ but $f'/g' = 1$.
- We can figure out what the correct rule must be by differentiating the equation $f = g(f/g)$ using the product rule.
- We have $f' = g'(f/g) + g(f/g)'$.
- Solving for $(f/g)'$, we have

$$\left(\frac{f}{g}\right)' = \frac{gf' - fg'}{g^2}$$

- The mantra “bottom times derivative of top minus top times derivative of bottom over bottom squared” may help you remember the quotient rule.

Example: Derivatives of Monomials

- We now know enough to do some complicated derivatives.

Example: Derivatives of Monomials

- We now know enough to do some complicated derivatives.
- Consider the derivative of x^2 . We can write x^2 as a product $x \cdot x$.

Example: Derivatives of Monomials

- We now know enough to do some complicated derivatives.
- Consider the derivative of x^2 . We can write x^2 as a product $x \cdot x$.
- The product rule then tells us that

$$\frac{d}{dx}x^2 = \frac{d}{dx}(x \cdot x) = \left(\frac{d}{dx}x\right)x + x\left(\frac{d}{dx}x\right) = x + x = 2x$$

Example: Derivatives of Monomials

- We now know enough to do some complicated derivatives.
- Consider the derivative of x^2 . We can write x^2 as a product $x \cdot x$.
- The product rule then tells us that

$$\frac{d}{dx}x^2 = \frac{d}{dx}(x \cdot x) = \left(\frac{d}{dx}x\right)x + x\left(\frac{d}{dx}x\right) = x + x = 2x$$

- Using the previous result, the product rule tells us that

$$\frac{d}{dx}x^3 = \frac{d}{dx}(x \cdot x^2) = \left(\frac{d}{dx}x\right)x^2 + x\left(\frac{d}{dx}x^2\right) = x^2 + x \cdot 2x = 3x^2$$

Example: Derivatives of Monomials

- We now know enough to do some complicated derivatives.
- Consider the derivative of x^2 . We can write x^2 as a product $x \cdot x$.
- The product rule then tells us that

$$\frac{d}{dx}x^2 = \frac{d}{dx}(x \cdot x) = \left(\frac{d}{dx}x\right)x + x\left(\frac{d}{dx}x\right) = x + x = 2x$$

- Using the previous result, the product rule tells us that

$$\frac{d}{dx}x^3 = \frac{d}{dx}(x \cdot x^2) = \left(\frac{d}{dx}x\right)x^2 + x\left(\frac{d}{dx}x^2\right) = x^2 + x \cdot 2x = 3x^2$$

- Continuing the pattern, we have $\frac{d}{dx}x^n = nx^{n-1}$ for $n = 1, 2, \dots$

Example: Derivatives of Monomials

- We now know enough to do some complicated derivatives.
- Consider the derivative of x^2 . We can write x^2 as a product $x \cdot x$.
- The product rule then tells us that

$$\frac{d}{dx}x^2 = \frac{d}{dx}(x \cdot x) = \left(\frac{d}{dx}x\right)x + x\left(\frac{d}{dx}x\right) = x + x = 2x$$

- Using the previous result, the product rule tells us that

$$\frac{d}{dx}x^3 = \frac{d}{dx}(x \cdot x^2) = \left(\frac{d}{dx}x\right)x^2 + x\left(\frac{d}{dx}x^2\right) = x^2 + x \cdot 2x = 3x^2$$

- Continuing the pattern, we have $\frac{d}{dx}x^n = nx^{n-1}$ for $n = 1, 2, \dots$
- That result is found in a more complicated way in the textbook.

Example: Power Rule for Derivatives

- We can now find derivatives of negative powers of x using the quotient rule.

Example: Power Rule for Derivatives

- We can now find derivatives of negative powers of x using the quotient rule.
- Suppose $f(x) = x^{-m}$ where m is some positive integer.

Example: Power Rule for Derivatives

- We can now find derivatives of negative powers of x using the quotient rule.
- Suppose $f(x) = x^{-m}$ where m is some positive integer.
- We can write $f(x) = 1/x^m$. Differentiating by the quotient rule we have

$$f'(x) = \frac{x^m(1)' - 1(x^m)'}{(x^m)^2} = \frac{-mx^{m-1}}{x^{2m}} = -m\frac{1}{x^{m+1}} = -mx^{-m-1}$$

Example: Power Rule for Derivatives

- We can now find derivatives of negative powers of x using the quotient rule.
- Suppose $f(x) = x^{-m}$ where m is some positive integer.
- We can write $f(x) = 1/x^m$. Differentiating by the quotient rule we have

$$f'(x) = \frac{x^m(1)' - 1(x^m)'}{(x^m)^2} = \frac{-mx^{m-1}}{x^{2m}} = -m\frac{1}{x^{m+1}} = -mx^{-m-1}$$

- Writing $n = -m$ we have $\frac{d}{dx}x^n = nx^{n-1}$ for $n = -1, -2, \dots$

Example: Power Rule for Derivatives

- We can now find derivatives of negative powers of x using the quotient rule.
- Suppose $f(x) = x^{-m}$ where m is some positive integer.
- We can write $f(x) = 1/x^m$. Differentiating by the quotient rule we have

$$f'(x) = \frac{x^m(1)' - 1(x^m)'}{(x^m)^2} = \frac{-mx^{m-1}}{x^{2m}} = -m\frac{1}{x^{m+1}} = -mx^{-m-1}$$

- Writing $n = -m$ we have $\frac{d}{dx}x^n = nx^{n-1}$ for $n = -1, -2, \dots$
- The case $n = 0$ is special, but the result still holds.

Example: Power Rule for Derivatives


- We can now find derivatives of negative powers of x using the quotient rule.
- Suppose $f(x) = x^{-m}$ where m is some positive integer.
- We can write $f(x) = 1/x^m$. Differentiating by the quotient rule we have

$$f'(x) = \frac{x^m(1)' - 1(x^m)'}{(x^m)^2} = \frac{-mx^{m-1}}{x^{2m}} = -m\frac{1}{x^{m+1}} = -mx^{-m-1}$$

- Writing $n = -m$ we have $\frac{d}{dx}x^n = nx^{n-1}$ for $n = -1, -2, \dots$
- The case $n = 0$ is special, but the result still holds.
- In general, $\frac{d}{dx}x^p = px^{p-1}$ for any real number p , but the proof is too complicated to give now.


Tangent Lines Revisited

- Now we can more easily solve derivative questions.




Tangent Lines Revisited

- Now we can more easily solve derivative questions.
- Consider $f(x) = x^2 + 1$. What is the tangent line at $a = 1$?




Tangent Lines Revisited

- Now we can more easily solve derivative questions.
- Consider $f(x) = x^2 + 1$. What is the tangent line at $a = 1$?
- Recall that the tangent line is $y - f(a) = f'(a)(x - a)$.




Tangent Lines Revisited

- Now we can more easily solve derivative questions.
- Consider $f(x) = x^2 + 1$. What is the tangent line at $a = 1$?
- Recall that the tangent line is $y - f(a) = f'(a)(x - a)$.
- We know $f(a)$. From our rules,
$$\frac{d}{dx}f(x) = \frac{d}{dx}(x^2 + 1) = \frac{d}{dx}x^2 + \frac{d}{dx}1 = 2x + 0 = 2x.$$




Tangent Lines Revisited

- Now we can more easily solve derivative questions.
- Consider $f(x) = x^2 + 1$. What is the tangent line at $a = 1$?
- Recall that the tangent line is $y - f(a) = f'(a)(x - a)$.
- We know $f(a)$. From our rules,
$$\frac{d}{dx}f(x) = \frac{d}{dx}(x^2 + 1) = \frac{d}{dx}x^2 + \frac{d}{dx}1 = 2x + 0 = 2x.$$
- So $f'(a) = f'(1) = 2(1) = 2$, and the tangent line is $y - 2 = 2(x - 1)$.




Perpendicular Lines

- We will need to find lines which are **perpendicular** to a tangent line.




Perpendicular Lines

- We will need to find lines which are **perpendicular** to a tangent line.
- Two lines are perpendicular if the product of their slopes is -1 .




Perpendicular Lines

- We will need to find lines which are **perpendicular** to a tangent line.
- Two lines are perpendicular if the product of their slopes is -1 .
- E.g., if we have two lines $y = m_1x + b$ and $y - y_0 = m_2(x - x_0)$, they are perpendicular if $m_1m_2 = -1$.




Perpendicular Lines

- We will need to find lines which are **perpendicular** to a tangent line.
- Two lines are perpendicular if the product of their slopes is -1 .
- E.g., if we have two lines $y = m_1x + b$ and $y - y_0 = m_2(x - x_0)$, they are perpendicular if $m_1m_2 = -1$.
- Given the slope m_1 of a line, the slope m_2 of any perpendicular line is the **negative reciprocal** of m_1 : $m_2 = -\frac{1}{m_1}$




Normal Lines

- Now we can find **normal lines** (lines perpendicular to a tangent at a point on a curve). Let $f(x) = x^2 + 1$ and $a = 1$.




Normal Lines

- Now we can find **normal lines** (lines perpendicular to a tangent at a point on a curve). Let $f(x) = x^2 + 1$ and $a = 1$.
- The slope of the tangent is $f'(1) = 2(1) = 2$; the slope of the normal line is $m_2 = -1/2$.




Normal Lines

- Now we can find **normal lines** (lines perpendicular to a tangent at a point on a curve). Let $f(x) = x^2 + 1$ and $a = 1$.
- The slope of the tangent is $f'(1) = 2(1) = 2$; the slope of the normal line is $m_2 = -1/2$.
- The normal line in point-slope form is $y - f(a) = m_2(x - a)$, or filling in the blanks,




Normal Lines

- Now we can find **normal lines** (lines perpendicular to a tangent at a point on a curve). Let $f(x) = x^2 + 1$ and $a = 1$.
- The slope of the tangent is $f'(1) = 2(1) = 2$; the slope of the normal line is $m_2 = -1/2$.
- The normal line in point-slope form is $y - f(a) = m_2(x - a)$, or filling in the blanks, $y - 2 = -\frac{1}{2}(x - 1)$.



Normal Lines

- Now we can find **normal lines** (lines perpendicular to a tangent at a point on a curve). Let $f(x) = x^2 + 1$ and $a = 1$.
- The slope of the tangent is $f'(1) = 2(1) = 2$; the slope of the normal line is $m_2 = -1/2$.
- The normal line in point-slope form is $y - f(a) = m_2(x - a)$, or filling in the blanks, $y - 2 = -\frac{1}{2}(x - 1)$.
- The normal line to $y = f(x)$ at $P(a, f(a))$ is $y - f(a) = -\frac{1}{f'(a)}(x - a)$.



Examples

1. Calculate y' where

1.1 $y = \sqrt{x} + \frac{1}{\sqrt[3]{x^4}}$

1.2 $y = \frac{3x-2}{\sqrt{x}}.$

1.3 $y = \frac{\sqrt{x}}{3x-2}.$

2. Find the equations of the tangent and normal lines to the curve

$y = \frac{x-1}{x+1}$ at the point $(0, -1)$.

3. Find the equations of the tangent and normal lines to the curve

$y = \frac{x^2-1}{x^2+1}$ at the point $(0, -1)$.

Solution to Example 3

- In this example we have

$$y' = \frac{(x^2 + 1)2x - (x^2 - 1)2x}{(x^2 + 1)^2} = \frac{4x}{(x^2 + 1)^2}, \text{ so } y'(0) = 0.$$

- The equation of the tangent line is easy enough to find:
 $y - f(a) = f'(a)(x - a)$ where $a = 0$, $f(a) = -1$, and $f'(a) = 0$
gives $y - (-1) = 0(x - 0)$ or in other words $y = -1$.
- The equation of the normal is not so clear in this case. Since the slope of the tangent is 0, the slope of the normal would be $-1/0$ which doesn't make any sense. However, it should seem clear that a line perpendicular to the horizontal line $y = -1$ is a vertical line. Since the vertical line must pass through $(0, 1)$, the equation of the normal line is $x = 0$. Draw it!

Now you should work on Problem Set 2.3. After you have finished it, you should try the following additional exercises from Section 2.3:

2.3 C-level: 1–22, 25–44, 51–52, 53–54, 55–58, 59–62, 63–66

B-level: 23–24, 45, 46–48, 49–50, 67, 68, 69–72, 73–74, 75–76,
77–82, 83–86, 93–96,

A-level: 87, 88–92, 97–100, 101–110