

# MATH 110 Review 0.1

## Review of Algebra

---

Edward Doolittle

Tuesday, January 6, 2026

Department of Indigenous Knowledge and Science  
First Nations University of Canada

## Review of Algebra

Arithmetic Operations

Factoring and Quadratics

Exponents and the Binomial Theorem

# Review of Algebra

---

# Commutativity of Addition

- If I add two numbers, say 2 and 3, the answer does not depend on the order in which I add the numbers.

# Commutativity of Addition

- If I add two numbers, say 2 and 3, the answer does not depend on the order in which I add the numbers.
- $2 + 3 = 5$ , but also  $3 + 2 = 5$ .

# Commutativity of Addition

- If I add two numbers, say 2 and 3, the answer does not depend on the order in which I add the numbers.
- $2 + 3 = 5$ , but also  $3 + 2 = 5$ .
- In other words we can write  $2 + 3 = 3 + 2$ .

# Commutativity of Addition

- If I add two numbers, say 2 and 3, the answer does not depend on the order in which I add the numbers.
- $2 + 3 = 5$ , but also  $3 + 2 = 5$ .
- In other words we can write  $2 + 3 = 3 + 2$ .
- The above result doesn't depend on the numbers 2 and 3. For example, I also have  $4 + 7 = 7 + 4$ , and so on.

# Commutativity of Addition

- If I add two numbers, say 2 and 3, the answer does not depend on the order in which I add the numbers.
- $2 + 3 = 5$ , but also  $3 + 2 = 5$ .
- In other words we can write  $2 + 3 = 3 + 2$ .
- The above result doesn't depend on the numbers 2 and 3. For example, I also have  $4 + 7 = 7 + 4$ , and so on.
- We say that the operation of addition is *commutative*.



# Commutative Law for Addition

- We can express that more general result using *algebraic* notation:  $a + b = b + a$  for any numbers  $a$  and  $b$ .

# Commutative Law for Addition

- We can express that more general result using *algebraic* notation:  $a + b = b + a$  for any numbers  $a$  and  $b$ .
- For example choosing  $a = 2$  and  $b = 3$  gives us the result  $2 + 3 = 3 + 2$ , and choosing  $a = 4$  and  $b = 7$  gives us the result  $4 + 7 = 7 + 4$ .

# Commutative Law for Addition

- We can express that more general result using *algebraic* notation:  $a + b = b + a$  for any numbers  $a$  and  $b$ .
- For example choosing  $a = 2$  and  $b = 3$  gives us the result  $2 + 3 = 3 + 2$ , and choosing  $a = 4$  and  $b = 7$  gives us the result  $4 + 7 = 7 + 4$ .
- The notation  $a + b = b + a$  is very powerful; it stands for an infinite number of different particular results.

# Commutative Law for Addition

- We can express that more general result using *algebraic* notation:  $a + b = b + a$  for any numbers  $a$  and  $b$ .
- For example choosing  $a = 2$  and  $b = 3$  gives us the result  $2 + 3 = 3 + 2$ , and choosing  $a = 4$  and  $b = 7$  gives us the result  $4 + 7 = 7 + 4$ .
- The notation  $a + b = b + a$  is very powerful; it stands for an infinite number of different particular results.
- We call that expression the *commutative law for addition*.

# Commutativity of Multiplication

- If I multiply two numbers, say 3 and 5, the answer does not depend on the order in which I multiply the numbers.

# Commutativity of Multiplication

- If I multiply two numbers, say 3 and 5, the answer does not depend on the order in which I multiply the numbers.
- We have  $3 \times 5 = 15$ , but we also have  $5 \times 3 = 15$

# Commutativity of Multiplication

- If I multiply two numbers, say 3 and 5, the answer does not depend on the order in which I multiply the numbers.
- We have  $3 \times 5 = 15$ , but we also have  $5 \times 3 = 15$
- In other words, we can write  $5 \times 3 = 3 \times 5$ .

# Commutativity of Multiplication

- If I multiply two numbers, say 3 and 5, the answer does not depend on the order in which I multiply the numbers.
- We have  $3 \times 5 = 15$ , but we also have  $5 \times 3 = 15$
- In other words, we can write  $5 \times 3 = 3 \times 5$ .
- The above result does not depend on the particular numbers 3 and 5. For example, we have  $2 \times 6 = 6 \times 2$ , and so on, for any two numbers I choose.



# Commutativity of Multiplication

- If I multiply two numbers, say 3 and 5, the answer does not depend on the order in which I multiply the numbers.
- We have  $3 \times 5 = 15$ , but we also have  $5 \times 3 = 15$
- In other words, we can write  $5 \times 3 = 3 \times 5$ .
- The above result does not depend on the particular numbers 3 and 5. For example, we have  $2 \times 6 = 6 \times 2$ , and so on, for any two numbers I choose.
- We say that the operation of multiplication is also commutative.

# Commutative Law for Multiplication

- We can express that more general result using *algebraic* notation:  $a \times b = b \times a$  for any numbers  $a$  and  $b$ .

# Commutative Law for Multiplication

- We can express that more general result using *algebraic* notation:  $a \times b = b \times a$  for any numbers  $a$  and  $b$ .
- As a short form, we usually drop the  $\times$  if we can do so without confusion, so we usually write  $ab = ba$ .

# Commutative Law for Multiplication

- We can express that more general result using *algebraic* notation:  $a \times b = b \times a$  for any numbers  $a$  and  $b$ .
- As a short form, we usually drop the  $\times$  if we can do so without confusion, so we usually write  $ab = ba$ .
- For example choosing  $a = 3$  and  $b = 5$  gives us the result  $3 \times 5 = 5 \times 3$ , and choosing  $a = 2$  and  $b = 6$  gives us the result  $2 \times 6 = 6 \times 2$ .

# Commutative Law for Multiplication

- We can express that more general result using *algebraic* notation:  $a \times b = b \times a$  for any numbers  $a$  and  $b$ .
- As a short form, we usually drop the  $\times$  if we can do so without confusion, so we usually write  $ab = ba$ .
- For example choosing  $a = 3$  and  $b = 5$  gives us the result  $3 \times 5 = 5 \times 3$ , and choosing  $a = 2$  and  $b = 6$  gives us the result  $2 \times 6 = 6 \times 2$ .
- The notation  $ab = ba$  is very powerful; it stands for an infinite number of different particular results.

# Commutative Law for Multiplication

- We can express that more general result using *algebraic* notation:  $a \times b = b \times a$  for any numbers  $a$  and  $b$ .
- As a short form, we usually drop the  $\times$  if we can do so without confusion, so we usually write  $ab = ba$ .
- For example choosing  $a = 3$  and  $b = 5$  gives us the result  $3 \times 5 = 5 \times 3$ , and choosing  $a = 2$  and  $b = 6$  gives us the result  $2 \times 6 = 6 \times 2$ .
- The notation  $ab = ba$  is very powerful; it stands for an infinite number of different particular results.
- We call that expression the *commutative law for multiplication*.

# The Commutative Laws

- In summary, we have the commutative law of addition:

$$a + b = b + a$$

for any numbers  $a$  and  $b$

# The Commutative Laws

- In summary, we have the commutative law of addition:

$$a + b = b + a$$

for any numbers  $a$  and  $b$

- And the commutative law of multiplication

$$ab = ba$$



# The Commutative Laws

- In summary, we have the commutative law of addition:

$$a + b = b + a$$

for any numbers  $a$  and  $b$

- And the commutative law of multiplication

$$ab = ba$$

- We obviously don't have to use the letters  $a$  and  $b$ . So we could write the *commutative laws* as

$$x + y = y + x$$

and

$$xy = yx$$

# The Commutative Laws

- In summary, we have the commutative law of addition:

$$a + b = b + a$$

for any numbers  $a$  and  $b$

- And the commutative law of multiplication

$$ab = ba$$

- We obviously don't have to use the letters  $a$  and  $b$ . So we could write the *commutative laws* as

$$x + y = y + x$$

and

$$xy = yx$$

- That is our first result in algebra.

# Associative Laws

- Similarly, the order in which we add three numbers does not matter.

# Associative Laws

- Similarly, the order in which we add three numbers does not matter.
- For example,  $(3 + 6) + 5 = 3 + (6 + 5)$ .

# Associative Laws

- Similarly, the order in which we add three numbers does not matter.
- For example,  $(3 + 6) + 5 = 3 + (6 + 5)$ .
- In general, we write  $(x + y) + z = x + (y + z)$ .

# Associative Laws

- Similarly, the order in which we add three numbers does not matter.
- For example,  $(3 + 6) + 5 = 3 + (6 + 5)$ .
- In general, we write  $(x + y) + z = x + (y + z)$ .
- Furthermore, the order in which we multiply three numbers does not matter.

# Associative Laws

- Similarly, the order in which we add three numbers does not matter.
- For example,  $(3 + 6) + 5 = 3 + (6 + 5)$ .
- In general, we write  $(x + y) + z = x + (y + z)$ .
- Furthermore, the order in which we multiply three numbers does not matter.
- For example,  $(2 \times 5) \times 3 = 2 \times (5 \times 3)$ .

# Associative Laws

- Similarly, the order in which we add three numbers does not matter.
- For example,  $(3 + 6) + 5 = 3 + (6 + 5)$ .
- In general, we write  $(x + y) + z = x + (y + z)$ .
- Furthermore, the order in which we multiply three numbers does not matter.
- For example,  $(2 \times 5) \times 3 = 2 \times (5 \times 3)$ .
- In general, we write  $(x \times y) \times z = x \times (y \times z)$ , or in short form  $(xy)z = x(yz)$ .



# Distributive Laws

- The previous laws applied to just one operation, addition or multiplication.

## Distributive Laws

- The previous laws applied to just one operation, addition or multiplication.
- We also have a law which involves both operations simultaneously, the distributive law.

## Distributive Laws

- The previous laws applied to just one operation, addition or multiplication.
- We also have a law which involves both operations simultaneously, the distributive law.
- Adding first and then multiplying is the same as multiplying first and then adding.

## Distributive Laws

- The previous laws applied to just one operation, addition or multiplication.
- We also have a law which involves both operations simultaneously, the distributive law.
- Adding first and then multiplying is the same as multiplying first and then adding.
- More precisely, in symbols, we write  $x(y + z) = (xy) + (xz)$ .

# Distributive Laws

- The previous laws applied to just one operation, addition or multiplication.
- We also have a law which involves both operations simultaneously, the distributive law.
- Adding first and then multiplying is the same as multiplying first and then adding.
- More precisely, in symbols, we write  $x(y + z) = (xy) + (xz)$ .
- Order of operations (BEDMAS) says that multiplication is done before addition so we can drop the brackets on the right hand side (RHS).

# Distributive Laws

- The previous laws applied to just one operation, addition or multiplication.
- We also have a law which involves both operations simultaneously, the distributive law.
- Adding first and then multiplying is the same as multiplying first and then adding.
- More precisely, in symbols, we write  $x(y + z) = (xy) + (xz)$ .
- Order of operations (BEDMAS) says that multiplication is done before addition so we can drop the brackets on the right hand side (RHS).
- The *distributive law* is usually written  $x(y + z) = xy + xz$ .

- The distributive law is the key to an important process in algebra known as *expanding*.

# Expanding

- The distributive law is the key to an important process in algebra known as *expanding*.
- Consider the expression  $3(t + 2)$ .



# Expanding

- The distributive law is the key to an important process in algebra known as *expanding*.
- Consider the expression  $3(t + 2)$ .
- Expanding means applying the distributive law to remove the brackets:  $3(t + 2) = 3 \times t + 3 \times 2 = 3t + 6$ .

# Expanding

- The distributive law is the key to an important process in algebra known as *expanding*.
- Consider the expression  $3(t + 2)$ .
- Expanding means applying the distributive law to remove the brackets:  $3(t + 2) = 3 \times t + 3 \times 2 = 3t + 6$ .
- Now consider  $(s + 3)(t + 2)$ .

# Expanding

- The distributive law is the key to an important process in algebra known as *expanding*.
- Consider the expression  $3(t + 2)$ .
- Expanding means applying the distributive law to remove the brackets:  $3(t + 2) = 3 \times t + 3 \times 2 = 3t + 6$ .
- Now consider  $(s + 3)(t + 2)$ .
- Apply the distributive law:  $(s + 3)(t + 2) = (s + 3)t + (s + 3)2$ .

# Expanding

- The distributive law is the key to an important process in algebra known as *expanding*.
- Consider the expression  $3(t + 2)$ .
- Expanding means applying the distributive law to remove the brackets:  $3(t + 2) = 3 \times t + 3 \times 2 = 3t + 6$ .
- Now consider  $(s + 3)(t + 2)$ .
- Apply the distributive law:  $(s + 3)(t + 2) = (s + 3)t + (s + 3)2$ .
- Apply it again:  $(s + 3)t + (s + 3)2 = st + 3t + (s + 3)2$ .

# Expanding

- The distributive law is the key to an important process in algebra known as *expanding*.
- Consider the expression  $3(t + 2)$ .
- Expanding means applying the distributive law to remove the brackets:  $3(t + 2) = 3 \times t + 3 \times 2 = 3t + 6$ .
- Now consider  $(s + 3)(t + 2)$ .
- Apply the distributive law:  $(s + 3)(t + 2) = (s + 3)t + (s + 3)2$ .
- Apply it again:  $(s + 3)t + (s + 3)2 = st + 3t + (s + 3)2$ .
- And again:  $st + 3t + (s + 3)2 = st + 3t + 2s + 6$ .

- We can apply any of the laws we know to simplify expressions.

# Simplifying

- We can apply any of the laws we know to simplify expressions.
- Consider  $4 - 3x + 6$ .

# Simplifying

- We can apply any of the laws we know to simplify expressions.
- Consider  $4 - 3x + 6$ .
- We can write  $4 - 3x + 6 = 4 + (-3 \times x) + 6$ .



# Simplifying

- We can apply any of the laws we know to simplify expressions.
- Consider  $4 - 3x + 6$ .
- We can write  $4 - 3x + 6 = 4 + (-3 \times x) + 6$ .
- Applying the commutative law of addition, we have  
 $4 + (-3 \times x) + 6 = 4 + 6 + (-3 \times x)$ .

# Simplifying

- We can apply any of the laws we know to simplify expressions.
- Consider  $4 - 3x + 6$ .
- We can write  $4 - 3x + 6 = 4 + (-3 \times x) + 6$ .
- Applying the commutative law of addition, we have  
 $4 + (-3 \times x) + 6 = 4 + 6 + (-3 \times x)$ .
- We can add the 4 and 6 to get 10:  
 $4 + 6 + (-3 \times x) = 10 + (-3 \times x)$ .

# Simplifying

- We can apply any of the laws we know to simplify expressions.
- Consider  $4 - 3x + 6$ .
- We can write  $4 - 3x + 6 = 4 + (-3 \times x) + 6$ .
- Applying the commutative law of addition, we have  
 $4 + (-3 \times x) + 6 = 4 + 6 + (-3 \times x)$ .
- We can add the 4 and 6 to get 10:  
 $4 + 6 + (-3 \times x) = 10 + (-3 \times x)$ .
- By Order of Operations we can drop the brackets:  
 $10 + (-3 \times x) = 10 - 3x$ .

# Simplifying

- We can apply any of the laws we know to simplify expressions.
- Consider  $4 - 3x + 6$ .
- We can write  $4 - 3x + 6 = 4 + (-3 \times x) + 6$ .
- Applying the commutative law of addition, we have  
 $4 + (-3 \times x) + 6 = 4 + 6 + (-3 \times x)$ .
- We can add the 4 and 6 to get 10:  
 $4 + 6 + (-3 \times x) = 10 + (-3 \times x)$ .
- By Order of Operations we can drop the brackets:  
 $10 + (-3 \times x) = 10 - 3x$ .
- In summary,  $4 - 3x + 6 = 10 - 3x$ .

# Multiplying Fractions

- We also need to remember rules for dealing with fractions.

# Multiplying Fractions

- We also need to remember rules for dealing with fractions.
- The simplest operation we can perform on fractions is multiplying them.

# Multiplying Fractions

- We also need to remember rules for dealing with fractions.
- The simplest operation we can perform on fractions is multiplying them.
- The rule for multiplying fractions is

$$\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}$$

# Multiplying Fractions

- We also need to remember rules for dealing with fractions.
- The simplest operation we can perform on fractions is multiplying them.
- The rule for multiplying fractions is

$$\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}$$

- We just *multiply the numerators and multiply the denominators* to get the answer.



# Dividing Fractions

- The next simplest operation for fractions is division.

# Dividing Fractions

- The next simplest operation for fractions is division.
- To divide fractions we *invert and multiply*:

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}$$

# Dividing Fractions

- The next simplest operation for fractions is division.
- To divide fractions we *invert and multiply*:

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}$$

- Sometimes we see the same procedure in slightly different notation:

$$\frac{a/b}{c/d} = \frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}$$

# Negating Fractions

- Negating a fraction is the same thing as multiplying it by

$$-1 = \frac{-1}{1} = \frac{1}{-1}$$

# Negating Fractions

- Negating a fraction is the same thing as multiplying it by

$$-1 = \frac{-1}{1} = \frac{1}{-1}$$

- So all three of the following are negatives of  $2/3$ :

$$-\frac{2}{3} = \frac{-2}{3} = \frac{2}{-3}$$

# Equivalent Fractions

- There are many different ways of representing the number 1 as a fraction:

$$1 = \frac{1}{1} = \frac{2}{2} = \frac{7}{7} = \frac{-3}{-3} = \dots$$

# Equivalent Fractions

- There are many different ways of representing the number 1 as a fraction:

$$1 = \frac{1}{1} = \frac{2}{2} = \frac{7}{7} = \frac{-3}{-3} = \dots$$

- Multiplying any fraction by any of those representations of 1 gives an equivalent fraction:

$$\frac{2}{3} = \frac{4}{6} = \frac{14}{21} = \frac{-6}{-9} = \dots$$

# Adding Fractions I

- The rule for adding fractions is surprisingly more complicated than the rule for multiplying fractions.



# Adding Fractions I

- The rule for adding fractions is surprisingly more complicated than the rule for multiplying fractions.
- To add fractions, we must find a common denominator.

# Adding Fractions I

- The rule for adding fractions is surprisingly more complicated than the rule for multiplying fractions.
- To add fractions, we must find a common denominator.
- We do that by finding appropriate equivalent fractions.

# Adding Fractions I

- The rule for adding fractions is surprisingly more complicated than the rule for multiplying fractions.
- To add fractions, we must find a common denominator.
- We do that by finding appropriate equivalent fractions.
- For example, to add  $2/3$  and  $4/7$ , we find fractions equivalent to  $2/3$ , and fractions equivalent to  $4/7$ , looking for a common denominator.

$$\frac{2}{3} = \frac{4}{6} = \cdots = \frac{14}{21} = \cdots$$

$$\frac{4}{7} = \frac{8}{14} = \frac{12}{21} = \cdots$$

# Adding Fractions I

- The rule for adding fractions is surprisingly more complicated than the rule for multiplying fractions.
- To add fractions, we must find a common denominator.
- We do that by finding appropriate equivalent fractions.
- For example, to add  $2/3$  and  $4/7$ , we find fractions equivalent to  $2/3$ , and fractions equivalent to  $4/7$ , looking for a common denominator.

$$\frac{2}{3} = \frac{4}{6} = \cdots = \frac{14}{21} = \cdots$$

$$\frac{4}{7} = \frac{8}{14} = \frac{12}{21} = \cdots$$

- We found a common denominator, 21.

## Adding Fractions II

- Now we can add the two fractions:

$$\frac{2}{3} + \frac{4}{7} = \frac{14}{21} + \frac{12}{21} = \frac{26}{21}$$

## Adding Fractions II

- Now we can add the two fractions:

$$\frac{2}{3} + \frac{4}{7} = \frac{14}{21} + \frac{12}{21} = \frac{26}{21}$$

- Note that, with addition, we just add the numerators and we leave the denominators alone.

## Adding Fractions II

- Now we can add the two fractions:

$$\frac{2}{3} + \frac{4}{7} = \frac{14}{21} + \frac{12}{21} = \frac{26}{21}$$

- Note that, with addition, we just add the numerators and we leave the denominators alone.
- Note that we can always find a common denominator:

$$\frac{a}{b} + \frac{c}{d} = \frac{ad}{bd} + \frac{bc}{bd} = \frac{ad + bc}{bd}$$

## Adding Fractions II

- Now we can add the two fractions:

$$\frac{2}{3} + \frac{4}{7} = \frac{14}{21} + \frac{12}{21} = \frac{26}{21}$$

- Note that, with addition, we just add the numerators and we leave the denominators alone.
- Note that we can always find a common denominator:

$$\frac{a}{b} + \frac{c}{d} = \frac{ad}{bd} + \frac{bc}{bd} = \frac{ad + bc}{bd}$$

- That is a general formula for adding fractions.



# Subtracting Fractions

- We follow a similar procedure to subtract fractions.

# Subtracting Fractions

- We follow a similar procedure to subtract fractions.
- We find a common denominator and then subtract only the numerators:

$$\frac{a}{b} - \frac{c}{d} = \frac{ad}{bd} - \frac{bc}{bd} = \frac{ad - bc}{bd}$$

# Subtracting Fractions

- We follow a similar procedure to subtract fractions.
- We find a common denominator and then subtract only the numerators:

$$\frac{a}{b} - \frac{c}{d} = \frac{ad}{bd} - \frac{bc}{bd} = \frac{ad - bc}{bd}$$

- It's better to remember the procedure for adding and subtracting fractions, rather than the formula.

## Expanding and Factoring

- The opposite of *expanding* is *factoring*.

# Expanding and Factoring

- The opposite of *expanding* is *factoring*.
- If I start with an expression like  $2x(x + 3)$  I can expand it to get

# Expanding and Factoring

- The opposite of *expanding* is *factoring*.
- If I start with an expression like  $2x(x + 3)$  I can expand it to get  $2x^2 + 6x$
- Sometimes expanding is appropriate, and sometimes factoring is appropriate.

## Expanding and Factoring

- The opposite of *expanding* is *factoring*.
- If I start with an expression like  $2x(x + 3)$  I can expand it to get  $2x^2 + 6x$
- On the other hand, if I start with  $2x^2 + 6x$  I can factor it to get  $2x(x + 3)$ .
- Sometimes expanding is appropriate, and sometimes factoring is appropriate.
- It's hard to give a rule to tell you which is better in a given situation. You need to use your judgment.

# Factoring Quadratics

- Many of the examples used in our course will require factoring quadratics.



# Factoring Quadratics

- Many of the examples used in our course will require factoring quadratics.
- A quadratic is a polynomial with highest degree term of degree 2, for example  $x^2 + 5x + 6$  or  $x^2 - 9$  or  $-3x^2 + 6x - 3$ .

# Factoring Quadratics

- Many of the examples used in our course will require factoring quadratics.
- A quadratic is a polynomial with highest degree term of degree 2, for example  $x^2 + 5x + 6$  or  $x^2 - 9$  or  $-3x^2 + 6x - 3$ .
- Often you can guess how to factor a quadratic of the form  $x^2 + ax + b$ .

# Factoring Quadratics

- Many of the examples used in our course will require factoring quadratics.
- A quadratic is a polynomial with highest degree term of degree 2, for example  $x^2 + 5x + 6$  or  $x^2 - 9$  or  $-3x^2 + 6x - 3$ .
- Often you can guess how to factor a quadratic of the form  $x^2 + ax + b$ .
- You need to find two numbers that add to give  $a$  and multiply to give  $b$ .

# Factoring Quadratics

- Many of the examples used in our course will require factoring quadratics.
- A quadratic is a polynomial with highest degree term of degree 2, for example  $x^2 + 5x + 6$  or  $x^2 - 9$  or  $-3x^2 + 6x - 3$ .
- Often you can guess how to factor a quadratic of the form  $x^2 + ax + b$ .
- You need to find two numbers that add to give  $a$  and multiply to give  $b$ .
- For the example  $x^2 + 5x + 6$ , you would eventually guess 2 and 3: their sum is 5 and their product is 6.

# Factoring Quadratics

- Many of the examples used in our course will require factoring quadratics.
- A quadratic is a polynomial with highest degree term of degree 2, for example  $x^2 + 5x + 6$  or  $x^2 - 9$  or  $-3x^2 + 6x - 3$ .
- Often you can guess how to factor a quadratic of the form  $x^2 + ax + b$ .
- You need to find two numbers that add to give  $a$  and multiply to give  $b$ .
- For the example  $x^2 + 5x + 6$ , you would eventually guess 2 and 3: their sum is 5 and their product is 6.
- So we guess  $x^2 + 5x + 6 = (x + 2)(x + 3)$ .

# Factoring Quadratics

- Many of the examples used in our course will require factoring quadratics.
- A quadratic is a polynomial with highest degree term of degree 2, for example  $x^2 + 5x + 6$  or  $x^2 - 9$  or  $-3x^2 + 6x - 3$ .
- Often you can guess how to factor a quadratic of the form  $x^2 + ax + b$ .
- You need to find two numbers that add to give  $a$  and multiply to give  $b$ .
- For the example  $x^2 + 5x + 6$ , you would eventually guess 2 and 3: their sum is 5 and their product is 6.
- So we guess  $x^2 + 5x + 6 = (x + 2)(x + 3)$ .
- Check by expanding the RHS.

# Difference of Squares

- One common case of quadratic polynomial is a difference of squares.

# Difference of Squares

- One common case of quadratic polynomial is a difference of squares.
- For example,  $x^2$  is a square and  $9 = 3^2$  is a square, and their difference is  $x^2 - 9$ .



# Difference of Squares

- One common case of quadratic polynomial is a difference of squares.
- For example,  $x^2$  is a square and  $9 = 3^2$  is a square, and their difference is  $x^2 - 9$ .
- A difference of squares can always be factored by taking the sum and difference of the square roots of the squares.

# Difference of Squares

- One common case of quadratic polynomial is a difference of squares.
- For example,  $x^2$  is a square and  $9 = 3^2$  is a square, and their difference is  $x^2 - 9$ .
- A difference of squares can always be factored by taking the sum and difference of the square roots of the squares.
- For example,  $x^2 - 9 = (x + 3)(x - 3)$ .

# Difference of Squares

- One common case of quadratic polynomial is a difference of squares.
- For example,  $x^2$  is a square and  $9 = 3^2$  is a square, and their difference is  $x^2 - 9$ .
- A difference of squares can always be factored by taking the sum and difference of the square roots of the squares.
- For example,  $x^2 - 9 = (x + 3)(x - 3)$ .
- Again, check by expanding the RHS.

## Difference of Cubes

- There is also a similar pattern for factoring a difference of cubes.

## Difference of Cubes

- There is also a similar pattern for factoring a difference of cubes.
- A difference of cubes is a polynomial of the form  $x^3 - a^3$ , for example,  $x^3 - 8 = x^3 - 2^3$ .

## Difference of Cubes

- There is also a similar pattern for factoring a difference of cubes.
- A difference of cubes is a polynomial of the form  $x^3 - a^3$ , for example,  $x^3 - 8 = x^3 - 2^3$ .
- The pattern is  $x^3 - a^3 = (x - a)(x^2 + ax + a^2)$ .

## Difference of Cubes

- There is also a similar pattern for factoring a difference of cubes.
- A difference of cubes is a polynomial of the form  $x^3 - a^3$ , for example,  $x^3 - 8 = x^3 - 2^3$ .
- The pattern is  $x^3 - a^3 = (x - a)(x^2 + ax + a^2)$ .
- We can check by expanding the RHS:

$$(x - a)(x^2 + ax + a^2) = x^3 + ax^2 + a^2x - ax^2 - a^2x - a^3 = x^3 - a^3$$

## Difference of Cubes

- There is also a similar pattern for factoring a difference of cubes.
- A difference of cubes is a polynomial of the form  $x^3 - a^3$ , for example,  $x^3 - 8 = x^3 - 2^3$ .
- The pattern is  $x^3 - a^3 = (x - a)(x^2 + ax + a^2)$ .
- We can check by expanding the RHS:

$$(x - a)(x^2 + ax + a^2) = x^3 + ax^2 + a^2x - ax^2 - a^2x - a^3 = x^3 - a^3$$

- For example,  $x^3 - 8 = (x - 2)(x^2 + 2x + 4)$ . (Check!)



## Sum of Cubes

- We can't factor a sum of squares like  $x^2 + 9$

## Sum of Cubes

- We can't factor a sum of squares like  $x^2 + 9$
- However, we can factor a sum of cubes using a trick.

## Sum of Cubes

- We can't factor a sum of squares like  $x^2 + 9$
- However, we can factor a sum of cubes using a trick.
- The trick is based on this observation:  $b^3 = -(-b^3) = -(-b)^3$

## Sum of Cubes

- We can't factor a sum of squares like  $x^2 + 9$
- However, we can factor a sum of cubes using a trick.
- The trick is based on this observation:  $b^3 = -(-b^3) = -(-b)^3$
- So a sum of cubes can be written as a difference of cubes:

$$x^3 + b^3 = x^3 - (-b)^3 = x^3 - a^3$$

where  $a = -b$ .

## Sum of Cubes

- We can't factor a sum of squares like  $x^2 + 9$
- However, we can factor a sum of cubes using a trick.
- The trick is based on this observation:  $b^3 = -(-b^3) = -(-b)^3$
- So a sum of cubes can be written as a difference of cubes:

$$x^3 + b^3 = x^3 - (-b)^3 = x^3 - a^3$$

where  $a = -b$ .

- Using our difference of cubes formula, we have  
$$x^3 - a^3 = (x - a)(x^2 + ax + a^2)$$

## Sum of Cubes

- We can't factor a sum of squares like  $x^2 + 9$
- However, we can factor a sum of cubes using a trick.
- The trick is based on this observation:  $b^3 = -(-b^3) = -(-b)^3$
- So a sum of cubes can be written as a difference of cubes:

$$x^3 + b^3 = x^3 - (-b)^3 = x^3 - a^3$$

where  $a = -b$ .

- Using our difference of cubes formula, we have
$$x^3 - a^3 = (x - a)(x^2 + ax + a^2)$$
- Putting back  $-b$  for  $a$  we have  $x^3 + b^3 = (x + b)(x^2 - bx + b^2)$ .  
Check!

# The Factor Theorem

- There is a connection between the roots of a polynomial and its factors.

# The Factor Theorem

- There is a connection between the roots of a polynomial and its factors.
- Note that 2 is a root of the polynomial  $x^2 - 5x + 6$ , i.e., when we substitute 2 for  $x$ , the result is 0.



# The Factor Theorem

- There is a connection between the roots of a polynomial and its factors.
- Note that 2 is a root of the polynomial  $x^2 - 5x + 6$ , i.e., when we substitute 2 for  $x$ , the result is 0.
- $2^2 - 5(2) + 6 = 4 - 10 + 6 = 0$

# The Factor Theorem

- There is a connection between the roots of a polynomial and its factors.
- Note that 2 is a root of the polynomial  $x^2 - 5x + 6$ , i.e., when we substitute 2 for  $x$ , the result is 0.
- $2^2 - 5(2) + 6 = 4 - 10 + 6 = 0$
- That means that  $x - 2$  should be a factor of  $x^2 - 5x + 6$ .

# The Factor Theorem

- There is a connection between the roots of a polynomial and its factors.
- Note that 2 is a root of the polynomial  $x^2 - 5x + 6$ , i.e., when we substitute 2 for  $x$ , the result is 0.
- $2^2 - 5(2) + 6 = 4 - 10 + 6 = 0$
- That means that  $x - 2$  should be a factor of  $x^2 - 5x + 6$ .
- Indeed it is: check  $x^2 - 5x + 6 = (x - 2)(x - 3)$

- If we know one factor of a number, we can find the other factor by division.

# Polynomial Division I

- If we know one factor of a number, we can find the other factor by division.
- For example, we know that 2 is a factor of 26, we can find the other factor by dividing:  $26/2 = 13$ .

# Polynomial Division I

- If we know one factor of a number, we can find the other factor by division.
- For example, we know that 2 is a factor of 26, we can find the other factor by dividing:  $26/2 = 13$ .
- We write  $26 = 2 \times 13$

# Polynomial Division I

- If we know one factor of a number, we can find the other factor by division.
- For example, we know that 2 is a factor of 26, we can find the other factor by dividing:  $26/2 = 13$ .
- We write  $26 = 2 \times 13$
- We can do the same with polynomials.

## Polynomial Division II

- I recommend you divide polynomials as follows. First Write down the form of the factorization:  $x^2 - 5x + 6 = (x - 2)(x + a)$



## Polynomial Division II

- I recommend you divide polynomials as follows. First Write down the form of the factorization:  $x^2 - 5x + 6 = (x - 2)(x + a)$
- Expand and gather the RHS:  
$$(x - 2)(x + a) = x^2 - 2x + ax - 2a = x^2 + (a - 2)x + (-2a)$$

## Polynomial Division II

- I recommend you divide polynomials as follows. First Write down the form of the factorization:  $x^2 - 5x + 6 = (x - 2)(x + a)$
- Expand and gather the RHS:  
$$(x - 2)(x + a) = x^2 - 2x + ax - 2a = x^2 + (a - 2)x + (-2a)$$
- Compare:  $x^2 - 5x + 6 = x^2 + (a - 2)x + (-2a)$  means  $a - 2 = -5$   
means  $a = -3$ , so the other factor should be  $x + (-3) = x - 3$

## Polynomial Division II

- I recommend you divide polynomials as follows. First Write down the form of the factorization:  $x^2 - 5x + 6 = (x - 2)(x + a)$
- Expand and gather the RHS:  
$$(x - 2)(x + a) = x^2 - 2x + ax - 2a = x^2 + (a - 2)x + (-2a)$$
- Compare:  $x^2 - 5x + 6 = x^2 + (a - 2)x + (-2a)$  means  $a - 2 = -5$   
means  $a = -3$ , so the other factor should be  $x + (-3) = x - 3$
- Check that  $x^2 - 5x + 6 = (x - 2)(x - 3)$

# Completing the Square I

- An important technique for dealing with quadratic polynomials is *completing the square*.

# Completing the Square I

- An important technique for dealing with quadratic polynomials is *completing the square*.
- Consider the polynomial  $x^2 + 6x + 5$ .

# Completing the Square I

- An important technique for dealing with quadratic polynomials is *completing the square*.
- Consider the polynomial  $x^2 + 6x + 5$ .
- We want to find a square that looks as much like that polynomial as possible.

# Completing the Square I

- An important technique for dealing with quadratic polynomials is *completing the square*.
- Consider the polynomial  $x^2 + 6x + 5$ .
- We want to find a square that looks as much like that polynomial as possible.
- We square  $(x + a)$  to obtain  $(x + a)^2 = x^2 + 2ax + a^2$ .

# Completing the Square I

- An important technique for dealing with quadratic polynomials is *completing the square*.
- Consider the polynomial  $x^2 + 6x + 5$ .
- We want to find a square that looks as much like that polynomial as possible.
- We square  $(x + a)$  to obtain  $(x + a)^2 = x^2 + 2ax + a^2$ .
- The first term is correct. For the second term to be correct we need  $2a = 6$ ,  $a = 6/2 = 3$ .



## Completing the Square II

- So we have  $x^2 + 6x + 5 = (x + 3)^2 + \text{other junk}$

## Completing the Square II

- So we have  $x^2 + 6x + 5 = (x + 3)^2 + \text{other junk}$
- We figure out what the other junk is by expanding and simplifying:  $x^2 + 6x + 5 = x^2 + 6x + 9 + \text{other junk}$  so  $-4 = \text{other junk}$

## Completing the Square II

- So we have  $x^2 + 6x + 5 = (x + 3)^2 + \text{other junk}$
- We figure out what the other junk is by expanding and simplifying:  $x^2 + 6x + 5 = x^2 + 6x + 9 + \text{other junk}$  so  $-4 = \text{other junk}$
- Altogether we have  $x^2 + 6x + 5 = (x + 3)^2 - 4$

## Completing the Square II

- So we have  $x^2 + 6x + 5 = (x + 3)^2 + \text{other junk}$
- We figure out what the other junk is by expanding and simplifying:  $x^2 + 6x + 5 = x^2 + 6x + 9 + \text{other junk}$  so  $-4 = \text{other junk}$
- Altogether we have  $x^2 + 6x + 5 = (x + 3)^2 - 4$
- Now that the quadratic is in that form, we can do other useful things with it, e.g., factoring it as a difference of squares.

## Completing the Square II

- So we have  $x^2 + 6x + 5 = (x + 3)^2 + \text{other junk}$
- We figure out what the other junk is by expanding and simplifying:  $x^2 + 6x + 5 = x^2 + 6x + 9 + \text{other junk}$  so  $-4 = \text{other junk}$
- Altogether we have  $x^2 + 6x + 5 = (x + 3)^2 - 4$
- Now that the quadratic is in that form, we can do other useful things with it, e.g., factoring it as a difference of squares.
- See the supplement for the procedure to follow when the coefficient of  $x^2$  is not 1, e.g., for  $2x^2 + 8x + 6$ .

# The Quadratic Formula

- Complete the square for the general quadratic  $ax^2 + bx + c$

# The Quadratic Formula

- Complete the square for the general quadratic  $ax^2 + bx + c$
- We obtain

$$ax^2 + bx + c = a \left( x + \frac{b}{2a} \right)^2 - \left( \frac{b^2}{4a} - c \right)$$

# The Quadratic Formula

- Complete the square for the general quadratic  $ax^2 + bx + c$
- We obtain

$$ax^2 + bx + c = a \left( x + \frac{b}{2a} \right)^2 - \left( \frac{b^2}{4a} - c \right)$$

- Now if we factor by difference of squares we get the roots of a general quadratic:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



# The Quadratic Formula

- Complete the square for the general quadratic  $ax^2 + bx + c$
- We obtain

$$ax^2 + bx + c = a \left( x + \frac{b}{2a} \right)^2 - \left( \frac{b^2}{4a} - c \right)$$

- Now if we factor by difference of squares we get the roots of a general quadratic:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- That is the quadratic formula, and it can save you a lot of work if you memorize it.

# The Discriminant

- Note that in the quadratic formula, we are asked to take the square root of the expression  $b^2 - 4ac$ .

# The Discriminant

- Note that in the quadratic formula, we are asked to take the square root of the expression  $b^2 - 4ac$ .
- If that expression is negative, we can't take the square root, so the quadratic polynomial  $ax^2 + bx + c$  has no roots in that case.

# The Discriminant

- Note that in the quadratic formula, we are asked to take the square root of the expression  $b^2 - 4ac$ .
- If that expression is negative, we can't take the square root, so the quadratic polynomial  $ax^2 + bx + c$  has no roots in that case.
- If that expression is zero, the quadratic equation has two identical roots.

# The Discriminant

- Note that in the quadratic formula, we are asked to take the square root of the expression  $b^2 - 4ac$ .
- If that expression is negative, we can't take the square root, so the quadratic polynomial  $ax^2 + bx + c$  has no roots in that case.
- If that expression is zero, the quadratic equation has two identical roots.
- If that expression is positive, the quadratic equation has two different roots.

# The Discriminant

- Note that in the quadratic formula, we are asked to take the square root of the expression  $b^2 - 4ac$ .
- If that expression is negative, we can't take the square root, so the quadratic polynomial  $ax^2 + bx + c$  has no roots in that case.
- If that expression is zero, the quadratic equation has two identical roots.
- If that expression is positive, the quadratic equation has two different roots.
- Because the expression  $b^2 - 4ac$  helps us discriminate between those three cases, it is called the discriminant of the quadratic equation.

## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.

## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.
- Following that pattern,  $x^1$  should be  $x$ .



## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.
- Following that pattern,  $x^1$  should be  $x$ .
- But what is  $x^0$ ? Let's run the pattern in reverse to find out.

## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.
- Following that pattern,  $x^1$  should be  $x$ .
- But what is  $x^0$ ? Let's run the pattern in reverse to find out.
- To go from  $x^3$  to  $x^2$ , we divide by  $x$ .

## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.
- Following that pattern,  $x^1$  should be  $x$ .
- But what is  $x^0$ ? Let's run the pattern in reverse to find out.
- To go from  $x^3$  to  $x^2$ , we divide by  $x$ .
- To go from  $x^2$  to  $x^1$ , we divide by  $x$ .

## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.
- Following that pattern,  $x^1$  should be  $x$ .
- But what is  $x^0$ ? Let's run the pattern in reverse to find out.
- To go from  $x^3$  to  $x^2$ , we divide by  $x$ .
- To go from  $x^2$  to  $x^1$ , we divide by  $x$ .
- To go from  $x^1$  to  $x^0$ , we should divide by  $x$ .

## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.
- Following that pattern,  $x^1$  should be  $x$ .
- But what is  $x^0$ ? Let's run the pattern in reverse to find out.
- To go from  $x^3$  to  $x^2$ , we divide by  $x$ .
- To go from  $x^2$  to  $x^1$ , we divide by  $x$ .
- To go from  $x^1$  to  $x^0$ , we should divide by  $x$ .
- $x/x = 1$  so  $x^0$  is 1.

## Powers of a Variable

- We have already seen a short form for  $xx$ , namely  $x^2$ . Similarly a short form for  $xxx$  is  $x^3$ , and so on.
- Following that pattern,  $x^1$  should be  $x$ .
- But what is  $x^0$ ? Let's run the pattern in reverse to find out.
- To go from  $x^3$  to  $x^2$ , we divide by  $x$ .
- To go from  $x^2$  to  $x^1$ , we divide by  $x$ .
- To go from  $x^1$  to  $x^0$ , we should divide by  $x$ .
- $x/x = 1$  so  $x^0$  is 1.
- Similarly,  $x^{-1}$  is  $1/x$ ,  $x^{-2}$  is  $(1/x)/x = 1/x^2$ , and so on.

# The Binomial Theorem

- Sometimes we have to deal with powers of more complicated expressions.

# The Binomial Theorem

- Sometimes we have to deal with powers of more complicated expressions.
- For example,  $(x + y)^2 = x^2 + 2xy + y^2$ .



# The Binomial Theorem

- Sometimes we have to deal with powers of more complicated expressions.
- For example,  $(x + y)^2 = x^2 + 2xy + y^2$ .
- Multiplying both sides again by  $x + y$ , and expanding and gathering the RHS,

$$(x + y)^3 = (x^2 + 2xy + y^2)(x + y) = \cdots = x^3 + 3x^2y + 3xy^2 + y^3$$

# The Binomial Theorem

- Sometimes we have to deal with powers of more complicated expressions.
- For example,  $(x + y)^2 = x^2 + 2xy + y^2$ .
- Multiplying both sides again by  $x + y$ , and expanding and gathering the RHS,

$$(x + y)^3 = (x^2 + 2xy + y^2)(x + y) = \dots = x^3 + 3x^2y + 3xy^2 + y^3$$

- Doing that again give

$$(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

# The Binomial Theorem

- Sometimes we have to deal with powers of more complicated expressions.
- For example,  $(x + y)^2 = x^2 + 2xy + y^2$ .
- Multiplying both sides again by  $x + y$ , and expanding and gathering the RHS,

$$(x + y)^3 = (x^2 + 2xy + y^2)(x + y) = \dots = x^3 + 3x^2y + 3xy^2 + y^3$$

- Doing that again give

$$(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

- There is a pattern to these results called the *binomial theorem*. See the textbook supplement for details.

# Laws of Exponents

- Consider  $x^2 \times x^3$ .

# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.

# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.
- The result should be  $x^5$ . But note  $2 + 3 = 5$ .

# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.
- The result should be  $x^5$ . But note  $2 + 3 = 5$ .
- In general, we have  $x^m \times x^n = x^{m+n}$ .

# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.
- The result should be  $x^5$ . But note  $2 + 3 = 5$ .
- In general, we have  $x^m \times x^n = x^{m+n}$ .
- Similarly  $x^m / x^n = x^{m-n}$ .



# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.
- The result should be  $x^5$ . But note  $2 + 3 = 5$ .
- In general, we have  $x^m \times x^n = x^{m+n}$ .
- Similarly  $x^m / x^n = x^{m-n}$ .
- Also  $(x^m)^n$  is  $n$  collections of  $m$   $x$ 's, so we have  $(x^m)^n = x^{mn}$ .

# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.
- The result should be  $x^5$ . But note  $2 + 3 = 5$ .
- In general, we have  $x^m \times x^n = x^{m+n}$ .
- Similarly  $x^m / x^n = x^{m-n}$ .
- Also  $(x^m)^n$  is  $n$  collections of  $m$   $x$ 's, so we have  $(x^m)^n = x^{mn}$ .
- Another common situation is like  $(xy)^2 = xyxy = xxyy = x^2y^2$ .

# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.
- The result should be  $x^5$ . But note  $2 + 3 = 5$ .
- In general, we have  $x^m \times x^n = x^{m+n}$ .
- Similarly  $x^m / x^n = x^{m-n}$ .
- Also  $(x^m)^n$  is  $n$  collections of  $m$   $x$ 's, so we have  $(x^m)^n = x^{mn}$ .
- Another common situation is like  $(xy)^2 = xyxy = xxyy = x^2y^2$ .
- In general,  $(xy)^m = x^m y^m$ .

# Laws of Exponents

- Consider  $x^2 \times x^3$ .
- That is the product of two  $x$ 's times the product of three  $x$ 's.
- The result should be  $x^5$ . But note  $2 + 3 = 5$ .
- In general, we have  $x^m \times x^n = x^{m+n}$ .
- Similarly  $x^m / x^n = x^{m-n}$ .
- Also  $(x^m)^n$  is  $n$  collections of  $m$   $x$ 's, so we have  $(x^m)^n = x^{mn}$ .
- Another common situation is like  $(xy)^2 = xyxy = xxyy = x^2y^2$ .
- In general,  $(xy)^m = x^m y^m$ .
- If you think about it, you'll see that those results hold for negative values of  $m$  and/or  $n$  as well.

# Fractional Powers

- We know what  $x^n$  means for  $n = 0, 1, 2, \dots$  and  $n = -1, -2, -3, \dots$

# Fractional Powers

- We know what  $x^n$  means for  $n = 0, 1, 2, \dots$  and  $n = -1, -2, -3, \dots$
- Now we would like to figure out what  $x^n$  means for fractional values of  $n$ , for example  $n = 1/2$ .

# Fractional Powers

- We know what  $x^n$  means for  $n = 0, 1, 2, \dots$  and  $n = -1, -2, -3, \dots$
- Now we would like to figure out what  $x^n$  means for fractional values of  $n$ , for example  $n = 1/2$ .
- We do that by crossing our fingers and hoping that the laws of exponents will apply to fractional exponents.

# Fractional Powers

- We know what  $x^n$  means for  $n = 0, 1, 2, \dots$  and  $n = -1, -2, -3, \dots$
- Now we would like to figure out what  $x^n$  means for fractional values of  $n$ , for example  $n = 1/2$ .
- We do that by crossing our fingers and hoping that the laws of exponents will apply to fractional exponents.
- If that's the case then  $(x^{1/2})^2$  should be  $x^{(1/2) \times 2} = x^1$ .



# Fractional Powers

- We know what  $x^n$  means for  $n = 0, 1, 2, \dots$  and  $n = -1, -2, -3, \dots$
- Now we would like to figure out what  $x^n$  means for fractional values of  $n$ , for example  $n = 1/2$ .
- We do that by crossing our fingers and hoping that the laws of exponents will apply to fractional exponents.
- If that's the case then  $(x^{1/2})^2$  should be  $x^{(1/2) \times 2} = x^1$ .
- In other words,  $x^{1/2}$  should be  $\sqrt{x}$ .

# Fractional Powers

- We know what  $x^n$  means for  $n = 0, 1, 2, \dots$  and  $n = -1, -2, -3, \dots$
- Now we would like to figure out what  $x^n$  means for fractional values of  $n$ , for example  $n = 1/2$ .
- We do that by crossing our fingers and hoping that the laws of exponents will apply to fractional exponents.
- If that's the case then  $(x^{1/2})^2$  should be  $x^{(1/2) \times 2} = x^1$ .
- In other words,  $x^{1/2}$  should be  $\sqrt{x}$ .
- It turns out that everything works out fine if we say  $x^{1/2} = \sqrt{x}$ ,  $x^{1/3} = \sqrt[3]{x}$ , and so on.

# Radicals

- It is better to write  $x^{1/2}$  instead of  $\sqrt{x}$ , but the latter expression is very common so we should learn how to deal with it.

# Radicals

- It is better to write  $x^{1/2}$  instead of  $\sqrt{x}$ , but the latter expression is very common so we should learn how to deal with it.
- For example, by the laws of exponents, we have

$$\sqrt{xy} = (xy)^{1/2} = x^{1/2}y^{1/2} = \sqrt{x}\sqrt{y}$$

# Radicals

- It is better to write  $x^{1/2}$  instead of  $\sqrt{x}$ , but the latter expression is very common so we should learn how to deal with it.
- For example, by the laws of exponents, we have

$$\sqrt{xy} = (xy)^{1/2} = x^{1/2}y^{1/2} = \sqrt{x}\sqrt{y}$$

- NOTE: you *cannot* write

$$\sqrt{x+y} = \sqrt{x} + \sqrt{y}$$

# Radicals

- It is better to write  $x^{1/2}$  instead of  $\sqrt{x}$ , but the latter expression is very common so we should learn how to deal with it.
- For example, by the laws of exponents, we have

$$\sqrt{xy} = (xy)^{1/2} = x^{1/2}y^{1/2} = \sqrt{x}\sqrt{y}$$

- NOTE: you *cannot* write

$$\sqrt{x+y} = \sqrt{x} + \sqrt{y}$$

- That is wrong, wrong, wrong! Try  $x = 9$  and  $y = 16$ .

# Radicals

- It is better to write  $x^{1/2}$  instead of  $\sqrt{x}$ , but the latter expression is very common so we should learn how to deal with it.
- For example, by the laws of exponents, we have

$$\sqrt{xy} = (xy)^{1/2} = x^{1/2}y^{1/2} = \sqrt{x}\sqrt{y}$$

- NOTE: you *cannot* write

$$\sqrt{x+y} = \sqrt{x} + \sqrt{y}$$

- That is wrong, wrong, wrong! Try  $x = 9$  and  $y = 16$ .
- (You could handle  $\sqrt{x+y}$  using an extended version of the binomial theorem, but that is beyond our powers at the moment.)

# Rationalizing

- There is a useful trick with radicals we should know. Note that the difference of squares factorization gives

$$(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) = x - y$$



# Rationalizing

- There is a useful trick with radicals we should know. Note that the difference of squares factorization gives

$$(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) = x - y$$

- Suppose we want to simplify the numerator in an expression like

$$\frac{\sqrt{x} + \sqrt{y}}{4}$$

# Rationalizing

- There is a useful trick with radicals we should know. Note that the difference of squares factorization gives

$$(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) = x - y$$

- Suppose we want to simplify the numerator in an expression like

$$\frac{\sqrt{x} + \sqrt{y}}{4}$$

- We can do that (at the expense of making the denominator more complicated) by multiplying by a specially chosen representation of 1 based on the *conjugate radical*  $\sqrt{x} - \sqrt{y}$ .

# Rationalizing

- There is a useful trick with radicals we should know. Note that the difference of squares factorization gives

$$(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) = x - y$$

- Suppose we want to simplify the numerator in an expression like

$$\frac{\sqrt{x} + \sqrt{y}}{4}$$

- We can do that (at the expense of making the denominator more complicated) by multiplying by a specially chosen representation of 1 based on the *conjugate radical*  $\sqrt{x} - \sqrt{y}$ .
- We have

$$\frac{\sqrt{x} + \sqrt{y}}{4} = \frac{\sqrt{x} + \sqrt{y}}{4} \times \frac{\sqrt{x} - \sqrt{y}}{\sqrt{x} - \sqrt{y}} = \frac{x - y}{4(\sqrt{x} - \sqrt{y})}$$