

MATH 110 Lecture 2.5

The Chain Rule

Edward Doolittle

Tuesday, February 10, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

The Chain Rule

Composition of Functions

Derivation of the Chain Rule

Application of the Chain Rule

Examples and Exercises

The Chain Rule

Review of Composition of Functions

- Recall the concept of chaining or composing functions:

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

Review of Composition of Functions

- Recall the concept of chaining or composing functions:

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

- We consider $x \mapsto f(g(x))$ a single function called the composition of f and g , sometimes written $f \circ g$.

Review of Composition of Functions

- Recall the concept of chaining or composing functions:

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

- We consider $x \mapsto f(g(x))$ a single function called the composition of f and g , sometimes written $f \circ g$.
- We already know a rule for limits of compositions of functions, namely

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x))$$

provided f is continuous at $\lim_{x \rightarrow a} g(x)$.

Review of Composition of Functions

- Recall the concept of chaining or composing functions:

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

- We consider $x \mapsto f(g(x))$ a single function called the composition of f and g , sometimes written $f \circ g$.
- We already know a rule for limits of compositions of functions, namely

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x))$$

provided f is continuous at $\lim_{x \rightarrow a} g(x)$.

- We would like to develop a rule for derivatives of compositions of functions.

Identifying a Composition of Functions

- Before we get into derivatives, however, it might be a good idea to better understand when we have a composition of functions.

Identifying a Composition of Functions

- Before we get into derivatives, however, it might be a good idea to better understand when we have a composition of functions.
- Compositions of functions will seldom, if ever, be presented in a form like $f \circ g$ where f and g are immediately obvious. Instead, we will have to find f and g .

Identifying a Composition of Functions

- Before we get into derivatives, however, it might be a good idea to better understand when we have a composition of functions.
- Compositions of functions will seldom, if ever, be presented in a form like $f \circ g$ where f and g are immediately obvious. Instead, we will have to find f and g .
- Consider $h(x) = \sin(2x + \pi)$. We read the expression on the right as

the sine of two x plus π

The separator between f and g is usually the word ‘of’. In this case $f(u) = \sin u$ and $g(x) = 2x + \pi$.

Identifying a Composition of Functions

- Before we get into derivatives, however, it might be a good idea to better understand when we have a composition of functions.
- Compositions of functions will seldom, if ever, be presented in a form like $f \circ g$ where f and g are immediately obvious. Instead, we will have to find f and g .
- Consider $h(x) = \sin(2x + \pi)$. We read the expression on the right as

the sine of two x plus π

The separator between f and g is usually the word ‘of’. In this case $f(u) = \sin u$ and $g(x) = 2x + \pi$.

- Another way of thinking of it is that $g(x) = 2x + \pi$ is the ‘inner function’, i.e., the function inside brackets. So $f(u) = \sin u$ is the outer function. The u is just a placeholder.

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

- Letting $u = g(x)$, $du = g'(x)dx$, we have

$$f(g(x + dx))$$

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

- Letting $u = g(x)$, $du = g'(x)dx$, we have

$$f(g(x + dx)) = f(g(x) + g'(x)dx + \dots)$$

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

- Letting $u = g(x)$, $du = g'(x)dx$, we have

$$\begin{aligned} f(g(x + dx)) &= f(g(x) + g'(x)dx + \dots) \\ &= f(u + du) \end{aligned}$$

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

- Letting $u = g(x)$, $du = g'(x)dx$, we have

$$\begin{aligned} f(g(x + dx)) &= f(g(x) + g'(x)dx + \dots) \\ &= f(u + du) = f(u) + f'(u)du + \dots \end{aligned}$$

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

- Letting $u = g(x)$, $du = g'(x)dx$, we have

$$\begin{aligned} f(g(x + dx)) &= f(g(x) + g'(x)dx + \dots) \\ &= f(u + du) = f(u) + f'(u)du + \dots \\ &= f(g(x)) + f'(g(x))g'(x)dx + \dots \end{aligned}$$

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

- Letting $u = g(x)$, $du = g'(x)dx$, we have

$$\begin{aligned} f(g(x + dx)) &= f(g(x) + g'(x)dx + \dots) \\ &= f(u + du) = f(u) + f'(u)du + \dots \\ &= f(g(x)) + f'(g(x))g'(x)dx + \dots \end{aligned}$$

- This suggests that the derivative of $f(g(x))$ is $f'(g(x)) \cdot g'(x)$.

Derivation of the Chain Rule

- We want to differentiate $f(g(x))$.
- Recall that we can write

$$f(u + du) = f(u) + f'(u)du + \dots$$

$$g(x + dx) = g(x) + g'(x)dx + \dots$$

where \dots represents terms with factors of du^2 and dx^2 .

- Letting $u = g(x)$, $du = g'(x)dx$, we have

$$\begin{aligned} f(g(x + dx)) &= f(g(x) + g'(x)dx + \dots) \\ &= f(u + du) = f(u) + f'(u)du + \dots \\ &= f(g(x)) + f'(g(x))g'(x)dx + \dots \end{aligned}$$

- This suggests that the derivative of $f(g(x))$ is $f'(g(x)) \cdot g'(x)$.
- The result is proven in the textbook, but you should skip the proof unless you are majoring in math.

The Chain Rule in Newton and Leibniz Notation

- In summary, the chain rule in Newton notation is

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

The Chain Rule in Newton and Leibniz Notation

- In summary, the chain rule in Newton notation is

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

- The derivative of a composition is the derivative of the outer function evaluated at the inner function, times the derivative of the inner function.

The Chain Rule in Newton and Leibniz Notation

- In summary, the chain rule in Newton notation is

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

- The derivative of a composition is the derivative of the outer function evaluated at the inner function, times the derivative of the inner function.
- The chain rule in Leibniz notation is

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

The Chain Rule in Newton and Leibniz Notation

- In summary, the chain rule in Newton notation is

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

- The derivative of a composition is the derivative of the outer function evaluated at the inner function, times the derivative of the inner function.
- The chain rule in Leibniz notation is

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

- In the Leibniz rule, y and u have changing meanings.

The Chain Rule in Newton and Leibniz Notation

- In summary, the chain rule in Newton notation is

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

- The derivative of a composition is the derivative of the outer function evaluated at the inner function, times the derivative of the inner function.
- The chain rule in Leibniz notation is

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

- In the Leibniz rule, y and u have changing meanings.
- The way I personally think of the chain rule is

$$\frac{d}{dx}f(g(x)) = f'(g(x)) \cdot \frac{d}{dx}g(x)$$

Application of the Chain Rule

- To apply the chain rule we need to

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g
 4. assemble the formula $f'(g(x)) \cdot g'(x)$

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g
 4. assemble the formula $f'(g(x)) \cdot g'(x)$
- Consider the function $h(x) = \sin(5x)$

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g
 4. assemble the formula $f'(g(x)) \cdot g'(x)$
- Consider the function $h(x) = \sin(5x)$
- We read this as ‘the sine *of* $5x$ ’.

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g
 4. assemble the formula $f'(g(x)) \cdot g'(x)$
- Consider the function $h(x) = \sin(5x)$
- We read this as ‘the sine *of* $5x$ ’.
- It is a composition with $f(u) = \sin u$, $g(x) = 5x$.

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g
 4. assemble the formula $f'(g(x)) \cdot g'(x)$
- Consider the function $h(x) = \sin(5x)$
- We read this as ‘the sine *of* $5x$ ’.
- It is a composition with $f(u) = \sin u$, $g(x) = 5x$.
- We have $f'(u) = \cos u$, $g'(x) = 5$.

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g
 4. assemble the formula $f'(g(x)) \cdot g'(x)$
- Consider the function $h(x) = \sin(5x)$
- We read this as ‘the sine *of* $5x$ ’.
- It is a composition with $f(u) = \sin u$, $g(x) = 5x$.
- We have $f'(u) = \cos u$, $g'(x) = 5$.
- The chain rule tells us the derivative of $h(x) = f(g(x))$ is
$$h'(x) = f'(g(x)) \cdot g'(x) = \cos(5x) \cdot 5.$$

Application of the Chain Rule

- To apply the chain rule we need to
 1. identify when we have a composition
 2. identify the inner function g and outer function f
 3. differentiate both f and g
 4. assemble the formula $f'(g(x)) \cdot g'(x)$
- Consider the function $h(x) = \sin(5x)$
- We read this as ‘the sine *of* $5x$ ’.
- It is a composition with $f(u) = \sin u$, $g(x) = 5x$.
- We have $f'(u) = \cos u$, $g'(x) = 5$.
- The chain rule tells us the derivative of $h(x) = f(g(x))$ is
$$h'(x) = f'(g(x)) \cdot g'(x) = \cos(5x) \cdot 5.$$
- Conventionally, we rearrange expressions like that to the form
$$h'(x) = 5 \cos 5x.$$

Application of the Chain Rule

- Consider the function $y = \cos^3 x$.

Application of the Chain Rule

- Consider the function $y = \cos^3 x$.
- That means $y = (\cos x)^3$. It is a composition of functions.

Application of the Chain Rule

- Consider the function $y = \cos^3 x$.
- That means $y = (\cos x)^3$. It is a composition of functions.
- The outer function is $f(u) = u^3$. The inner function is $g(x) = \cos x$.

Application of the Chain Rule

- Consider the function $y = \cos^3 x$.
- That means $y = (\cos x)^3$. It is a composition of functions.
- The outer function is $f(u) = u^3$. The inner function is $g(x) = \cos x$.
- Taking derivatives, we have $f'(u) = 3u^2$, $g'(x) = -\sin x$.

Application of the Chain Rule

- Consider the function $y = \cos^3 x$.
- That means $y = (\cos x)^3$. It is a composition of functions.
- The outer function is $f(u) = u^3$. The inner function is $g(x) = \cos x$.
- Taking derivatives, we have $f'(u) = 3u^2$, $g'(x) = -\sin x$.
- Note that we have $f'(g(x)) = 3(g(x))^2 = 3(\cos x)^2$.

Application of the Chain Rule

- Consider the function $y = \cos^3 x$.
- That means $y = (\cos x)^3$. It is a composition of functions.
- The outer function is $f(u) = u^3$. The inner function is $g(x) = \cos x$.
- Taking derivatives, we have $f'(u) = 3u^2$, $g'(x) = -\sin x$.
- Note that we have $f'(g(x)) = 3(g(x))^2 = 3(\cos x)^2$.
- Substituting into the chain rule, we have

$$y' = f'(g(x)) \cdot g'(x) = 3(\cos x)^2 \cdot -\sin x$$

Application of the Chain Rule

- Consider the function $y = \cos^3 x$.
- That means $y = (\cos x)^3$. It is a composition of functions.
- The outer function is $f(u) = u^3$. The inner function is $g(x) = \cos x$.
- Taking derivatives, we have $f'(u) = 3u^2$, $g'(x) = -\sin x$.
- Note that we have $f'(g(x)) = 3(g(x))^2 = 3(\cos x)^2$.
- Substituting into the chain rule, we have

$$y' = f'(g(x)) \cdot g'(x) = 3(\cos x)^2 \cdot -\sin x$$

- We can rearrange if we like:

$$y' = -3\cos^2 x \sin x$$

Special Case: Power-chain rule

- A common special case of the chain rule is when the outer function is a power function of the form u^p .

Special Case: Power-chain rule

- A common special case of the chain rule is when the outer function is a power function of the form u^p .
- The previous example $y = (\cos x)^3$ was of this form.

Special Case: Power-chain rule

- A common special case of the chain rule is when the outer function is a power function of the form u^p .
- The previous example $y = (\cos x)^3$ was of this form.
- Another example is $h(x) = (3x + 2)^{2.5}$, with derivative

$$\begin{aligned} h'(x) &= 2.5(3x + 2)^{1.5} \cdot \frac{d}{dx}(3x + 2) = 2.5(3x + 2)^{1.5} \cdot 3 \\ &= 7.5(3x + 2)^{1.5} \end{aligned}$$

Special Case: Power-chain rule

- A common special case of the chain rule is when the outer function is a power function of the form u^p .
- The previous example $y = (\cos x)^3$ was of this form.
- Another example is $h(x) = (3x + 2)^{2.5}$, with derivative

$$\begin{aligned} h'(x) &= 2.5(3x + 2)^{1.5} \cdot \frac{d}{dx}(3x + 2) = 2.5(3x + 2)^{1.5} \cdot 3 \\ &= 7.5(3x + 2)^{1.5} \end{aligned}$$

- The special case power-chain rule can be written

$$\frac{d}{dx}[g(x)]^p = p[g(x)]^{p-1} \cdot \frac{d}{dx}g(x)$$

Longer Chains of Functions

- Consider the function $y = \sin(\cos(\tan x))$, which is a chain of three functions.

Longer Chains of Functions

- Consider the function $y = \sin(\cos(\tan x))$, which is a chain of three functions.
- First, \sin is the outer function, $\cos(\tan x)$ is the inner function:

$$y' = \frac{d}{dx} \sin(\cos(\tan x))$$

Longer Chains of Functions

- Consider the function $y = \sin(\cos(\tan x))$, which is a chain of three functions.
- First, \sin is the outer function, $\cos(\tan x)$ is the inner function:

$$\begin{aligned} y' &= \frac{d}{dx} \sin(\cos(\tan x)) \\ &= \sin'(\cos(\tan x)) \cdot \frac{d}{dx} \cos(\tan x) \end{aligned}$$

Longer Chains of Functions

- Consider the function $y = \sin(\cos(\tan x))$, which is a chain of three functions.
- First, \sin is the outer function, $\cos(\tan x)$ is the inner function:

$$\begin{aligned}y' &= \frac{d}{dx} \sin(\cos(\tan x)) \\&= \sin'(\cos(\tan x)) \cdot \frac{d}{dx} \cos(\tan x) \\&= \sin'(\cos(\tan x)) \cdot \cos'(\tan x) \cdot \frac{d}{dx} \tan x\end{aligned}$$

Longer Chains of Functions

- Consider the function $y = \sin(\cos(\tan x))$, which is a chain of three functions.
- First, \sin is the outer function, $\cos(\tan x)$ is the inner function:

$$\begin{aligned}y' &= \frac{d}{dx} \sin(\cos(\tan x)) \\&= \sin'(\cos(\tan x)) \cdot \frac{d}{dx} \cos(\tan x) \\&= \sin'(\cos(\tan x)) \cdot \cos'(\tan x) \cdot \frac{d}{dx} \tan x \\&= \cos(\cos(\tan x)) \cdot -\sin(\tan x) \cdot \sec^2 x\end{aligned}$$

Longer Chains of Functions

- Consider the function $y = \sin(\cos(\tan x))$, which is a chain of three functions.
- First, \sin is the outer function, $\cos(\tan x)$ is the inner function:

$$\begin{aligned}y' &= \frac{d}{dx} \sin(\cos(\tan x)) \\&= \sin'(\cos(\tan x)) \cdot \frac{d}{dx} \cos(\tan x) \\&= \sin'(\cos(\tan x)) \cdot \cos'(\tan x) \cdot \frac{d}{dx} \tan x \\&= \cos(\cos(\tan x)) \cdot -\sin(\tan x) \cdot \sec^2 x\end{aligned}$$

- If you have a function of the form $y = \cos^3 x^2$, you might try re-writing it before applying the chain rule.

Longer Chains of Functions

- Consider the function $y = \sin(\cos(\tan x))$, which is a chain of three functions.
- First, \sin is the outer function, $\cos(\tan x)$ is the inner function:

$$\begin{aligned}y' &= \frac{d}{dx} \sin(\cos(\tan x)) \\&= \sin'(\cos(\tan x)) \cdot \frac{d}{dx} \cos(\tan x) \\&= \sin'(\cos(\tan x)) \cdot \cos'(\tan x) \cdot \frac{d}{dx} \tan x \\&= \cos(\cos(\tan x)) \cdot -\sin(\tan x) \cdot \sec^2 x\end{aligned}$$

- If you have a function of the form $y = \cos^3 x^2$, you might try re-writing it before applying the chain rule.
- $y = f(\cos(h(x)))$ where $f(u) = u^3$ and $h(x) = x^2$.

Examples

1. Calculate y' where $y = \cos(\tan x)$.
2. If $f(t) = \sqrt{4t+1}$, find $f''(2)$.
3. Find an equation for the tangent line and normal line to $y = 4\sin^2 x$ at the point $(\pi/6, 1)$ on the curve.
4. Differentiate the trig identities
 - 4.1 $\cos 2x = \cos^2 x - \sin^2 x$
 - 4.2 $\sin(x+a) = \sin x \cos a + \cos x \sin a$to obtain new trig identities.

Now you should work on Problem Set 2.5. After you have finished it, you should try the following additional exercises from Section 2.5:

2.5 C-level: 1–6, 7–46, 47–50, 51–54, 55–57, 59–60, 61–64, 67–68, 69–72

B-level: 58, 65–66, 73–74, 75–76, 77–80

A-level: 83–90