

MATH 110 Lecture 2.4

Derivatives of Trigonometric Functions

Edward Doolittle

Thursday, February 5, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Derivatives of Trigonometric Functions

The Basic Trig Limits

Other Trig Limits

Derivatives of Trig Functions

Examples and Exercises

Derivatives of Trigonometric Functions

Strategy for Finding Derivatives of Trig Functions

- In this course we will be studying the following trigonometric functions:

\sin \cos \tan

\csc \sec \cot

Strategy for Finding Derivatives of Trig Functions

- In this course we will be studying the following trigonometric functions:

\sin \cos \tan

\csc \sec \cot

- We would like to be able to differentiate those functions.

Strategy for Finding Derivatives of Trig Functions

- In this course we will be studying the following trigonometric functions:

\sin \cos \tan

\csc \sec \cot

- We would like to be able to differentiate those functions.
- The strategy we will employ is to reduce everything to \sin and \cos .

Strategy for Finding Derivatives of Trig Functions

- In this course we will be studying the following trigonometric functions:

\sin \cos \tan

\csc \sec \cot

- We would like to be able to differentiate those functions.
- The strategy we will employ is to reduce everything to \sin and \cos .
- We still need to differentiate those functions. In order to do so, we will have to use the definition of the derivative in terms of limits.

Strategy for Finding Derivatives of Trig Functions

- In this course we will be studying the following trigonometric functions:

\sin \cos \tan

\csc \sec \cot

- We would like to be able to differentiate those functions.
- The strategy we will employ is to reduce everything to \sin and \cos .
- We still need to differentiate those functions. In order to do so, we will have to use the definition of the derivative in terms of limits.
- That means we will need to be able to calculate limits of \sin and \cos , which is the problem to which we now turn.

The Basic Trig Limits

- We have already seen two basic trig limits in connection with our study of the continuity of sin and cos:

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \text{and} \quad \lim_{h \rightarrow 0} \cos h = 1$$

The Basic Trig Limits

- We have already seen two basic trig limits in connection with our study of the continuity of \sin and \cos :

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \text{and} \quad \lim_{h \rightarrow 0} \cos h = 1$$

- If we are going to calculate $\sin'(0)$, we'll need a new trig limit.

The Basic Trig Limits

- We have already seen two basic trig limits in connection with our study of the continuity of \sin and \cos :

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \text{and} \quad \lim_{h \rightarrow 0} \cos h = 1$$

- If we are going to calculate $\sin'(0)$, we'll need a new trig limit.
- By the definition of derivative,

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin(0)}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h}$$

The Basic Trig Limits

- We have already seen two basic trig limits in connection with our study of the continuity of \sin and \cos :

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \text{and} \quad \lim_{h \rightarrow 0} \cos h = 1$$

- If we are going to calculate $\sin'(0)$, we'll need a new trig limit.
- By the definition of derivative,

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin(0)}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h}$$

- So we need the latter limit. Recall that $\sin x \approx x$ (in radians), so we expect

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

The Basic Trig Limits

- We have already seen two basic trig limits in connection with our study of the continuity of \sin and \cos :

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \text{and} \quad \lim_{h \rightarrow 0} \cos h = 1$$

- If we are going to calculate $\sin'(0)$, we'll need a new trig limit.
- By the definition of derivative,

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin(0)}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h}$$


- So we need the latter limit. Recall that $\sin x \approx x$ (in radians), so we expect

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

- All of our results for trig functions will follow from those limits.


The Big Idea Behind the Proof of a Basic Trig Limit

Consider the diagram on the right.




The Big Idea Behind the Proof of a Basic Trig Limit

We calculate the lengths of various lines in the diagram in terms of the angle x .




The Big Idea Behind the Proof of a Basic Trig Limit

We calculate the lengths of various lines in the diagram in terms of the angle x .




The Big Idea Behind the Proof of a Basic Trig Limit

We calculate the lengths of various lines in the diagram in terms of the angle x .




The Big Idea Behind the Proof of a Basic Trig Limit

We calculate the lengths of various lines in the diagram in terms of the angle x .




The Big Idea Behind the Proof of a Basic Trig Limit

We calculate the lengths of various lines in the diagram in terms of the angle x .




The Big Idea Behind the Proof of a Basic Trig Limit

Now consider the areas of various regions in the diagram.




The Big Idea Behind the Proof of a Basic Trig Limit

Now consider the areas of various regions in the diagram.



The Big Idea Behind the Proof of a Basic Trig Limit


Now consider the areas of various regions in the diagram.



The Big Idea Behind the Proof of a Basic Trig Limit

It should be clear from the diagram that


$$\text{area}(A) \leq \text{area}(B) \leq \text{area}(C)$$



The Big Idea Behind the Proof of a Basic Trig Limit

It should be clear from the diagram that


$$\text{area}(A) \leq \text{area}(B) \leq \text{area}(C)$$



The Big Idea Behind the Proof of a Basic Trig Limit

It should be clear from the diagram that


$$\text{area}(A) \leq \text{area}(B) \leq \text{area}(C)$$



The Big Idea Behind the Proof of a Basic Trig Limit

Now we calculate the various areas.


$$\text{area}(A) \leq \text{area}(B) \leq \text{area}(C)$$



The Big Idea Behind the Proof of a Basic Trig Limit

Now we calculate the various areas.


$$\frac{1}{2} \sin x \cos x \leq \text{area}(B) \leq \text{area}(C)$$



The Big Idea Behind the Proof of a Basic Trig Limit

Now we calculate the various areas.


$$\frac{1}{2} \sin x \cos x \leq \frac{1}{2} x \leq \text{area}(C)$$



The Big Idea Behind the Proof of a Basic Trig Limit

Now we calculate the various areas.


$$\frac{1}{2} \sin x \cos x \leq \frac{1}{2} x \leq \frac{1}{2} \tan x$$



The Big Idea Behind the Proof of a Basic Trig Limit

Now we calculate the various areas.


$$\frac{1}{2} \sin x \cos x \leq \frac{1}{2} x \leq \frac{1}{2} \frac{\sin x}{\cos x}$$



The Big Idea Behind the Proof of a Basic Trig Limit

Simplifying,


$$\frac{1}{2} \sin x \cos x \leq \frac{1}{2} x \leq \frac{1}{2} \frac{\sin x}{\cos x}$$



The Big Idea Behind the Proof of a Basic Trig Limit

Simplifying,


$$\sin x \cos x \leq x \leq \frac{\sin x}{\cos x}$$



The Big Idea Behind the Proof of a Basic Trig Limit

Simplifying,


$$\cos x \leq \frac{x}{\sin x} \leq \frac{1}{\cos x}$$



The Big Idea Behind the Proof of a Basic Trig Limit

Taking limits,


$$\lim_{x \rightarrow 0} \cos x \leq \lim_{x \rightarrow 0} \frac{x}{\sin x} \leq \lim_{x \rightarrow 0} \frac{1}{\cos x}$$



The Big Idea Behind the Proof of a Basic Trig Limit

Applying the squeeze theorem,

$$1 \leq \lim_{x \rightarrow 0} \frac{x}{\sin x} \leq 1$$



The Limit $\lim_{h \rightarrow 0} \frac{\tan h}{h}$

- The basic trig limits

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \lim_{h \rightarrow 0} \cos h = 1 \quad \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

allow us to compute more complicated limits of trig functions.

The Limit $\lim_{h \rightarrow 0} \frac{\tan h}{h}$

- The basic trig limits

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \lim_{h \rightarrow 0} \cos h = 1 \quad \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

allow us to compute more complicated limits of trig functions.

- For example,

$$\lim_{h \rightarrow 0} \frac{\tan h}{h}$$

The Limit $\lim_{h \rightarrow 0} \frac{\tan h}{h}$

- The basic trig limits

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \lim_{h \rightarrow 0} \cos h = 1 \quad \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

allow us to compute more complicated limits of trig functions.

- For example,

$$\lim_{h \rightarrow 0} \frac{\tan h}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} \frac{1}{\cos h}$$

The Limit $\lim_{h \rightarrow 0} \frac{\tan h}{h}$

- The basic trig limits

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \lim_{h \rightarrow 0} \cos h = 1 \quad \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

allow us to compute more complicated limits of trig functions.

- For example,

$$\lim_{h \rightarrow 0} \frac{\tan h}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} \frac{1}{\cos h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} \lim_{h \rightarrow 0} \frac{1}{\cos h}$$

The Limit $\lim_{h \rightarrow 0} \frac{\tan h}{h}$

- The basic trig limits

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \lim_{h \rightarrow 0} \cos h = 1 \quad \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

allow us to compute more complicated limits of trig functions.

- For example,

$$\lim_{h \rightarrow 0} \frac{\tan h}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} \frac{1}{\cos h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} \lim_{h \rightarrow 0} \frac{1}{\cos h} = 1 \cdot \frac{1}{1} = 1$$

The Limit $\lim_{h \rightarrow 0} \frac{\tan h}{h}$

- The basic trig limits

$$\lim_{h \rightarrow 0} \sin h = 0 \quad \lim_{h \rightarrow 0} \cos h = 1 \quad \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

allow us to compute more complicated limits of trig functions.

- For example,

$$\lim_{h \rightarrow 0} \frac{\tan h}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} \frac{1}{\cos h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} \lim_{h \rightarrow 0} \frac{1}{\cos h} = 1 \cdot \frac{1}{1} = 1$$

- In general, we try to use algebra to rearrange expressions so that the basic trig limits appear.

The Limit $\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h}$

- Sometimes we have to be clever about rearranging expressions to calculate trig limits.

The Limit $\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h}$

- Sometimes we have to be clever about rearranging expressions to calculate trig limits.
- For example,

$$\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h}$$

The Limit $\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h}$

- Sometimes we have to be clever about rearranging expressions to calculate trig limits.
- For example,

$$\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h} = \lim_{h \rightarrow 0} \frac{\sin 2h}{2h} \cdot \frac{2h}{h} \cdot \frac{h}{\tan h}$$

The Limit $\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h}$

- Sometimes we have to be clever about rearranging expressions to calculate trig limits.
- For example,

$$\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h} = \lim_{h \rightarrow 0} \frac{\sin 2h}{2h} \cdot \frac{2h}{h} \cdot \frac{h}{\tan h} = \lim_{h \rightarrow 0} \frac{\sin 2h}{2h} \cdot 2 \cdot \frac{1}{\lim_{h \rightarrow 0} \frac{\tan h}{h}}$$

The Limit $\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h}$

- Sometimes we have to be clever about rearranging expressions to calculate trig limits.
- For example,

$$\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h} = \lim_{h \rightarrow 0} \frac{\sin 2h}{2h} \cdot \frac{2h}{h} \cdot \frac{h}{\tan h} = \lim_{h \rightarrow 0} \frac{\sin 2h}{2h} \cdot 2 \cdot \frac{1}{\lim_{h \rightarrow 0} \frac{\tan h}{h}} = 2$$

The Limit $\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h}$

- Sometimes we have to be clever about rearranging expressions to calculate trig limits.
- For example,

$$\lim_{h \rightarrow 0} \frac{\sin 2h}{\tan h} = \lim_{h \rightarrow 0} \frac{\sin 2h}{2h} \cdot \frac{2h}{h} \cdot \frac{h}{\tan h} = \lim_{h \rightarrow 0} \frac{\sin 2h}{2h} \cdot 2 \cdot \frac{1}{\lim_{h \rightarrow 0} \frac{\tan h}{h}} = 2$$

- In general, we want to divide and multiply by the arguments of any sin functions that appear.

The Limit $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$

- Another important limit for which we have to be clever is

$$\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$$

The Limit $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$

- Another important limit for which we have to be clever is

$$\lim_{h \rightarrow 0} \frac{\cos h - 1}{h} = \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \cdot \frac{\cos h + 1}{\cos h + 1}$$

The Limit $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$

- Another important limit for which we have to be clever is

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{\cos h - 1}{h} &= \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \cdot \frac{\cos h + 1}{\cos h + 1} \\ &= \lim_{h \rightarrow 0} \frac{\cos^2 h - 1}{h} \cdot \frac{1}{\cos h + 1}\end{aligned}$$

The Limit $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$

- Another important limit for which we have to be clever is

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{\cos h - 1}{h} &= \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \cdot \frac{\cos h + 1}{\cos h + 1} \\ &= \lim_{h \rightarrow 0} \frac{\cos^2 h - 1}{h} \cdot \frac{1}{\cos h + 1} \\ &= \lim_{h \rightarrow 0} \frac{-\sin^2 h}{h} \cdot \frac{1}{\cos h + 1}\end{aligned}$$

The Limit $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$

- Another important limit for which we have to be clever is

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{\cos h - 1}{h} &= \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \cdot \frac{\cos h + 1}{\cos h + 1} \\&= \lim_{h \rightarrow 0} \frac{\cos^2 h - 1}{h} \cdot \frac{1}{\cos h + 1} \\&= \lim_{h \rightarrow 0} \frac{-\sin^2 h}{h} \cdot \frac{1}{\cos h + 1} \\&= - \lim_{h \rightarrow 0} \frac{\sin h}{h} \cdot \frac{\sin h}{\cos h + 1}\end{aligned}$$

The Limit $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$

- Another important limit for which we have to be clever is

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{\cos h - 1}{h} &= \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \cdot \frac{\cos h + 1}{\cos h + 1} \\&= \lim_{h \rightarrow 0} \frac{\cos^2 h - 1}{h} \cdot \frac{1}{\cos h + 1} \\&= \lim_{h \rightarrow 0} \frac{-\sin^2 h}{h} \cdot \frac{1}{\cos h + 1} \\&= - \lim_{h \rightarrow 0} \frac{\sin h}{h} \cdot \frac{\sin h}{\cos h + 1} \\&= - \lim_{h \rightarrow 0} \frac{\sin h}{h} \cdot \lim_{h \rightarrow 0} \frac{\sin h}{\cos h + 1}\end{aligned}$$

The Limit $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}$

- Another important limit for which we have to be clever is

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{\cos h - 1}{h} &= \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \cdot \frac{\cos h + 1}{\cos h + 1} \\&= \lim_{h \rightarrow 0} \frac{\cos^2 h - 1}{h} \cdot \frac{1}{\cos h + 1} \\&= \lim_{h \rightarrow 0} \frac{-\sin^2 h}{h} \cdot \frac{1}{\cos h + 1} \\&= - \lim_{h \rightarrow 0} \frac{\sin h}{h} \cdot \frac{\sin h}{\cos h + 1} \\&= - \lim_{h \rightarrow 0} \frac{\sin h}{h} \cdot \lim_{h \rightarrow 0} \frac{\sin h}{\cos h + 1} = -1 \cdot \frac{0}{1 + 1} = 0\end{aligned}$$

The Derivatives of \sin and \cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

The Derivatives of \sin and \cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0)$$

The Derivatives of \sin and \cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h}$$

The Derivatives of \sin and \cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h}$$

The Derivatives of \sin and \cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

The Derivatives of sin and cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

- We would like to find the derivative of sin at other numbers x .

The Derivatives of sin and cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

- We would like to find the derivative of sin at other numbers x .
- For that we will need the angle addition formula

$$\sin(x + h)$$

The Derivatives of sin and cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

- We would like to find the derivative of sin at other numbers x .
- For that we will need the angle addition formula

$$\sin(x + h) = \sin x \cos h + \cos x \sin h$$

The Derivatives of sin and cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

- We would like to find the derivative of sin at other numbers x .
- For that we will need the angle addition formula

$$\sin(x + h) = \sin x \cos h + \cos x \sin h$$

- Now we have

$$\sin'(x)$$

The Derivatives of sin and cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

- We would like to find the derivative of sin at other numbers x .
- For that we will need the angle addition formula

$$\sin(x + h) = \sin x \cos h + \cos x \sin h$$

- Now we have

$$\sin'(x) = \lim_{h \rightarrow 0} \frac{\sin(x + h) - \sin x}{h}$$

The Derivatives of sin and cos

- We have learned the basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$, which essentially expresses the fact that $\sin'(0) = 1$:

$$\sin'(0) = \lim_{h \rightarrow 0} \frac{\sin(0 + h) - \sin 0}{h} = \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

- We would like to find the derivative of sin at other numbers x .
- For that we will need the angle addition formula

$$\sin(x + h) = \sin x \cos h + \cos x \sin h$$

- Now we have

$$\sin'(x) = \lim_{h \rightarrow 0} \frac{\sin(x + h) - \sin x}{h} = \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h}$$

The Derivatives of \sin and \cos , continued

- Applying the difference and constant multiple limit laws ($\sin x$ and $\cos x$ are considered constants inside this calculation),

$$\sin'(x)$$

The Derivatives of \sin and \cos , continued

- Applying the difference and constant multiple limit laws ($\sin x$ and $\cos x$ are considered constants inside this calculation),

$$\sin'(x) = \lim_{h \rightarrow 0} \frac{\sin x (\cos h - 1)}{h} + \frac{\cos x \sin h}{h}$$

The Derivatives of \sin and \cos , continued

- Applying the difference and constant multiple limit laws ($\sin x$ and $\cos x$ are considered constants inside this calculation),

$$\begin{aligned}\sin'(x) &= \lim_{h \rightarrow 0} \frac{\sin x (\cos h - 1)}{h} + \frac{\cos x \sin h}{h} \\ &= \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h}\end{aligned}$$

The Derivatives of \sin and \cos , continued

- Applying the difference and constant multiple limit laws ($\sin x$ and $\cos x$ are considered constants inside this calculation),

$$\begin{aligned}\sin'(x) &= \lim_{h \rightarrow 0} \frac{\sin x (\cos h - 1)}{h} + \frac{\cos x \sin h}{h} \\ &= \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= \sin x \cdot 0 + \cos x \cdot 1\end{aligned}$$

The Derivatives of \sin and \cos , continued

- Applying the difference and constant multiple limit laws ($\sin x$ and $\cos x$ are considered constants inside this calculation),

$$\begin{aligned}\sin'(x) &= \lim_{h \rightarrow 0} \frac{\sin x (\cos h - 1)}{h} + \frac{\cos x \sin h}{h} \\ &= \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= \sin x \cdot 0 + \cos x \cdot 1 \\ &= \cos x\end{aligned}$$

The Derivatives of \sin and \cos , continued

- Applying the difference and constant multiple limit laws ($\sin x$ and $\cos x$ are considered constants inside this calculation),

$$\begin{aligned}\sin'(x) &= \lim_{h \rightarrow 0} \frac{\sin x (\cos h - 1)}{h} + \frac{\cos x \sin h}{h} \\ &= \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= \sin x \cdot 0 + \cos x \cdot 1 \\ &= \cos x\end{aligned}$$

- In summary, $\sin' x = \cos x$.

The Derivatives of \sin and \cos , continued

- Applying the difference and constant multiple limit laws ($\sin x$ and $\cos x$ are considered constants inside this calculation),

$$\begin{aligned}\sin'(x) &= \lim_{h \rightarrow 0} \frac{\sin x(\cos h - 1)}{h} + \frac{\cos x \sin h}{h} \\ &= \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= \sin x \cdot 0 + \cos x \cdot 1 \\ &= \cos x\end{aligned}$$

- In summary, $\sin' x = \cos x$.
- An analogous argument gives $\cos' x = -\sin x$. Try it!

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.
- We use radian measure because it is more natural in the math setting than other angle units.

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.
- We use radian measure because it is more natural in the math setting than other angle units.
- The basic trig limit in degree measure is

$$\lim_{h \rightarrow 0} \frac{\sindeg h}{h}$$

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.
- We use radian measure because it is more natural in the math setting than other angle units.
- The basic trig limit in degree measure is

$$\lim_{h \rightarrow 0} \frac{\sindeg h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{h}$$

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.
- We use radian measure because it is more natural in the math setting than other angle units.
- The basic trig limit in degree measure is

$$\lim_{h \rightarrow 0} \frac{\sindeg h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{\frac{\pi}{180} h} \cdot \frac{\pi}{180}$$

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.
- We use radian measure because it is more natural in the math setting than other angle units.
- The basic trig limit in degree measure is

$$\lim_{h \rightarrow 0} \frac{\sindeg h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{\frac{\pi}{180} h} \cdot \frac{\pi}{180} = \frac{\pi}{180}$$

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.
- We use radian measure because it is more natural in the math setting than other angle units.
- The basic trig limit in degree measure is

$$\lim_{h \rightarrow 0} \frac{\sindeg h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{\frac{\pi}{180} h} \cdot \frac{\pi}{180} = \frac{\pi}{180}$$

- That crazy factor of $\pi/180$ will screw up every other calculus formula involving trig functions.

A Word on Radian Measure

- The basic trig limit $\lim_{h \rightarrow 0} \frac{\sin h}{h}$, and everything that follows it, depends on using **radian measure** for angles.
- We use radian measure because it is more natural in the math setting than other angle units.
- The basic trig limit in degree measure is

$$\lim_{h \rightarrow 0} \frac{\sindeg h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{h} = \lim_{h \rightarrow 0} \frac{\sin \frac{\pi}{180} h}{\frac{\pi}{180} h} \cdot \frac{\pi}{180} = \frac{\pi}{180}$$

- That crazy factor of $\pi/180$ will screw up every other calculus formula involving trig functions.
- For another example, try differentiating \sindeg . We'll take another look at that in section 3.5.

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.

The Derivatives of tan and sec

- The derivatives of other trig functions can now be found by writing them in terms of sin and cos and applying the quotient rule.
- For example,

$$\frac{d}{dx} \tan x$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\frac{d}{dx} \tan x = \frac{d}{dx} \frac{\sin x}{\cos x}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\frac{d}{dx} \tan x = \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x}\end{aligned}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x}\end{aligned}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

- The answer can be expressed in numerous ways depending on which trig identities are applied.

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

- The answer can be expressed in numerous ways depending on which trig identities are applied.
- The derivative of $\sec x$ is similar:

$$\frac{d}{dx} \sec x$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

- The answer can be expressed in numerous ways depending on which trig identities are applied.
- The derivative of $\sec x$ is similar:

$$\frac{d}{dx} \sec x = \frac{d}{dx} \frac{1}{\cos x}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

- The answer can be expressed in numerous ways depending on which trig identities are applied.
- The derivative of $\sec x$ is similar:

$$\frac{d}{dx} \sec x = \frac{d}{dx} \frac{1}{\cos x} = \frac{\cos x \cdot 0 - 1(-\sin x)}{\cos^2 x}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

- The answer can be expressed in numerous ways depending on which trig identities are applied.
- The derivative of $\sec x$ is similar:

$$\frac{d}{dx} \sec x = \frac{d}{dx} \frac{1}{\cos x} = \frac{\cos x \cdot 0 - 1(-\sin x)}{\cos^2 x} = \frac{\sin x}{\cos^2 x}$$

The Derivatives of \tan and \sec

- The derivatives of other trig functions can now be found by writing them in terms of \sin and \cos and applying the quotient rule.
- For example,

$$\begin{aligned}\frac{d}{dx} \tan x &= \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \sin' x - \sin x \cos' x}{\cos^2 x} \\ &= \frac{\cos x \cos x + \sin x \sin x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

- The answer can be expressed in numerous ways depending on which trig identities are applied.
- The derivative of $\sec x$ is similar:

$$\frac{d}{dx} \sec x = \frac{d}{dx} \frac{1}{\cos x} = \frac{\cos x \cdot 0 - 1(-\sin x)}{\cos^2 x} = \frac{\sin x}{\cos^2 x} = \sec x \tan x$$

Table of Trig Derivatives

The traditional formulas for the derivatives of the basic trig functions are as follows:

$\sin' x = \cos x$	
$\cos' x = -\sin x$	

You should memorize that table as well as you can.

Table of Trig Derivatives

The traditional formulas for the derivatives of the basic trig functions are as follows:

$\sin' x = \cos x$	
$\cos' x = -\sin x$	
$\tan' x = \sec^2 x$	

You should memorize that table as well as you can.

Table of Trig Derivatives

The traditional formulas for the derivatives of the basic trig functions are as follows:

$\sin' x = \cos x$	
$\cos' x = -\sin x$	
$\tan' x = \sec^2 x$	

You should memorize that table as well as you can.

Table of Trig Derivatives

The traditional formulas for the derivatives of the basic trig functions are as follows:

$\sin' x = \cos x$	
$\cos' x = -\sin x$	$\sec' x = \sec x \tan x$
$\tan' x = \sec^2 x$	

You should memorize that table as well as you can.

Table of Trig Derivatives

The traditional formulas for the derivatives of the basic trig functions are as follows:

$\sin' x = \cos x$	$\csc' x = -\csc x \cot x$
$\cos' x = -\sin x$	$\sec' x = \sec x \tan x$
$\tan' x = \sec^2 x$	$\cot' x = -\csc^2 x$

You should memorize that table as well as you can.

Examples

1. Find the limit $\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\sin \theta}$.
2. Find the derivative of $y = \frac{\sec x}{1 + \tan x}$.
3. Find equations of the tangent and normal lines to $y = \sec x$ at the point $(\pi/3, 2)$.
4. Find the numbers x at which the tangent to the curve $y = \sin x + \cos x$ is horizontal.

Now you should work on Problem Set 2.4. After you have finished it, you should try the following additional exercises from Section 2.4:

2.4 C-level: 1–16, 21–24, 25–26, 27–30, 33–34, 39–50;

B-level: 17–20, 31, 32, 35–38;

A-level: 51–52, 53–58