

MATH 110 Lecture 3.7

Optimization Problems

Edward Doolittle

Thursday, March 19, 2026

Department of Indigenous Knowledge and Science
First Nations University of Canada

Optimization Problems

Steps in Solving Optimization Problems

Examples from Algebra

Examples from Geometry and Physics

Examples from Business and Economics

Examples and Exercises

Optimization Problems

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.
- Steps required to solve such problems typically include:

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.
- Steps required to solve such problems typically include:
 1. **Understand the problem.** You will need to read carefully and understand the statement of the problem before you can solve it. You may have to return to this step repeatedly as you move through the other steps.

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.
- Steps required to solve such problems typically include:
 1. **Understand the problem.** You will need to read carefully and understand the statement of the problem before you can solve it. You may have to return to this step repeatedly as you move through the other steps.
 2. **Draw a diagram.** Essential for geometry-related problems.

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.
- Steps required to solve such problems typically include:
 1. **Understand the problem.** You will need to read carefully and understand the statement of the problem before you can solve it. You may have to return to this step repeatedly as you move through the other steps.
 2. **Draw a diagram.** Essential for geometry-related problems.
 3. **Introduce notation.** Sometimes it is given, usually you need to introduce it yourself.

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.
- Steps required to solve such problems typically include:
 1. **Understand the problem.** You will need to read carefully and understand the statement of the problem before you can solve it. You may have to return to this step repeatedly as you move through the other steps.
 2. **Draw a diagram.** Essential for geometry-related problems.
 3. **Introduce notation.** Sometimes it is given, usually you need to introduce it yourself.
 4. **Find a formula for Q , the quantity to be maximized.**

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.
- Steps required to solve such problems typically include:
 1. **Understand the problem.** You will need to read carefully and understand the statement of the problem before you can solve it. You may have to return to this step repeatedly as you move through the other steps.
 2. **Draw a diagram.** Essential for geometry-related problems.
 3. **Introduce notation.** Sometimes it is given, usually you need to introduce it yourself.
 4. **Find a formula for Q , the quantity to be maximized.**
 5. **Find formulas for constraints.** Use them to reduce Q to a function of one variable.

Steps in Solving Optimization Problems

- Optimization problems are word problems in which your goal is to maximize or minimize some quantity: time, cost, profit, etc.
- Steps required to solve such problems typically include:
 1. **Understand the problem.** You will need to read carefully and understand the statement of the problem before you can solve it. You may have to return to this step repeatedly as you move through the other steps.
 2. **Draw a diagram.** Essential for geometry-related problems.
 3. **Introduce notation.** Sometimes it is given, usually you need to introduce it yourself.
 4. **Find a formula for Q , the quantity to be maximized.**
 5. **Find formulas for constraints.** Use them to reduce Q to a function of one variable.
 6. **Use the methods of 3.1, 3.3, 3.4 to find the *absolute* maximum or minimum of Q .** Check end behaviour!

Skills Required

- In nearly every problem, four major steps are required:

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .
 3. Calculus: Find the absolute maximum or minimum value of Q .

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .
 3. Calculus: Find the absolute maximum or minimum value of Q .
 4. Interpretation: Re-read the question and answer it in words.

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .
 3. Calculus: Find the absolute maximum or minimum value of Q .
 4. Interpretation: Re-read the question and answer it in words.
- The skills required in each case are different:

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .
 3. Calculus: Find the absolute maximum or minimum value of Q .
 4. Interpretation: Re-read the question and answer it in words.
- The skills required in each case are different:
 1. Analysis: understanding, visualization, and conversion of words to symbols.

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .
 3. Calculus: Find the absolute maximum or minimum value of Q .
 4. Interpretation: Re-read the question and answer it in words.
- The skills required in each case are different:
 1. Analysis: understanding, visualization, and conversion of words to symbols.
 2. Algebra: algebraic manipulations.

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .
 3. Calculus: Find the absolute maximum or minimum value of Q .
 4. Interpretation: Re-read the question and answer it in words.
- The skills required in each case are different:
 1. Analysis: understanding, visualization, and conversion of words to symbols.
 2. Algebra: algebraic manipulations.
 3. Calculus: essentially the same skills as for curve sketching.

Skills Required

- In nearly every problem, four major steps are required:
 1. Analysis: Go from words to formulas for objective Q and constraint(s) C .
 2. Algebra: Eliminate dependent variables in Q by solving C .
 3. Calculus: Find the absolute maximum or minimum value of Q .
 4. Interpretation: Re-read the question and answer it in words.
- The skills required in each case are different:
 1. Analysis: understanding, visualization, and conversion of words to symbols.
 2. Algebra: algebraic manipulations.
 3. Calculus: essentially the same skills as for curve sketching.
 4. Interpretation: understanding and conversion of symbols to words.

Advice

- Optimization problems are some of the most useful applications of calculus we will study.

Advice

- Optimization problems are some of the most useful applications of calculus we will study.
- Optimization problems are difficult because they test a wide spectrum of skills and abilities.

Advice

- Optimization problems are some of the most useful applications of calculus we will study.
- Optimization problems are difficult because they test a wide spectrum of skills and abilities.
- With optimization problems, as with all word problems, lots of practice is necessary to build skill.

Advice

- Optimization problems are some of the most useful applications of calculus we will study.
- Optimization problems are difficult because they test a wide spectrum of skills and abilities.
- With optimization problems, as with all word problems, lots of practice is necessary to build skill.
- The problems we will look at divide into categories (dealing with numbers, geometry with rectangles, geometry with circles and rectangles, three dimensional geometry with boxes, trigonometry, etc.)

Advice

- Optimization problems are some of the most useful applications of calculus we will study.
- Optimization problems are difficult because they test a wide spectrum of skills and abilities.
- With optimization problems, as with all word problems, lots of practice is necessary to build skill.
- The problems we will look at divide into categories (dealing with numbers, geometry with rectangles, geometry with circles and rectangles, three dimensional geometry with boxes, trigonometry, etc.)
- Once you have done a few problems in each category, you will be able to solve others in that category more quickly.

Advice

- Optimization problems are some of the most useful applications of calculus we will study.
- Optimization problems are difficult because they test a wide spectrum of skills and abilities.
- With optimization problems, as with all word problems, lots of practice is necessary to build skill.
- The problems we will look at divide into categories (dealing with numbers, geometry with rectangles, geometry with circles and rectangles, three dimensional geometry with boxes, trigonometry, etc.)
- Once you have done a few problems in each category, you will be able to solve others in that category more quickly.
- Start with easier problems to build skill and then work through as many of the harder problems as you can.

Sum is Constraint, Product is Objective

- Find two numbers, the sum of which is 23 and the product of which is maximum.

Sum is Constraint, Product is Objective

- Find two numbers, the sum of which is 23 and the product of which is maximum.
- We want two numbers, so let's call them x and y .

Sum is Constraint, Product is Objective

- Find two numbers, the sum of which is 23 and the product of which is maximum.
- We want two numbers, so let's call them x and y .
- The sum of the two numbers is 23, so $x + y = 23$. That is the constraint in the problem.

Sum is Constraint, Product is Objective

- Find two numbers, the sum of which is 23 and the product of which is maximum.
- We want two numbers, so let's call them x and y .
- The sum of the two numbers is 23, so $x + y = 23$. That is the constraint in the problem.
- The product should be a maximum, so our objective is to maximize $Q = xy$.

Sum is Constraint, Product is Objective

- Find two numbers, the sum of which is 23 and the product of which is maximum.
- We want two numbers, so let's call them x and y .
- The sum of the two numbers is 23, so $x + y = 23$. That is the constraint in the problem.
- The product should be a maximum, so our objective is to maximize $Q = xy$.
- We use the constraint to eliminate one of the variables, e.g., $y = 23 - x$ so we can write $Q(x) = xy = x(23 - x)$.

Sum is Constraint, Product is Objective


- Find two numbers, the sum of which is 23 and the product of which is maximum.
- We want two numbers, so let's call them x and y .
- The sum of the two numbers is 23, so $x + y = 23$. That is the constraint in the problem.
- The product should be a maximum, so our objective is to maximize $Q = xy$.
- We use the constraint to eliminate one of the variables, e.g., $y = 23 - x$ so we can write $Q(x) = xy = x(23 - x)$.
- To maximize Q , we consider $Q'(x) = 23 - 2x$. The max will occur where $Q'(x) = 0$, or where $Q'(x)$ doesn't exist (but it always exists), or at an endpoint of the domain (there is no endpoint). Solving $Q'(x) = 0$ we have $23 - 2x = 0$ or $x = 23/2$.

Sum is Constraint, Product is Objective, ctd.

- We have found $Q(x) = xy = x(23 - x)$ has a critical point at $x = 23/2$.


Sum is Constraint, Product is Objective, ctd.

- We have found $Q(x) = xy = x(23 - x)$ has a critical point at $x = 23/2$.
- We need to check that $Q(x)$ has a **maximum** at $x = 23/2$.




Sum is Constraint, Product is Objective, ctd.

- We have found $Q(x) = xy = x(23 - x)$ has a critical point at $x = 23/2$.
- We need to check that $Q(x)$ has a **maximum** at $x = 23/2$.
- $Q'(x) > 0$ for $x < 23/2$, and $Q'(x) < 0$ for $x > 23/2$, so $Q(x)$ increases to $x = 23/2$ then decreases.




Sum is Constraint, Product is Objective, ctd.

- We have found $Q(x) = xy = x(23 - x)$ has a critical point at $x = 23/2$.
- We need to check that $Q(x)$ has a **maximum** at $x = 23/2$.
- $Q'(x) > 0$ for $x < 23/2$, and $Q'(x) < 0$ for $x > 23/2$, so $Q(x)$ increases to $x = 23/2$ then decreases.
- By the first derivative test, Q has an absolute maximum at $x = 23/2$.



Sum is Constraint, Product is Objective, ctd.

- We have found $Q(x) = xy = x(23 - x)$ has a critical point at $x = 23/2$.
- We need to check that $Q(x)$ has a **maximum** at $x = 23/2$.
- $Q'(x) > 0$ for $x < 23/2$, and $Q'(x) < 0$ for $x > 23/2$, so $Q(x)$ increases to $x = 23/2$ then decreases.
- By the first derivative test, Q has an absolute maximum at $x = 23/2$.
- Now we can answer the problem: The two numbers which add to 23 and have the largest possible product are $x = 23/2$ and $y = 23 - x = 23/2$.



Sum is Objective, Product is Constraint

- Find two positive numbers, the product of which is 100 and the sum of which is minimum.

Sum is Objective, Product is Constraint

- Find two positive numbers, the product of which is 100 and the sum of which is minimum.
- Again, let our two numbers be x and y .

Sum is Objective, Product is Constraint

- Find two positive numbers, the product of which is 100 and the sum of which is minimum.
- Again, let our two numbers be x and y .
- We have the constraint $xy = 100$.

Sum is Objective, Product is Constraint

- Find two positive numbers, the product of which is 100 and the sum of which is minimum.
- Again, let our two numbers be x and y .
- We have the constraint $xy = 100$.
- We have the objective to minimize $Q = x + y$.

Sum is Objective, Product is Constraint

- Find two positive numbers, the product of which is 100 and the sum of which is minimum.
- Again, let our two numbers be x and y .
- We have the constraint $xy = 100$.
- We have the objective to minimize $Q = x + y$.
- We also have the constraint that x and y must be positive; in symbols, $x > 0$ and $y > 0$.

Sum is Objective, Product is Constraint

- Find two positive numbers, the product of which is 100 and the sum of which is minimum.
- Again, let our two numbers be x and y .
- We have the constraint $xy = 100$.
- We have the objective to minimize $Q = x + y$.
- We also have the constraint that x and y must be positive; in symbols, $x > 0$ and $y > 0$.
- Eliminating the variable $y = 100/x$, we reduce the problem to minimizing $Q(x) = x + y = x + 100/x$, $x > 0$. Note that the domain of Q is not the whole real line, only the half line $(0, \infty)$.

Sum is Objective, Product is Constraint

- Find two positive numbers, the product of which is 100 and the sum of which is minimum.
- Again, let our two numbers be x and y .
- We have the constraint $xy = 100$.
- We have the objective to minimize $Q = x + y$.
- We also have the constraint that x and y must be positive; in symbols, $x > 0$ and $y > 0$.
- Eliminating the variable $y = 100/x$, we reduce the problem to minimizing $Q(x) = x + y = x + 100/x$, $x > 0$. Note that the domain of Q is not the whole real line, only the half line $(0, \infty)$.
- The derivative is $Q'(x) = 1 - 100/x^2$, with critical points where $Q'(x) = 0$, $1 - 100/x^2 = 0$, $x = \pm 10$.

Sum is Objective, Product is Constraint


- Find two positive numbers, the product of which is 100 and the sum of which is minimum.
- Again, let our two numbers be x and y .
- We have the constraint $xy = 100$.
- We have the objective to minimize $Q = x + y$.
- We also have the constraint that x and y must be positive; in symbols, $x > 0$ and $y > 0$.
- Eliminating the variable $y = 100/x$, we reduce the problem to minimizing $Q(x) = x + y = x + 100/x$, $x > 0$. Note that the domain of Q is not the whole real line, only the half line $(0, \infty)$.
- The derivative is $Q'(x) = 1 - 100/x^2$, with critical points where $Q'(x) = 0$, $1 - 100/x^2 = 0$, $x = \pm 10$.
- We throw away $x = -10$ because it is outside of the domain of Q .

Sum is Objective, Product is Constraint, ctd.

- Once again, we need to check that we have an absolute minimum at $x = 10$.


Sum is Objective, Product is Constraint, ctd.

- Once again, we need to check that we have an absolute minimum at $x = 10$.
- We have $Q'(x) = 1 - 100/x^2 < 0$ for $x < 10$, and $Q'(x) > 0$ for $x > 10$.




Sum is Objective, Product is Constraint, ctd.

- Once again, we need to check that we have an absolute minimum at $x = 10$.
- We have $Q'(x) = 1 - 100/x^2 < 0$ for $x < 10$, and $Q'(x) > 0$ for $x > 10$.
- So $Q(x)$ decreases to $x = 10$ and then increases.




Sum is Objective, Product is Constraint, ctd.

- Once again, we need to check that we have an absolute minimum at $x = 10$.
- We have $Q'(x) = 1 - 100/x^2 < 0$ for $x < 10$, and $Q'(x) > 0$ for $x > 10$.
- So $Q(x)$ decreases to $x = 10$ and then increases.
- By the First Derivative Test, Q has an absolute minimum at $x = 10$.



Sum is Objective, Product is Constraint, ctd.

- Once again, we need to check that we have an absolute minimum at $x = 10$.
- We have $Q'(x) = 1 - 100/x^2 < 0$ for $x < 10$, and $Q'(x) > 0$ for $x > 10$.
- So $Q(x)$ decreases to $x = 10$ and then increases.
- By the First Derivative Test, Q has an absolute minimum at $x = 10$.
- The answer to the question is: the two numbers, the product of which is 100 and the sum of which is a minimum, are $x = 10$ and $y = 100/x = 10$.




Plane Geometry

- Find a rectangle with perimeter 100 m for which the area is a maximum.


Plane Geometry

- Find a rectangle with perimeter 100 m for which the area is a maximum.
- To understand this problem better, we should draw a picture.




Plane Geometry

- Find a rectangle with perimeter 100 m for which the area is a maximum.
- To understand this problem better, we should draw a picture.
- We introduce labels for quantities.




Plane Geometry

- Find a rectangle with perimeter 100 m for which the area is a maximum.
- To understand this problem better, we should draw a picture.
- We introduce labels for quantities.
- We use the geometry of the diagram to label the other sides.




Plane Geometry

- Find a rectangle with perimeter 100 m for which the area is a maximum.
- To understand this problem better, we should draw a picture.
- We introduce labels for quantities.
- We use the geometry of the diagram to label the other sides.
- Geometry tells us that the area is $A = xy$ and the perimeter is $P = x + y + x + y = 2x + 2y$.




Plane Geometry

- Find a rectangle with perimeter 100 m for which the area is a maximum.
- To understand this problem better, we should draw a picture.
- We introduce labels for quantities.
- We use the geometry of the diagram to label the other sides.
- Geometry tells us that the area is $A = xy$ and the perimeter is
$$P = x + y + x + y = 2x + 2y.$$
- Our constraint is $P = 100$, i.e.,
$$x + y = 50.$$



Plane Geometry


- Find a rectangle with perimeter 100 m for which the area is a maximum.
- To understand this problem better, we should draw a picture.
- We introduce labels for quantities.
- We use the geometry of the diagram to label the other sides.
- Geometry tells us that the area is $A = xy$ and the perimeter is
$$P = x + y + x + y = 2x + 2y.$$
- Our constraint is $P = 100$, i.e.,
$$x + y = 50.$$
- Our objective is to maximize $A = xy$.



- Find the point on the line $y = 4x + 7$ that is closest to the origin.


Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0, 0)$ and $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0,0)$ and $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0, 0)$ and $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0,0)$ and $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0,0)$ and $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0, 0)$ and $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0,0)$ and $(x, 4x + 7)$.




Analytic Geometry

- Find the point on the line $y = 4x + 7$ that is closest to the origin.
- Again, we draw a diagram, including both the line and the origin.
- We introduce a variable x to describe a point on the line. Then the coordinates of the point on the line are $(x, 4x + 7)$.
- We try to imagine how we might find the point that is closest to the origin as x moves. We need a formula for the length of the line segment connecting $(0,0)$ and $(x, 4x + 7)$.




Analytic Geometry, ctd

- Recall that the distance between the two points $(0,0)$ and $(x, 4x+7)$ is $d(x) = \sqrt{x^2 + (4x+7)^2}$ by the Pythagorean theorem.




Analytic Geometry, ctd

- Recall that the distance between the two points $(0,0)$ and $(x, 4x + 7)$ is $d(x) = \sqrt{x^2 + (4x + 7)^2}$ by the Pythagorean theorem.
- The usual trick in these problems is to minimize the function $D(x) = (d(x))^2$. Then we don't have to worry about square roots.




Analytic Geometry, ctd

- Recall that the distance between the two points $(0,0)$ and $(x, 4x+7)$ is $d(x) = \sqrt{x^2 + (4x+7)^2}$ by the Pythagorean theorem.
- The usual trick in these problems is to minimize the function $D(x) = (d(x))^2$. Then we don't have to worry about square roots.
- We have $D(x) = x^2 + (4x+7)^2 = 17x^2 + 56x + 49$.




Analytic Geometry, ctd

- Recall that the distance between the two points $(0,0)$ and $(x, 4x+7)$ is $d(x) = \sqrt{x^2 + (4x+7)^2}$ by the Pythagorean theorem.
- The usual trick in these problems is to minimize the function $D(x) = (d(x))^2$. Then we don't have to worry about square roots.
- We have $D(x) = x^2 + (4x+7)^2 = 17x^2 + 56x + 49$.
- $D'(x) = 34x + 56$.




Analytic Geometry, ctd

- Recall that the distance between the two points $(0,0)$ and $(x, 4x+7)$ is $d(x) = \sqrt{x^2 + (4x+7)^2}$ by the Pythagorean theorem.
- The usual trick in these problems is to minimize the function $D(x) = (d(x))^2$. Then we don't have to worry about square roots.
- We have $D(x) = x^2 + (4x+7)^2 = 17x^2 + 56x + 49$.
- $D'(x) = 34x + 56$.
- $D'(x) = 0$ implies $x = -56/34 = -28/17$.




Analytic Geometry, ctd

- Recall that the distance between the two points $(0,0)$ and $(x, 4x+7)$ is $d(x) = \sqrt{x^2 + (4x+7)^2}$ by the Pythagorean theorem.
- The usual trick in these problems is to minimize the function $D(x) = (d(x))^2$. Then we don't have to worry about square roots.
- We have $D(x) = x^2 + (4x+7)^2 = 17x^2 + 56x + 49$.
- $D'(x) = 34x + 56$.
- $D'(x) = 0$ implies $x = -56/34 = -28/17$.
- The first derivative test tells us there's an absolute min at $x = -28/17$.



Analytic Geometry, ctd

- Recall that the distance between the two points $(0,0)$ and $(x, 4x+7)$ is $d(x) = \sqrt{x^2 + (4x+7)^2}$ by the Pythagorean theorem.
- The usual trick in these problems is to minimize the function $D(x) = (d(x))^2$. Then we don't have to worry about square roots.
- We have $D(x) = x^2 + (4x+7)^2 = 17x^2 + 56x + 49$.
- $D'(x) = 34x + 56$.
- $D'(x) = 0$ implies $x = -56/34 = -28/17$.
- The first derivative test tells us there's an absolute min at $x = -28/17$.
- The point on the line that is closest to the origin is $(x,y) = (-28/17, 4(-28/17) + 7)$.



- If 1200 cm^2 of material is available to make a box with a square base and an open top, find the largest possible volume of the box.

- A boat leaves a dock at 2 pm and travels south at a speed of 20 km/h. Another boat has been heading east at 15km/h and reaches the dock at 3pm. At what time were the boats closest together?

Optimizing Revenue

- The demand function $p(x)$ is the price per unit that a company can charge if it wants to sell x units.

Optimizing Revenue

- The demand function $p(x)$ is the price per unit that a company can charge if it wants to sell x units.
- The revenue function at sales level x is $R(x) = x \cdot p(x)$.

Optimizing Revenue

- The demand function $p(x)$ is the price per unit that a company can charge if it wants to sell x units.
- The revenue function at sales level x is $R(x) = x \cdot p(x)$.
- A baseball team plays in a stadium that holds 55,000 spectators. With ticket prices at \$10, the average attendance had been 27,000. When ticket prices were lowered to \$8, the average attendance rose to 33,000.

Optimizing Revenue

- The demand function $p(x)$ is the price per unit that a company can charge if it wants to sell x units.
- The revenue function at sales level x is $R(x) = x \cdot p(x)$.
- A baseball team plays in a stadium that holds 55,000 spectators. With ticket prices at \$10, the average attendance had been 27,000. When ticket prices were lowered to \$8, the average attendance rose to 33,000.
 1. Find the demand function, assuming that it is linear.

Optimizing Revenue

- The demand function $p(x)$ is the price per unit that a company can charge if it wants to sell x units.
- The revenue function at sales level x is $R(x) = x \cdot p(x)$.
- A baseball team plays in a stadium that holds 55,000 spectators. With ticket prices at \$10, the average attendance had been 27,000. When ticket prices were lowered to \$8, the average attendance rose to 33,000.
 1. Find the demand function, assuming that it is linear.
 2. How should ticket prices be set to maximize revenue?

Optimizing Profit

- Profit is revenue minus cost at sales level x , i.e.,
$$P(x) = x \cdot p(x) - C(x).$$

Optimizing Profit

- Profit is revenue minus cost at sales level x , i.e.,
$$P(x) = x \cdot p(x) - C(x).$$
- A manufacturer has been selling 1000 television sets per week at \$450 each. A market survey indicates that for each \$10 rebate offered to the buyer, the number of sets sold will increase by 100 per week.

Optimizing Profit

- Profit is revenue minus cost at sales level x , i.e.,
$$P(x) = x \cdot p(x) - C(x).$$
- A manufacturer has been selling 1000 television sets per week at \$450 each. A market survey indicates that for each \$10 rebate offered to the buyer, the number of sets sold will increase by 100 per week.
 1. Find the demand function.

Optimizing Profit

- Profit is revenue minus cost at sales level x , i.e.,
$$P(x) = x \cdot p(x) - C(x).$$
- A manufacturer has been selling 1000 television sets per week at \$450 each. A market survey indicates that for each \$10 rebate offered to the buyer, the number of sets sold will increase by 100 per week.
 1. Find the demand function.
 2. How large a rebate should the company offer to maximize its revenue?

Optimizing Profit

- Profit is revenue minus cost at sales level x , i.e.,
$$P(x) = x \cdot p(x) - C(x).$$
- A manufacturer has been selling 1000 television sets per week at \$450 each. A market survey indicates that for each \$10 rebate offered to the buyer, the number of sets sold will increase by 100 per week.
 1. Find the demand function.
 2. How large a rebate should the company offer to maximize its revenue?
 3. If its weekly cost function is $C(x) = 68,000 + 150x$, how should the manufacturer set the size of the rebate in order to maximize its profit?

Examples and Exercises

Examples

1. A farmer wants to fence an area 1.5 million square feet in rectangular field and then divide it in half with a fence parallel to one of the sides. How can he do this so as to minimize the cost of the fence?
2. Find the points on the ellipse $4x^2 + y^2 = 4$ that are farthest away from the point $(0, 1)$.
3. A right circular cylinder is inscribed in a sphere of radius r . Find the largest possible volume of such a cylinder.
4. The illumination of an object by a light source is directly proportional to the strength of the source and inversely proportional to the square of the distance from the source. If two light sources, one three times as strong as the other, are placed 10 ft apart, where should the object be placed on the line between the sources so as to receive the least illumination?

Now you should work on Problem Set 3.7. After you have finished it, you should try the following additional exercises from Section 3.7:

3.7 C-level: 1–64;

B-level: 65–67, 69–70;

A-level: 71–80