

Quantum Computing Cheatsheet

Edoardo Riggio

June 6, 2022

Quantum Computing - S.P. 2022
Computer Science
Università della Svizzera Italiana, Lugano

Contents

1	What is Quantum Informatics	2
1.1	Information and Physics	2
1.2	Second Law of Thermodynamics	2
1.3	The Stern/Gerlach Experiment	2
1.4	Superposition	4
2	Glossary	4
2.1	Magnetic Dipole Moment	4

1 What is Quantum Informatics

1.1 Information and Physics

Experience, observation, and physical discourse are in the form of information.
"It from Bit" – John Wheeler

Information representation, processing, and transmission are physical processes.
"Information is physical" – Rolf Landauer

The representation of a bit must be physical. Moreover, digitalization comes very naturally with **quantization**. In classical physics, digitalization has to be enforced somehow (e.g., switched).

1.2 Second Law of Thermodynamics


The second law of thermodynamics states that, in a closed system, entropy does not increase.

Entropy can be defined as a measure of disorder. Given n binary memory cells containing random bits, if we erase all of the bits – i.e., set them to 0 – then the entropy in the set of memory cells drops.

1.3 The Stern/Gerlach Experiment

This experiment was proposed in 1921 by Otto Stern and later carried out in 1922 by Walther Gerlach.


This is one of the most important experiments to understand the structure and properties of the basic building block of quantum information processing, the **Qbit**.



This experiment consisted of the measurement of the **magnetic dipole moment** of silver atoms. These silver atoms are sent as a stream (2) coming from an oven (1). Each atom is deflected from the path through an inhomogeneous magnetic field (3), and each atom is deflected from the path (5). This deflection is proportional to its dipole in the direction of the magnets.

This experiment revealed no detection in the middle of the screen (4) but rather two sharp peaks at equal distances from the center (5). The quantity measured by the experiment is known in quantum mechanics as **spin**.


In the case of a single measurement, for example, in the z -direction, it will result in two identical rays.



If the exact measurement is repeated for only one of the rays – say z_+ , then all the atoms are deflected again in the $+$ direction.



Finally, if the magnet is rotated and a x -direction measurement of the z_+ ray is made, another z -direction measurement, a 50-50 distribution arises again. This puts the stability and the independence of the properties in question.



1.4 Superposition

Quantum superposition is a fundamental principle of quantum mechanics. It states that any two – or more – quantum states can be added together, and the result will be another valid quantum state.


The question of whether a silver atom is in the state $|z_-\rangle$ or in the state $|z_+\rangle$ are complementary to one another. They can be regarded as two answers to the same question – i.e., the Z measurement.

If after performing an X measurement, we want to know whether the silver atom is in a state $|x_+\rangle$ or $|x_-\rangle$. Both are equal superpositions

$$|x_+\rangle = \frac{1}{\sqrt{2}}|z_+\rangle + \frac{1}{\sqrt{2}}|z_-\rangle$$

$$|x_-\rangle = \frac{1}{\sqrt{2}}|z_+\rangle - \frac{1}{\sqrt{2}}|z_-\rangle$$

No matter if we obtain one measurement or the other in the Z measurement, the X -measurement either $|x_+\rangle$ or $|x_-\rangle$ with equal probability. This is also known as a **quantum jump**.



2 Glossary

2.1 Magnetic Dipole Moment

The magnetic dipole moment is a vector that represents the strength and orientation of a magnet or other object that produces a magnetic field (e.g., an electron).