

# Bayesian inference on Poisson-Gamma random measures in Spatial statistics

Edoardo Tarcisio Visconti

March 2024

## Contents

<b>Abstract</b>	<b>2</b>
<b>1 Random measures and their simulation</b>	<b>3</b>
1.1 Poisson random measure . . . . .	3
1.2 Completely Random Measures and the Gamma Process . . . . .	5
1.3 Simulation of the Gamma Process . . . . .	7
1.4 Cox process and the Poisson-Gamma random measure . . . . .	9
<b>2 Bayesian Inference</b>	<b>13</b>
2.1 The Bayesian framework . . . . .	13
2.2 Markov chain Monte Carlo . . . . .	13
2.2.1 Metropolis-Hastings algorithm . . . . .	14
2.2.2 Gibbs Sampler . . . . .	15
<b>3 Bayesian inference for Poisson-Gamma Random measures</b>	<b>16</b>
3.1 On the discrete Gamma Process . . . . .	16
3.2 Sampling from the discrete Gamma Process . . . . .	18
3.3 The Metropolis within Gibbs algorithm . . . . .	19
3.4 The algorithm with more observations (stationary case) . . . . .	22
3.5 The algorithm with more observations (time-varying case) . . . . .	25
<b>4 An application: inference on forest fires in the Amazon Forest</b>	<b>26</b>
4.1 The stationary case . . . . .	26
4.2 The time-varying case . . . . .	29
<b>5 Further work</b>	<b>32</b>
<b>References</b>	<b>32</b>

## Abstract

In this bachelor’s final project, we begin by providing some background on Random Measures, with a focus on Poisson Point Processes, Completely Random Measures, and Gamma Processes. Then, the Poisson-Gamma Random Measure, a shot noise Cox process introduced by Wolpert and Ickstadt in [4], is presented along with theorems for its simulation. Additionally, we offer illustrative examples along with numerical methodologies to facilitate implementation.

This stochastic process is well-suited for explaining and modeling point processes occurring on subsets of  $\mathbb{R}^2$  where clustering is notable, such as in instances of forest fires, taxi pickups, and violent crimes.

In the second section, a concise overview of Markov Chain Monte Carlo (MCMC) algorithms is provided, with an emphasis on methodologies relevant to the subsequent discussion.

In the third segment, we present our contribution to the topic: the development of a Bayesian inference algorithm tailored specifically for Poisson-Gamma Random Measures.

A more generalized model for Bayesian inference on such measures has already been proposed in [4]. Nonetheless, we introduce an alternative approach based on discretizing the underlying Gamma process, namely a discrete Gamma process. We are confident that our model offers computational advantages, increased stability, and enables sampling from the posterior while observing multiple realizations of the same process simultaneously (unlike in [4], where only one realization is considered). Additionally, we believe that a partial result can be proved to show the efficacy of our model even in cases where the latent Gamma process is non-discrete.

Finally, we demonstrate the practical application of our approach using georeferenced wildfire data. This example serves to showcase the properties of the model and inference techniques, both in a constant scale and on a time-varying scale.

The code and the dataset are available here: **Project’s Github**.

# 1 Random measures and their simulation

## 1.1 Poisson random measure

This subsection offers a brief but rigorous overview the Poisson random measure, commonly referred to as the Poisson point process.

**Notation:** given the space  $(E, \mathcal{E})$  we consider the space  $(\mathcal{M}, \mathcal{B}(\mathcal{M}))$  where

$$\mathcal{M} = \{\text{sigma-finite measures } \eta \text{ on } E \text{ such that } \forall B \in \mathcal{E}, \eta(B) \in \mathbb{N} \cup \{\infty\}\}$$

and  $\mathcal{B}(\mathcal{M})$  is the  $\sigma$ -algebra generated by the sets

$$\{\eta \in \mathcal{M}: \eta(B) = k\}$$

for  $B \subset E$  measurable and  $k \in \mathbb{N}$

**Definition 1.1.** A Point Process on  $(E, \mathcal{E})$  is a stochastic process

$$M = (M(B))_{B \in \mathcal{E}}$$

with values in  $\mathbb{N} \cup \{\infty\}$  such that  $M \in \mathcal{M}$  a.s.

Intuitively, A point process indicates how many points there are in each region  $B$  of the space, i.e. for fixed  $B$ , the random integer  $M(B)$  represents the number of points in  $B$ . The condition  $M \in \mathcal{M}$  a.s. ensures that all the numbers of points in different regions are compatible with each other

**Definition 1.2.** A Poisson Point Process (or Poisson random measure) on  $(E, \mathcal{E})$  with intensity measure  $\mu$  ( $\sim ppp(\mu)$ ) is a Point process  $M$  such that:

1. For all  $B_1, \dots, B_k \subset E$  measurable and disjoint,  $M(B_1), \dots, M(B_k)$  are independent
2. For all  $B \subset E$  measurable,  $M(B) \sim Pois(\mu(B))$

**Remark 1.1.** From 2. we deduce that  $\mathbb{E}[M(B)] = \mu(B) \forall B$  measurable, which is the expected number of points in the region  $B$ .

**Theorem 1.1.** Let  $M$  be a  $ppp(\mu)$  on  $(E, \mathcal{E})$ . Let  $\tau = M(E)$  i.e. the total number of points in  $E$ . There exists some random variables  $X_n \in E, n > 0$  such that:

$$M = \sum_{n=1}^{\tau} \delta_{X_n} \text{ a.s.}$$

**Theorem 1.2.** Let  $\mu(E) < \infty$  and  $Z, (X_i)_{i \geq 1}$  be independent random variables such that:

$$Z \sim Poisson(\mu(E)), \quad X_i \sim \frac{\mu(\cdot)}{\mu(E)}.$$

Then:  $M = \sum_{n=1}^Z \delta_{X_n}$  is a  $ppp(\mu)$  on  $E$

**Remark 1.2.** This theorem allows us to simulate  $M \sim ppp(\mu)$  with the following scheme:

---

**Algorithm 1** Simulation of  $M \sim ppp(\mu)$

---

- 1: Evaluate  $A = \mu(E)$
  - 2: Draw one sample  $z$  from  $Z \sim \text{Poisson}(A)$
  - 3: Draw  $z$  independent samples from  $X_i \sim \frac{\mu(\cdot)}{\mu(E)}$
- 

Where Step 3. can be performed with an Acceptance-Rejection Method or any other suitable methods.

As for standard random variables, Point processes can be characterised by their Laplace Functional:

**Definition 1.3.** Let  $M$  be a point process on  $(E, \mathcal{E})$ , for every  $u : E \rightarrow \mathbb{R}_+$  measurable we define the Laplace Functional as:

$$L_M(u) = \mathbb{E} \left[ \exp \left( - \int u(dx) M(dx) \right) \right]$$

**Remark 1.3.** We can interpret  $\int u(dx) M(dx)$  as:  $\sum_{x^*} \text{"points of } M" u(x^*)$  counted with multiplicities.

**Theorem 1.3.** (Characterization via Laplace Functional). Let  $\mu$  be a  $\sigma$ -finite measure on  $(E, \mathcal{E})$ . Let  $M$  be a point process on  $E$ . The following are equivalent:

1.  $M \sim ppp(\mu)$
2.  $u : E \rightarrow \mathbb{R}_+$  measurable:

$$L_M(u) = \exp \left( - \int_E \left( 1 - e^{-u(x)} \right) \mu(dx) \right)$$

This theorem can be seen as a definition for a Poisson Point Process.

Proof for Theorem 1.3 and other material on the Poisson Process is available in [1].

## 1.2 Completely Random Measures and the Gamma Process

In this subsection, a short and non rigorous overview of Completely random measures is provided. Then the Gamma Process is introduced.

For more read: [1], [2], [3].

**Definition 1.4.** A Completely Random Measure (CRM) on  $(E, \mathcal{E})$  is a random measure  $M$  such that, for all  $B_1, \dots, B_k \subset E$  measurable and disjoint,  $M(B_1), \dots, M(B_k)$  are independent.

Under general assumptions, a CRM  $M$  can be written as the sum of three independent random measures:

- a deterministic measure;
- an atomistic random measure  $M_D$  with fixed atoms  $\sum_{i \geq 1} S_i \delta_{\theta_i^*}$  where the points  $\theta_1^*, \theta_2^*, \dots$  are fixed in  $E$  and  $S_i$  are independent infinitely divisible positive random variables (like a Gamma RV);
- the ordinary component  $M_O$ , i.e. a discrete random measure  $M_O$  without fixed atoms.

For our purposes, we can restrict to a CRM  $M$  without deterministic component.

Then,  $M$  is characterized by the Laplace functional:

$$L_M(u) = \exp \left( - \sum_i \psi_{S_i}(v) f(\theta_i^*) - \int_{\mathbb{R}_{>0} \times E} (1 - e^{-vu(x)}) \mu(dvdx) \right) \quad (1)$$

where  $\psi_{S_i} = -\log(\mathbb{E}[\exp(-vS_j)])$  and  $\mu$ , the Levy measure, is a  $\sigma$ -finite diffuse measure on the Borel sets of  $\mathbb{R}_{>0} \times E$ , such that, for every bounded measurable set  $A \subset E$ , it holds:

$$\int_{\mathbb{R}_{>0} \times A} \min(1, v) \mu(dvdx) < \infty$$

We will write:

$$M \sim CRM(\mu, \underline{\theta}^*, \Psi)$$

In the following, we will allow  $\mu$  to have a deterministic component, i.e.  $\mu = \mu(dv d\theta) + \sum_{\theta_i^* \in \underline{\theta}^*} \mu(dv, \theta_i^*)$  and accordingly, we can write  $M \sim (\mu, \emptyset)$  if

$$L_M(u) = \exp \left( - \int_{\mathbb{R}_{>0} \times E} (1 - e^{-vu(x)}) \mu(dvdx) \right). \quad (2)$$

Note that this does not necessarily mean that  $M$  has no fixed atoms, since they may arise from  $\mu$ . The atoms that arise from  $\mu$ , if they exist, they have necessarily infinite divisible weights.

**Theorem 1.4.** If  $M$  is a CRM with Laplace functional (2) then:

$$M(A) = \int_{\mathbb{R}_{>0} \times A} v N(dv dx) = \sum_{j \geq 1} v_j \delta_{x_j}(A)$$

for a suitable Poisson random measure  $N = \sum_{j \geq 1} \delta_{(v_j, x_j)}$  on  $\mathbb{R}_{>0} \times E$  with intensity measure  $\mu$ .

**Definition 1.5.** A Homogeneous completely random measure of parameter  $(\nu, H)$ , for short hCRM $(\nu, H)$ , is a CRM with Laplace functional (2) in the special case:

$$\mu(dv dx) = \nu(dv) H(dx),$$

where  $H$  is a boundedly finite measure on  $E$  and  $\nu$  a  $\sigma$ -finite measure on  $\mathbb{R}_{>0}$  such that

$$\int_{\mathbb{R}_{>0} \times A} \min(1, v) \nu(dv) < \infty$$

**Remark 1.4.** A Poisson random measure (or Poisson point process)  $M$  on  $E$  with intensity measure  $H$ ,  $M \sim ppp(H)$ , can be seen as special case of hCRM where the Levy intensity is  $\nu(dv) = \delta_1(dv)$ . Then the Levy measure is  $\mu(dv dx) = \delta_1(dv) H(dx)$ .

**Definition 1.6.** A Gamma random measure (or Gamma process) on  $E$  with base measure  $H$  and rate  $\beta > 0$ , written  $\mathcal{G}aP(H, \beta)$ , is a hCRM with Levy intensity  $\nu(dv) = v^{-1} e^{-\beta v} dv$ . In this case the Levy measure is  $\mu(dv dx) = v^{-1} e^{-\beta v} dv H(dx)$  and the Laplace functional (2) reduces to:

$$L_M(f) = \exp \left( - \int_E \log \left( 1 + \frac{f(x)}{\beta} \right) H(dx) \right).$$

**Corollary 1.1.** From Theorem 1.4, if  $H$  is boundedly finite and  $W \sim \mathcal{G}aP(H, \beta)$ , then:

$$W(A) = \int_{\mathbb{R}_{>0} \times A} v N(dv dx) = \sum_{j \geq 1} v_j \delta_{x_j}(A)$$

where  $N = \sum_{j \geq 1} v_j \delta_{(v_j, x_j)}$  is a Poisson Process on  $\mathbb{R}_{>0} \times E$  with intensity measure  $\mu(dv dx) = v^{-1} e^{-\beta v} dv H(dx)$ .

From 1.6, we can generalize the Gamma Process to non-constant  $\beta(x) > 0$ . Note that now this process is no more a hCRM as the Levy measure does not factorise.

**Definition 1.7.** A generalised Gamma process (or non-homogeneous Gamma process)  $W$  with parameters  $H$  and  $\beta(x) > 0$ , written  $W \sim \mathcal{G}aP(H(dx), \beta(x))$  is characterised by the Laplace functional:

$$L_W(f) = \exp \left( - \int_E \log \left( 1 + \frac{f(x)}{\beta(x)} \right) H(dx) \right).$$

### 1.3 Simulation of the Gamma Process

Let's see now how to simulate a generalised Gamma process, where  $E_1(t) = \int_t^\infty e^{-u} u^{-1} du$ :

**Theorem 1.5.** Let  $\alpha(x) \geq 0$  and  $\beta(x) > 0$  be measurable functions on the space  $E$ . Let  $\{\sigma_m\}_{m \in \mathbb{N}}$  be identical distributed draws for any probability  $\Pi(dx)$  on  $E$ , and let  $\{\tau_m\}_{m \in \mathbb{N}}$  be the successive jumps of a standard Poisson Process (on the real line). Set  $\tau(u, x) = E_1[u\beta(x)]\alpha(x)$ , and  $v_m = \sup\{u \geq 0 : \tau(u, \sigma_m) \geq \tau_m\}$ , that is:

$$v_m = E_1^{-1} \left[ \frac{\tau_m}{\alpha(\sigma_m)} \right] \beta(\sigma_m)^{-1}$$

or  $v_m = 0$  if  $\alpha(\sigma_m) = 0$ .

Then, the random measure defined by  $\Gamma(dx) := \sum_{m \in \mathbb{N}} v_m \delta_{\sigma_m}(dx)$  is a generalised Gamma process  $\Gamma \sim \mathcal{G}aP(\alpha(dx), \beta(x))$  where  $\alpha(dx) = \Pi(dx)\alpha(x)$

In our setting, we will refer to the  $\{v_m\}$  as the weights of the Gamma process and to the  $\sigma_m$  as the locations.

**Corollary 1.2.** The Gamma process  $\Gamma \sim \mathcal{G}aP(\alpha(dx), \beta(x))$  can be approximated in distribution to arbitrarily high accuracy by the following:

---

**Algorithm 2** Inverse Levy measure algorithm

---

- 1: Initialisation: Fix a large integer  $M$  and choose  $\Pi(dx)$ , a convenient distribution on  $E$ , such that  $\alpha(x) = \frac{\alpha(dx)}{\Pi(dx)}$
  - 2: Generate  $M$  independent identically distributed draws  $\{\sigma_m\}_{m \leq M}$  from  $\Pi(dx)$
  - 3: Generate  $\{\tau_m\}_{m \leq M}$ , the first  $M$  jumps of a standard Poisson process on the real line (1)
  - 4: Set  $v_m = E_1^{-1} \left[ \frac{\tau_m}{\alpha(\sigma_m)} \right] \beta(\sigma_m)^{-1}$  (2)
  - 5: Set  $\Gamma_m(dx) \approx \Gamma_M(dx) = \sum_{m \leq M} v_m \delta_{\sigma_m}(dx)$
- 

**Remark 1.5.** A standard Poisson process on the real line can be simulated adding independent exponential random variables.

**Remark 1.6.** For numerical simulation it is useful to know that:  $E_1$  can be seen as

$$E_1(x) = \lim_{a \rightarrow 0} \left[ \int_x^\infty e^{-t} t^{a-1} dt \right]$$

and therefore it can be computed as the limit of rescaled  $\chi^2$ -tail probabilities as the degrees of freedom  $d = 2a$  tend to zero.

$$E_1(x) = \lim_{d \rightarrow 0} \Gamma(d/2) Pr[\chi_d^2 > 2x] = \lim_{d \rightarrow 0} (2/d) Pr[\chi_d^2 > 2x]$$

Therefore, if we know the inverse  $CDF_d(x)$  of a  $\chi_d^2$ ,  $PPF_d(x)$  (it is implemented in many scripting languages):

$$E_1^{-1}(x) = \lim_{d \rightarrow 0} \frac{1}{2} PPF_d(1 - (d/2) * y)$$

Context for Theorem 1.5 and and its proof is available in [4], while more on Remark 1.6 can be found in [5].

Let's see now a numerical example:

**Example 1.1.**  $E = [0, 2] \times [0, 2]$

$M = 1000$

$\Pi(dx) = \frac{1}{4}dx$  i.e. the uniform measure on the square

$\alpha(x) = 10\|x\|_1$

$\beta(x) = 2 + x_1$




Figure 1: Gamma Process  $W$  on  $E$ .



## 1.4 Cox process and the Poisson-Gamma random measure

Ref: <https://hpaulkeeler.com/cox-point-process/>

The Poisson Gamma random measure is a special case of Cox point process:

**Definition 1.8.** Let  $\Lambda(dx)$  be a non-negative, a.s. locally integrable, random measure on  $E$ . Then a point Process  $N$  defined on  $E$ , is a Cox point process driven by  $\Lambda$ , if the conditional distribution of  $N$  given  $\Lambda$  is a Poisson point process with intensity function  $\Lambda$ .

$$N(dx)|\Lambda \sim ppp(\Lambda(dx))$$

Note that  $\Lambda$  and  $N$  can be defined on different spaces, for our purposes they are the same.

The Cox process is also known as doubly-stochastic Poisson Process, as two degrees of uncertainty are introduced: first, there is a underlying random measure  $\Lambda$ , then only a Poisson Point Process  $N$  is observed (whose intensity measure is  $\Lambda$ ).

In particular, we need to define shot noise Cox processes, where  $\Lambda(dx)$  is defined as following:

1. First we consider a point process  $W$  on  $E$  to create a shot noise term (shot noise is just the sum of some function over all the points of a process).
2. Then we introduce a (non-negative) kernel function  $k_\theta(\cdot, \cdot)$  on  $E$ .
3. Now,  $\Lambda(x) = \int_E k_\theta(x, y)W(dy)$  and  $\Lambda(dx) = \Lambda(x)w(dx)$  for some reference measure  $w(dx)$  on  $E$  (we will use the Lebesgue measure or a rescaled version of it).

We are now ready for:

**Definition 1.9.** A Poisson-Gamma random measure is a shot noise Cox process  $N(dx)$ , where  $W$  is a Gamma process, and  $\Lambda(dx)$  is derived as above from  $\Lambda(x) = \int_E k_\theta(x, y)W(dy)$  for some integral kernel  $k_\theta(\cdot, \cdot)$ .

**Example 1.2.** Setting  $E = [0, 100] \times [0, 100]$ ,  $M = 1000$ ,  $\beta^{-1} = 1.5 * 10^5$ ,  $H(dx) = \alpha(dx) = 70 * \beta$ ;

First, we simulate the Gamma process  $W \sim \mathcal{GaP}(\alpha(dx), \beta(x))$ , with Algorithm 2:




Figure 2:  $\Lambda(dx)$

In this project, as in [4], we will model  $k_\theta(x, y)$  as a Gaussian kernel density:  $k_\theta(x, y) = \frac{1}{\pi\theta^2} \exp\left(-\frac{|x-y|}{\theta^2}\right)$ . Therefore,  $\Lambda$  can be seen as a "mixture" of Gaussian kernels whose weights are the weights of the underlying Gamma process.

Choosing  $\theta = 5$ ,  $w(dx) = 10^{-4}dx$ , we can now compute  $\Lambda(dx)$  (as explained after Definition 1.8) and plot it:




Figure 3:  $\Lambda(dx)$

In the end, using Algorithm 1, we simulate  $N \sim ppp(\Lambda)$ . In particular:

1. to compute  $\Lambda(E)$ , instead of a bidimensional quadrature, it's easier and faster to take advantage of the geometry of the domain and of the properties of Gaussian kernels. Therefore, if  $E = [a, b] \times [c, d]$  and  $\Phi(x)$  is the CDF of the standard normal distribution:

$$\Lambda(E) = \sum_{m \leq M} \underbrace{\left\{ v_m \left[ \Phi\left(\frac{b - \sigma_m^x}{\theta_3}\right) - \Phi\left(\frac{a - \sigma_m^x}{\theta_3}\right) \right] \left[ \Phi\left(\frac{d - \sigma_m^y}{\theta_3}\right) - \Phi\left(\frac{c - \sigma_m^y}{\theta_3}\right) \right] \right\}}_{w_m}$$

where  $\{v_m\}_{m \leq M}$  are the weights and  $\{\sigma_m\}_{m \leq M} = \{(\sigma_m^x, \sigma_m^y)\}_{m \leq M}$  are

the locations of latent the Gamma process.

2. To sample from  $X_i \sim \frac{\Lambda(\cdot)}{\Lambda(E)}$  one can use a common Acceptance-Rejection method; however, it is (again) faster and easier to take advantage of the structure of  $\Lambda$  and follow the scheme:

---

**Algorithm 3** Sampling from  $\frac{\Lambda(\cdot)}{\Lambda(E)}$

---

- 1: Sample  $i \leq M$  with probability  $p_m = \frac{w_m}{\Lambda(E)}$
  - 2: **while**  $x = (x_i, y_i) \notin E$  **do**
  - 3:   Sample  $x_i \sim \mathcal{N}(\sigma_i^x, \theta_3^2)$
  - 4:   Sample  $y_i \sim \mathcal{N}(\sigma_i^y, \theta_3^2)$
  - 5: **end while**
- 

Note that step 1 can be seen as the choice of which point of the Gamma process generates the  $i$ -th observation of the Cox process.

Eventually:




Figure 4: Points of the Poisson-Gamma Process  $N$  over the contour plot of  $\Lambda(x)$

## 2 Bayesian Inference

### 2.1 The Bayesian framework

Given a sample  $\underline{X} = (X_1, \dots, X_n)$  for a density  $f_\theta(x)$ , with an unknown parameter  $\theta \in \Theta$ , the associated likelihood function is:

$$\mathcal{L}(\underline{X}|\theta) = \prod_i^n f_\theta(x_i).$$

This quantity is a fundamental entity for the analysis of the information provided about  $\theta$  by the sample  $\underline{X}$ , and Bayesian analysis relies on this function to draw inference on  $\theta$ . The major input of the Bayesian approach, compared with a standard likelihood approach, is that it modifies the likelihood into a posterior distribution, which is a probability distribution on  $\Theta$  defined by:

$$\pi(\theta|\underline{X}) = \frac{\mathcal{L}(\underline{X}|\theta)\pi(\theta)}{\int_\Theta \mathcal{L}(\underline{X}|\theta)\pi(\theta)d\theta}.$$

The factor  $\pi(\theta)$  is called the "prior" and it has to be "chosen" to start the analysis. A first motivation for this approach is that the prior distribution summarizes the information that is available on  $\theta$  prior to the observation of the sample  $\underline{X}$ . However, the choice of  $\pi(\theta)$  is often decided on practical grounds rather than strong subjective beliefs or prior information.

The concept that is at the core of Bayesian analysis is that one should provide an inferential assessment conditional on the realized value of  $\underline{X}$ , and Bayesian inference gives a proper probabilistic meaning to this conditioning by allocating to  $\theta$  a probability distribution.

### 2.2 Markov chain Monte Carlo

The posterior distribution is often not directly computable in closed form. Therefore, to sample from such a probability distribution, Markov Chain Monte Carlo (MCMC) methods are employed.

By creating a Markov chain  $\{\theta^{(t)}\}_{t \geq 1}$  where the desired distribution  $\pi(\theta)$  is its equilibrium distribution, we can generate a sample of  $\pi$  by recording states from the chain. If an algorithm that generates such a chain can be constructed, the ergodic theorem guarantees that, in almost all settings, the empirical average (over time):  $\frac{1}{T} \sum_{t=1}^T g(\theta^{(t)})$  converges to  $\mathbb{E}(g(\theta))$ , independently from the starting value.

Several algorithms exist for constructing such chains, including Metropolis-Hastings and Gibbs sampling, among others. These algorithms iteratively generate states of the Markov chain, ensuring that over time, the states converge to the desired distribution.

### 2.2.1 Metropolis-Hastings algorithm

We want to sample from a target distribution  $\pi(\theta)$  (known up to a normalizing constant). Typically (in this project) it will be the posterior, that is  $\pi(\theta|\underline{X})$  but for simplicity we omit the dependence on  $\underline{X}$ . Note that the method is suitable to sample from any distribution.

To do so, we need to:

1. define a proposal kernel, i.e. a probability density of moving from  $\theta_{old}$  to  $\theta_{new}$ :

$$q(\theta_{new}|\theta_{old})$$

2. define an acceptance probability:

$$P(\theta_{new}, \theta_{old}) = \min \left( 1, \frac{\pi(\theta_{new})q(\theta_{old}|\theta_{new})}{\pi(\theta_{old})q(\theta_{new}|\theta_{old})} \right)$$

Then, the Metropolis-Hastings algorithm is as follows:

---

#### Algorithm 4 Generic Metropolis-Hastings Sampler

---

- 1: Initialisation: Choose an arbitrary starting value  $\theta^{(0)}$
  - 2: **for**  $t \geq 1$  **do**
  - 3:   Given  $\theta^{(t-1)}$ , sample  $\hat{\theta}$  from  $q(\theta|\theta^{(t-1)})$
  - 4:   Compute  $P = P(\hat{\theta}, \theta^{(t-1)})$  as above
  - 5:   With probability  $P$  accept  $\hat{\theta}$  and set  $\theta_t = \hat{\theta}$ ; otherwise, reject  $\hat{\theta}$  and set  $\theta^{(t)} = \theta^{(t-1)}$
  - 6: **end for**
- 

**Remark 2.1.** While theoretical guarantees that the algorithm converges are high, the choice of  $q$  remains essential in practice. Poor choices of  $q$  may indeed result either in a very high rejection rate, meaning that the Markov chain hardly moves, or in a myopic exploration of the support of  $\pi$ , that is, in a dependence on the starting value  $\theta_0$  such that the chain is stuck in a neighborhood region of  $\theta_0$ . A particular choice of proposal  $q$  may thus work well for one target density but be extremely poor for another one. While the algorithm is indeed universal, it is impossible to prescribe application-independent strategies for choosing  $q$ .

Choosing particular  $q(\theta_{new}|\theta_{old})$  we can have:

- The **Independence Sampler**: when  $q$  is independent from  $\theta_{old}$ , i.e.

$$q(\theta_{new}|\theta_{old}) = q(\theta_{new}),$$

and then  $P$  simplifies to

$$P(\theta_{new}, \theta_{old}) = \min \left( 1, \frac{\pi(\theta_{new})q(\theta_{old})}{\pi(\theta_{old})q(\theta_{new})} \right).$$

- The **Random Walk Sampler**: when  $q$  is a symmetric transition kernel i.e.:

$$q(\theta_{new}|\theta_{old}) = f(\theta_{new} - \theta_{old}),$$

where  $f$  is a symmetric density function and then  $P$  simplifies to

$$P(\theta_{new}, \theta_{old}) = \min \left( 1, \frac{\pi(\theta_{new})}{\pi(\theta_{old})} \right).$$

For example  $f$  can be a normal density with mean zero and given variance (covariance matrix).

### 2.2.2 Gibbs Sampler

Let  $\theta = (\theta_1, \dots, \theta_p) \in \mathbb{R}^p$  and set  $\theta_{-i} = (\theta_1, \dots, \theta_{i-1}, \theta_{i+1}, \dots, \theta_p)$ .

This algorithm specifically works for sampling from  $\pi(\theta|\underline{X})$ . To do so, however we should be able to compute (by hand) all the full conditional distributions:

$$\pi_i(\theta_i|\underline{X}, \theta_{-i}) \quad \forall i = 1, \dots, p$$

---

#### Algorithm 5 Gibbs Sampler

---

- 1: Initialisation: Choose an arbitrary starting value  $\theta^{(0)} = (\theta_1^{(0)}, \dots, \theta_p^{(0)})$
  - 2: **for**  $t \geq 1$  **do**
  - 3:   Given  $\theta^{(t-1)}$ , we sample  $\theta^t$  with the following scheme:
  - 4:     1)  $\theta_1^{(t)} \sim \pi_1(\theta_1|\underline{X}, \theta_{-1})$
  - 5:     2)  $\theta_i^{(t)} \sim \pi_i(\theta_i|\underline{X}, \theta_{-i})$
  - 6:     ...
  - 7:     p)  $\theta_p^{(t)} \sim \pi_p(\theta_p|\underline{X}, \theta_{-p})$
  - 8: **end for**
- 

#### Remark 2.2. Metropolis-within-Gibbs algorithm

Several MCMC algorithms can be mixed together within a single algorithm using either a circular or a random design. While this construction is often suboptimal (in that the inefficient algorithms in the mixture are still used on a regular basis), it almost always brings an improvement compared with its individual components. A special case where a mixed scenario is used is the Metropolis-within-Gibbs algorithm: when building a Gibbs sampler, it may happen that it is difficult or impossible to simulate from one or several of the conditional distributions. In that case, a single Metropolis step associated with this conditional distribution (as its target) can be used instead.

### 3 Bayesian inference for Poisson-Gamma Random measures

A Poisson-Gamma random measure  $N(dx)$  can be modelled as Cox process with the following hierarchy:

$$\begin{aligned} \alpha, \beta, \theta &\sim \pi(\alpha, \beta, \theta) & W(dx)|\alpha, \beta &\sim \mathcal{G}aP(\alpha(dx), \beta) \\ \Lambda(dx) &= \int_E k_\theta(dx, y)W(dy) & N(dx)|\alpha, \beta, \theta, W &\sim ppp(\Lambda(dx)) \end{aligned}$$

Now, given one or more observation of  $N$  we want to sample from the posterior distribution of  $(\alpha, \beta)$ . On the other side, we suppose that the spatial interaction  $\theta$  is known (it can be derived from the physical nature of the phenomenon we would like to study).

A model for Bayesian inference on Poisson Gamma-random measures has been already proposed in [4], where  $H = \alpha(dx)$  is supposed to be a continuous measure on  $E$ . However, usually, no prior information is available on the latent process  $W(dx)$  and  $\alpha(dx)$  is chosen as the Lebesgue measure on  $E$  rescaled by a factor  $\alpha = \text{constant}$ . In the following we propose a different method, where  $H(dx)$  is a discrete measure and we will show a result that (partially) connects the two methods.

Moreover, we believe our model, although "simpler", is computationally less expensive, more stable and will allow us to sample from the posterior observing more realizations of the same process at once (while in [4] only one realization is taken in consideration at a time). In the following we also refer to this version of the latent Gamma Process as a discrete Gamma Process (although it is still a Gamma process and a CRM).

#### 3.1 On the discrete Gamma Process

**Definition 3.1.** A discrete Gamma process is a Gamma process  $\mathcal{G}aP(H_D, \beta)$  on a space  $E$  when  $H_D(dx)$  is a discrete measure on  $E$ .

**Proposition 3.1.** A discrete Gamma process  $W(dx) \sim \mathcal{G}aP(H_D, \beta)$  on a space  $E$  where  $H_D(dx) = \sum_{m=1}^M \alpha_m \delta_{\sigma_m}(dx)$  and  $\{\sigma_m\}_{m \leq M}$  are fixed points of  $E$ , can be written as:

$$W(dx) = \sum_{m=1}^M v_m \delta_{\sigma_m}(dx)$$

where  $v_m \sim \mathcal{G}a(\alpha_m, \beta) \forall m = 1, \dots, M$  and  $(v_1, \dots, v_M)$  are independent.

*Proof.* Since a Gamma Process is a CRM, for all  $A_1, \dots, A_k \subset E$  measurable and disjoint,  $W(A_1), \dots, W(A_k)$  are independent.

Moreover,  $\forall A \in E$  and  $\forall t > 0$ ,  $tW(A) = \int_E t\mathbb{I}_A(x)W(dx)$ .



Therefore:

$$\begin{aligned}
L_{W(A)}(t) &= \mathbb{E}[e^{-tW(A)}] \\
&= \mathbb{E}[e^{-\int_E f(x)W(dx)}] \quad f(x) = t\mathbb{I}_A(x) \\
&= \exp \left\{ - \int \log \left( 1 + \frac{t\mathbb{I}_A(x)}{\beta} \right) H_D(dx) \right\} \\
&= \exp \left\{ - \log \left( 1 + \frac{t}{\beta} \right) H_D(A) \right\} \\
&= \left( 1 + \frac{t}{\beta} \right)^{-H_D(A)}
\end{aligned}$$

Which is equivalent to:  $W(A) \sim \mathcal{Ga}(H_D(A), \beta)$  if  $H_D(A) > 0$  or  $W(A) = 0$  a.s. if  $H_D(A) = 0$ .

Now, since  $H_D(dx) = \sum_{m=1}^M \alpha_m \delta_{\sigma_m}(dx)$ :

$$\begin{aligned}
\Rightarrow H_D(A) &= \sum_{m: \sigma_m \in A} \alpha_m \\
\Rightarrow W(A) &= 0 \text{ a.s. if } \sigma_m \notin A \quad \forall m = 1, \dots, M \\
\Rightarrow W_\omega(A) &= \sum_{m=1}^M v_m(\omega) \delta_{\sigma_m}(A) \text{ a.s.}
\end{aligned}$$

which means that  $v_m = W(A_m)$  if  $\sigma_m \in A_m$  and  $\sigma_k \notin A_m \quad \forall k \neq m$ .

Therefore,  $v_m \sim \mathcal{Ga}(\alpha_m, \beta) \quad \forall m = 1, \dots, M$  and  $(v_1, \dots, v_M)$  are independent.  $\square$

When modeling a Poisson Gamma-random measure with the previous hierarchy, no prior information is usually evident or available on the latent Gamma process, and a good choice is:  $\alpha(dx) = \alpha \cdot m(dx)$  (where  $m(dx)$  is the Lebesgue measure on  $E$ ). This is equivalent to believing that, a priori, no subset of  $E$  will have a bigger probability to "generate" points of the process  $N(dx)$ .

In this situation, we believe it can also be chosen to model  $\alpha(dx)$  as:  $\alpha_D(dx) = \sum_{m=1}^M \frac{\alpha}{M} \delta_{\sigma_m}(dx)$  where  $\{\sigma_m\}_{m \leq M}$  are equally spaced on  $E$ .

In fact, if:

$$\begin{aligned}
W_D &\sim \mathcal{GaP}(\alpha_D(dx), \beta) \\
W &\sim \mathcal{GaP}(\alpha \cdot m(dx), \beta)
\end{aligned}$$

It can be proved that  $\alpha_D(x) \xrightarrow{\mathcal{L}} \alpha \cdot m(dx)$ ,

and therefore follows that:

$$\mathbb{E} \left[ e^{-tW_D(A)} \right] \longrightarrow \mathbb{E} \left[ e^{-tW(A)} \right]$$

for every  $f : E \rightarrow \mathbb{R}_+$  measurable.

An in-depth theoretical analysis is not the intent of this project, however, we believe that what we have pointed out shows that we can expect a similar behaviour of  $W_D$  and  $W$  when  $M$  is chosen big.

Further results on the convergence of the process can follow from the results in chapter 14 of [8].

### 3.2 Sampling from the discrete Gamma Process

We propose a different way to sample from the discrete Gamma Process and from a Poisson-Gamma random measure with it as the latent process. The properties shown here are key to compute the likelihood of the model.

Let's see in detail how it is done.

We start by recalling some properties of the Gamma random variable:

If

$$v_1, \dots, v_M \stackrel{\text{IND}}{\sim} \mathcal{Ga}(\alpha_m, \beta) \quad \forall m = 1, \dots, M$$

and

$$\bar{v} := \sum_{m=1}^M v_m, \quad D_m := \frac{v_m}{\bar{v}} \quad \forall m = 1, \dots, M$$

then:

- $\bar{v}$  and  $(D_1, \dots, D_M)$  are independent,
- $\bar{v} \sim \mathcal{Ga} \left( \sum_{m=1}^M \alpha_m, \beta \right)$
- $(D_1, \dots, D_M) \sim \text{DIR}(\alpha_1, \dots, \alpha_M)$ ,

where  $\text{DIR}$  is the notation for the Dirichlet distribution, or multivariate Beta distribution, which is a generalisation on  $A \subset \mathbb{R}^n$  of the Beta distribution.

Therefore, for our purpose the latent (discrete Gamma) process reduces to:

$$\begin{aligned} W(dx) &= \sum_{m=1}^M v_m \delta_{\sigma_m}(dx) \\ &= \bar{v} \sum_{m=1}^M D_m \delta_{\sigma_m}(dx) \end{aligned}$$

where  $v_m, \bar{v}, D_m$  are distributed as above.

We are confident that it is possible to prove

Therefore, drawing samples from the Poisson-Gamma random measure  $N(dx)$  reduces to:

0. Choose arbitrarily:

- $M$ , the number of fixed locations  $\{\sigma_m\}_{m \leq M}$  and the coordinates of each  $\sigma_m$ ,
- $\alpha_i \quad \forall m = 1, \dots, M$ , the shape parameter of each  $v_m$  and set  $\alpha = \sum_{m=1}^M \alpha_m$
- $\beta$ , the (same) rate parameter of each  $v_m$
- $\theta$ , the standard deviation of the Gaussian Kernel  $k_\theta(x, y) = \frac{1}{\pi\theta^2} \exp\left(-\frac{|x-y|}{\theta^2}\right)$

1. Sample

$$\bar{v} \sim \mathcal{Ga}(\alpha, \beta)$$

2. Sample

$$(D_1, \dots, D_M) \sim \text{DIR}(\alpha_1, \dots, \alpha_M)$$

3. Sample

$$N \sim \text{Pois}(\Lambda(E))$$

where  $\Lambda(E) = \bar{v} \sum_{m=1}^M D_m k_\theta(E|\sigma_m)$ .

4. Choose the locations  $S_1, \dots, S_n$  that generates each of the  $n$  (the realization of  $N$ ) with probability:

$$P(S_i = \sigma_m | N = n, D_1, \dots, D_M) = \frac{D_m k_\theta(E|\sigma_m)}{\sum_{m=1}^M D_m k_\theta(E|\sigma_m)} \quad \forall i = 1, \dots, n \quad \forall m = 1, \dots, M$$

5. Sample the points  $\{X_i\}_{i=1}^n = \{(x_i, y_i)\}_{i=1}^n$  independently with probability:

$$P(X_i \in dx | N = n, S_i) = \frac{k_\theta(x|S_i)dx}{k_\theta(E|S_i)} \quad \forall i = 1, \dots, n$$

**Example 3.1.** Choosing:  $E = [0, 10] \times [0.10]$ ,  $M = 64$ ,  $\{\sigma_m\}_{m \leq M}$  equally spaced on  $E$ ,  $\theta = 0.5$ ,  $\beta = 2.5 \cdot 10^{-2}$ ,  $\alpha = 20$  and  $\alpha_m = \alpha/M \quad \forall m = 1, \dots, M$ , we can sample as above and plot the process:

### 3.3 The Metropolis within Gibbs algorithm

Now, we want to compute the likelihood of the model:

$$\begin{aligned} \mathcal{L}(\underline{X} | \alpha_1, \dots, \alpha_M, \beta) &= f_{\bar{v}}(\bar{v}) f_D(D_1, \dots, D_M) f_N(N | \Lambda(E)) \\ &\quad \times f_{X,S}(X_1, \dots, X_N, S_1, \dots, S_N | N, D) \end{aligned} \quad (3)$$




Figure 5: A realization of the simplified process

where:

$$\alpha = \sum_{m=1}^M \alpha_m$$

$$f_{\bar{v}}(\bar{v}) = \frac{\bar{v}^{\alpha-1} e^{-\beta \bar{v}} \beta^{\alpha}}{\Gamma(\alpha)}$$

$$f_D(D_1, \dots, D_M) = \Gamma(\alpha) \cdot \prod_{m=1}^M \frac{D_m^{\alpha_m-1}}{\Gamma(\alpha_m)}$$

$$\Lambda(E) = \bar{v} \sum_{m=1}^M D_m k_{\theta}(E|\sigma_m)$$

$$f_N(N|\Lambda(E)) = \frac{\Lambda(E)^N e^{-\Lambda(E)}}{N!} \mathbb{I}_{\mathbb{N}}(N)$$

$$f_{X,S}(X_1, \dots, X_N, S_1, \dots, S_N|N, D) = \prod_{i=1}^N \frac{D_i k_{\theta}(E|S_i)}{\sum_{m=1}^M D_m k_{\theta}(E|\sigma_m)} \frac{k_{\theta}(X_i|S_i)}{k_{\theta}(E|S_i)}$$

Therefore, supposing  $(\alpha_1, \dots, \alpha_M)$  and  $\beta$  independent, simplifying and rear-

ranging the terms, the joint likelihood becomes:

$$\begin{aligned} \mathcal{L}(\underline{X}|\alpha_1, \dots, \alpha_M, \beta) \pi(\alpha_1, \dots, \alpha_M, \beta) &= \pi(\alpha_1, \dots, \alpha_M) \pi(\beta) \bar{v}^{\alpha-1} \bar{v}^N e^{-\beta \bar{v}} \\ &\times \exp \left\{ -\bar{v} \sum_{m=1}^M D_m k_\theta(E|\sigma_m) \right\} \times \beta^\alpha \frac{1}{N!} \prod_{m=1}^M \frac{D_m^{\alpha_m-1}}{\Gamma(\alpha_m)} \prod_{i=1}^N D_i k_\theta(X_i|S_i) \end{aligned} \quad (4)$$

We can now compute the full conditionals:

1.  $S_i|All$ :

$$\mathbb{P}(S_i = \sigma_m) = \frac{D_i k_\theta(X_i|S_i)}{\sum_{m=1}^M D_m k_\theta(X_i|\sigma_m)} \quad \forall i = 1, \dots, N \quad \forall m = 1, \dots, M$$

2.  $(D_1, \dots, D_M)|All$ :

$$\begin{aligned} &\propto \prod_{m=1}^M D_m^{\alpha_m-1} D_m^{n_m} \cdot e^{-\Lambda_D(E)} \\ &\propto DIR(\alpha_1 + n_1, \dots, \alpha_M + n_M) \cdot e^{-\Lambda_D(E)} \end{aligned}$$

where  $n_m = \#\{j = 1, \dots, N | S_j = \sigma_m\}$ .

This full conditional is not known in closed form; nonetheless, we will sample with a step of an Independence Sampler. Choosing as a proposal  $q(D_{new}|D_{old}) = q(D_{new}) = DIR(\alpha_1 + n_1, \dots, \alpha_M + n_M)$ , then the acceptance probability becomes:

$$P(D_{new}, D_{old}) = \min \left( 1, \frac{e^{-\Lambda_{D_{new}}(E)}}{e^{-\Lambda_{D_{old}}(E)}} \right)$$

3.  $\bar{v}|All$ :

$$\propto \bar{v}^{\alpha-1} \bar{v}^N e^{-\beta \bar{v}} \times \exp \left\{ -\bar{v} \sum_{m=1}^M D_m k_\theta(E|\sigma_m) \right\}$$

and therefore we recognise that:

$$\bar{v}|All \sim \mathcal{Ga} \left( \alpha + N, \beta + \sum_{m=1}^M D_m k_\theta(E|\sigma_m) \right)$$

4.  $\beta|All$ :

$$\propto \pi(\beta) e^{-\beta \bar{v}} \beta^\alpha$$

Choosing  $\pi(\beta) \sim \mathcal{Ga}(a_0, b_0)$  the model is conjugate and

$$\beta|All \sim \mathcal{Ga}(a_0 + \alpha, b_0 + \bar{v})$$

5. For simplicity we will suppose  $\alpha_m = \frac{\alpha}{M} \quad \forall m = 1, \dots, M$  and we will sample from the posterior of  $\alpha$ .

$\alpha|All$ :

$$\propto \pi(\alpha) \bar{v}^{\alpha-1} \beta^\alpha \prod_{m=1}^M \frac{D_m^{\alpha_m-1}}{\Gamma(\alpha_m)}$$

Since we don't recognise any known distribution we use a Random-walk Metropolis-Hastings algorithm.

Having computed all the full conditionals, the Metropolis within Gibbs Sampler is ready to be implemented.

### 3.4 The algorithm with more observations (stationary case)

We can modify the the results above to sample from the posterior of  $(\alpha, \beta)$  after we have seen multiple realizations of the same phenomenon. In the following it is supposed that the latent Gamma Process doesn't change over time (so do not  $\alpha$  and  $\beta$ ) and each observation is independent from the others.

If  $T$  is the number of realizations,  $N_t$  the number of each observations at time  $t = 1, \dots, T$ , then the joint posterior becomes:

$$\begin{aligned} \mathcal{L}(\underline{X}_1, \dots, \underline{X}_T | \alpha_1, \dots, \alpha_M, \beta) \pi(\alpha_1, \dots, \alpha_M, \beta) &= \pi(\alpha_1, \dots, \alpha_M) \pi(\beta) \left( \prod_{t=1}^T \bar{v}_t \right)^{\alpha-1} \\ &\times \prod_{t=1}^T \bar{v}_t^{N_t} \times \exp \left( -\beta \sum_{t=1}^T \bar{v}_t \right) \times \prod_{t=1}^T \exp \left( -\bar{v}_t \sum_{m=1}^M D_{m,t} k_\theta(E|\sigma_m) \right) \\ &\times \beta^{\alpha T} \times \prod_{t=1}^T \frac{1}{N_t!} \times \prod_{t=1}^T \prod_{m=1}^M \frac{D_{m,t}^{\alpha_m-1}}{\Gamma(\alpha_m)} \times \prod_{t=1}^T \prod_{i=1}^{N_t} D_{i,t} k_\theta(X_{i,t} | S_{i,t}) \end{aligned}$$

Therefore, the full conditionals of  $S_{i,t}$ ,  $D_{i,t}$ ,  $\bar{v}_t \quad \forall i = 1, \dots, N_t$  are the same as above  $\forall t = 1, \dots, T$ ; while, now:

1.  $\beta|All$ :

$$\propto \pi(\beta) \exp \left( -\beta \sum_{t=1}^T \bar{v}_t \right) \beta^{\alpha T}$$

Choosing again  $\pi(\beta) \sim \mathcal{Ga}(a_0, b_0)$  the model is still conjugate and

$$\beta|All \sim \mathcal{Ga} \left( a_0 + \alpha T, b_0 + \sum_{t=1}^T \bar{v}_t \right)$$

2. Same as before, for simplicity we will suppose  $\alpha_m = \frac{\alpha}{M}$  and we will sample from the posterior of  $\alpha$ .

$\alpha|All$ :

$$\propto \pi(\alpha) \left( \prod_{t=1}^T \bar{v}_t \right)^{\alpha-1} \beta^{\alpha T} \times \prod_{t=1}^T \prod_{m=1}^M \frac{D_{m,t}^{\alpha_m-1}}{\Gamma(\alpha_m)}$$

We don't recognise any known distribution, then we use a Random-walk Metropolis-Hastings algorithm. In order to evaluate the acceptance probability it is advisable to calculate the logarithm of the targets (see below).

The final algorithm to sample from the posterior of  $(\alpha, \beta)$  then becomes:

---

**Algorithm 6** Metropolis within Gibbs Sampler for discrete Poisson-Gamma random measures

---

1: Initialisation:

- choose arbitrary starting values for:  $\bar{v}_t^{(0)}, D_{m,t}^{(0)} \quad \forall t = 1, \dots, T \quad \forall m = 1, \dots, M$ ;
- choose arbitrary starting values for  $\alpha^{(0)}, \beta^{(0)}$ ;
- choose wisely  $\pi(\alpha)$ , a prior for  $\alpha$ ,  $\sigma_{MC}^2$ , the variance for the proposal of  $\alpha_{new}$ , and  $a_0, b_0$  small  $\in \mathbb{R}$ , shape and rate for the prior  $\pi(\beta) \sim \mathcal{Ga}(a_0, b_0)$ .

2: **for**  $it \geq 1$  **do**

3:   **for**  $t = 1, \dots, T$  **do**

4:     Assign  $S_{i,t}^{(it)}$  with:

$$\mathbb{P}(S_{i,t}^{(it)} = \sigma_m) = \frac{D_{i,t}^{(it-1)} k_\theta(X_i | S_i)}{\sum_{m=1}^M D_{m,t}^{(it-1)} k_\theta(X_i | \sigma_m)} \quad \forall i = 1, \dots, N_t \quad \forall m = 1, \dots, M$$

5:     Sample:

$$D_{new,t} = (D_{new,t,1}, \dots, D_{new,t,M}) \sim \text{DIR}(\alpha_1^{(it-1)} + n_{1,t}, \alpha_M^{(it-1)} + n_{M,t})$$

and set  $D_t^{(it)} = D_{new,t}$  with probability  $P_1$ , otherwise  $D_t^{(it)} = D_t^{(it-1)}$ , where:

$$P_1(D_{new,t}, D_t^{(it-1)}) = \min \left( 1, \frac{e^{-\Lambda_{D_{new,t}}(E)}}{e^{-\Lambda_{D_t^{(it-1)}}(E)}} \right)$$

6:     Sample:

$$\bar{v}_t^{(it)} \sim \mathcal{Ga} \left( \alpha^{(it-1)} + N_t, \beta^{(it-1)} + \sum_{m=1}^M D_{m,t}^{(it)} k_\theta(E | \sigma_m) \right)$$

7:   **end for**

8:     Sample:

$$\beta^{(it)} \sim \mathcal{Ga} \left( a_0 + \alpha^{(it-1)} T, b_0 + \sum_{t=1}^T \bar{v}_t^{(it)} \right)$$

9:     Sample:

$$\alpha_{new} \sim \mathcal{N}(\alpha^{(it-1)}, \sigma_{MC}^2)$$

and set  $\alpha^{(it)} = \alpha_{new}$  with probability  $P_2$ , otherwise  $\alpha^{(it)} = \alpha^{(it-1)}$ , where:

$$\log P_2(\alpha_{new}, \alpha^{(it-1)}) = \min \left( 0, \log t(\alpha_{new}) - \log t(\alpha^{(it-1)}) \right)$$

and:

$$\log t(\alpha) = \log(\pi(\alpha)) + (\alpha - 1) \sum_{t=1}^T \log \left( \bar{v}_t^{(it)} \right) + \alpha T \log \left( \beta^{(it)} \right)$$

24

$$+ \frac{\alpha - M}{M} \sum_{t=1}^T \sum_{m=1}^M \log \left( D_{m,t}^{(it)} \right) - MT \log \left( \Gamma \left( \frac{\alpha}{M} \right) \right)$$

10: **end for**

---



### 3.5 The algorithm with more observations (time-varying case)

If we believe the latent process changes over time we can capture this aspect allowing  $\alpha$  to evolve over time. In particular, if  $\beta$  is stationary then  $\alpha_t$  is directly linked with the number of points observed at each time  $t$ ; in fact, we can approximate:  $\hat{N}_t = \mathbb{E}(\Lambda_t(E)) \simeq \frac{\alpha_t}{\beta}$ , which is an estimator of the number of observed points at time  $t$ .

With  $\alpha$  time-varying the full conditional becomes:

$\alpha_1, \dots, \alpha_T | All:$

$$\propto \pi(\alpha_1, \dots, \alpha_T) \left( \prod_{t=1}^T \bar{v}_t^{\alpha_t-1} \right) \beta^{\sum_{t=1}^T \alpha_t} \times \prod_{t=1}^T \prod_{m=1}^M \frac{D_{m,t}^{\alpha_{t,m}-1}}{\Gamma(\alpha_{t,m})}$$

Having in mind the application in 4.2, one could model  $\alpha_t$  as  $\alpha_t = e^{\eta_0 + \eta_1 I(t) + \eta_2 A(t)}$  where:

- $I(t)$  is a dummy variable with values in  $\{0, 1\}$  that allows to capture seasonality, such as dry vs wet season (if we model forest fires) or working days vs public holidays (if we model taxi pickups).
- $A(t)$  is a variable that allows to capture trend, such as increasing/decreasing number of forest fires, taxi pick-ups over weeks or years.

Now we want to sample from the posteriors of  $(\eta_0, \eta_1, \eta_2)$  and we do so by replacing Step 9 of Algorithm 6 with three steps (one for each  $\eta_i$ ) of a RW-Metropolis sampler or one for the all three at the same time.

Step 9 of Algorithm 6 then becomes:

1. Sample:  $(\eta_{0,new}, \eta_{1,new}, \eta_{2,new}) \sim \mathcal{N}\left(\left(\eta_0^{(it-1)}, \eta_1^{(it-1)}, \eta_2^{(it-1)}\right), \sigma_{MC}^2 \mathbb{I}_3\right)$
2. Compute:  $\alpha_{new}$  and  $\alpha^{(it-1)}$
3. Set  $\eta_i^{(it)} = \eta_{i,new} \quad \forall i = 0, 1, 2$  with probability  $P_2$ , otherwise  $\eta_i^{(it)} = \eta_i^{(it-1)}$ , where, once computed  $\alpha_{new}$  and  $\alpha^{(it-1)}$ :

$$\log P_2(\alpha_{new}, \alpha^{(it-1)}) = \min\left(0, \log t(\alpha_{new}) - \log t(\alpha^{(it)})\right)$$

and:

$$\begin{aligned} \log t(\alpha) &= \log(\pi(\eta_0)\pi(\eta_1)\pi(\eta_2)) + \sum_{t=1}^T (\alpha_t - 1) \log\left(\bar{v}_t^{(it)}\right) + \left(\sum_{t=1}^T \alpha_t\right) \log\left(\beta^{(it)}\right) \\ &+ \sum_{t=1}^T \sum_{m=1}^M \left(\frac{\alpha_t}{M} - 1\right) \log\left(D_{m,t}^{(it)}\right) - M \sum_{t=1}^T \log\left(\Gamma\left(\frac{\alpha_t}{M}\right)\right) \end{aligned}$$

## 4 An application: inference on forest fires in the Amazon Forest

### 4.1 The stationary case

In this last section we propose an application of the algorithm above. Given the coordinates of forest fires in a portion of the Amazon Forest (Latitude in  $30^\circ\text{S}$ - $20^\circ\text{S}$  and Longitude in  $65^\circ\text{W}$ - $55^\circ\text{W}$ ) over three different months of the dry season (August 2016, September 2017, September 2018) we would like to model the latent process generating the fires and sample from the posterior of  $(\alpha, \beta)$ .




Figure 6: Geoplot of the Coordinates of the forest fires over three months.

For simplicity we decided not to take in consideration the curvature of the Earth, then we translated the observations in the square  $E = [0, 10] \times [0, 10]$ .

Assigning  $\theta = 0.8$  a fair value for the spatial interaction, and choosing arbitrary values for the starting parameters (see the code for details), then running the Sampler for 5000 iterations, we can estimate the posterior distribution of  $\alpha$  and  $\beta$ .

For every month  $t = 1, \dots, T$  (here  $T = 3$ ), the algorithm samples from the posterior of  $D_t, \bar{v}_t$ . We can then plot the results for every single month (below an example with the third month).




Figure 7: Histogram (Burn-in: 2000) and Traceplot of  $\alpha^{(it)}$ .




Figure 8: Histogram (Burn-in: 2000) and Traceplot of  $\beta^{(it)}$ .

Figure 9: On the left: plot of the coordinates of the fires with estimates for the weights  $v_m$  of the latent process  $\Gamma(dx)$ ; On the right: surface plot of  $\Lambda(dx)$ .





Figure 10: Plot of the coordinates of the fires over the contour plot of  $\Lambda(E)$ .

## 4.2 The time-varying case

Now, we want to analyse the same dataset on a different time-frame and take in consideration 33 months, from January 2018 to September 2020 (the coordinates for each month are available here: **Project's Github**). Plotting the number of fires by month we can hypothesise that latent process changes over time and therefore model  $\alpha$  as:

$$\alpha_t = e^{\eta_0 + \eta_1 I(t) + \eta_2 A(t)}$$

where:

- $I(t) = 1$  if the  $t^{th}$  month belongs to the dry season (which spans from August to November) and  $I(t) = 0$  otherwise.
- $A(t) = 0, 1, 2$  if the  $t^{th}$  month belongs to the first, second, third year (2018, 2019, 2020); however, we consider December part of the previous year as it is the first month of the wet season.

Note that this way of modelling  $\alpha_t$  reflects the idea that in the same season of the same year the latent process has the same distribution.




Figure 11: Number of forest fires observed in the  $t^{th}$  month.

Applying the same transformations as in the stationary case and assigning  $\theta = 0,8$  we can now run the Metropolis within Gibbs sampler proposed in Algorithm 6 with the modification proposed in subsection 3.4 and sample from the posteriors of  $\eta_0, \eta_1, \eta_2, \beta$ . We run it for 10000 iterations to ensure reaching stationarity.




Figure 12: Traceplots of  $\eta_0, \eta_1, \eta_2, \beta$  and relative means (with a burn-in of 8000 iterations)

We can approximate an estimator for the number of points observed in each month with the empirical mean of  $\Lambda(E)$ , i.e  $\hat{N}_t = \mathbb{E}(\Lambda_t(E)) \simeq \frac{1}{L} \sum_{it=1}^L \Lambda_t^{(it)}(E)$

A similar way to do so is:  $\hat{N}_t = \mathbb{E}(\Lambda_t(E)) \simeq \frac{\alpha_t}{\beta} \simeq \frac{1}{L} \sum_{it=1}^L \frac{\alpha_t^{(it)}}{\beta^{(it)}}$  which is an estimator of the number of points within a fixed year and season (actually,  $N_t | \Lambda_t(E) \sim \text{Pois}(\Lambda_t(E))$  as seen before).




Figure 13

## 5 Further work

We believe the following areas present promising directions for future research:

1. **Convergence Analysis:** Prove a stronger result for the convergence of the Poisson-Gamma Random measure when the underlying discrete Gamma process converges to a non discrete one.
2. **Algorithmic Refinement:** Focus on optimizing the existing code and implementing an adaptive Metropolis-Hastings algorithm to enhance efficiency and accuracy.
3. **Exploration of Diverse Datasets:** Extend the study to analyze various datasets with the aim of improving the predictive performance of the algorithm. Examples from spatial statistics, such as taxi pickups in urban areas or instances of violent crime in neighborhoods, offer promising avenues for investigation. Utilize available online datasets to conduct comprehensive analyses and validate the algorithm's effectiveness across different scenarios.

## References

- [1] G. Last, M. Penrose. *Lectures on the Poisson process*. Cambridge University Press, 2018.
- [2] D. J. Daley, D. Vere-Jones. *An introduction to the theory of point processes, volume I. Elementary Theory and Methods*. Springer, 2003.
- [3] D. J. Daley D. Vere-Jones. *An introduction to the theory of point processes, volume II. General Theory and Structure*. Springer, 2008.
- [4] R. L. Wolpert, K. Ickstadt. *Poisson/Gamma random field models for spatial statistics*. Biometrika 85(2):251–267, 1998.
- [5] M. Abramowitz, I. A. Stegun. *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*. Applied Mathematics Series, 55. Washington, DC: National Bureau of Standards, 1964.
- [6] J. M. Marin, C. P. Robert. *A Bayesian Core: A Practical Approach to Computational Bayesian Statistics*. Springer, 2007.
- [7] F. Bassetti, R. Casarin, M. Iacopini. *A Spatiotemporal Gamma Shot Noise Cox Process*. ArXiv eprint arXiv:2308.08481, 2023.
- [8] O. Kallenberg. *Foundations of Modern Probability*. Springer, 2021.