

## Open Log and Read in Data

- Set your working directory and create a log file to divert your codes and results
- Read the external Stata data file gsscum7212Teach.dta into R.

```
>setwd("/Users/burrisfaculty/Desktop/DSCode/SOC686")
>sink("Shepherd_asgn07.log", split=T)
>rm(list=ls(all=TRUE))
>mygss <- read.dta("gsscum7212teach.dta", convert.factor=F)
```

## Manage Covariates Data

- Keep six variables, include mental health (mntlhlth), age (age), sex (sex), race (race), education (educ), and income (inclk)
- Create dummy variables for sex and race. Note that the race variable has three categories, so please create three dummy variables for race (Alternatively, you can the factor function to turn the race variable into a factor variable and use it directly in the regression). Also you need to be careful and clear about 1) how many of these three dummy variables, all measuring race, are usually used in a regression model and 2) how to interpret the results/corresponding coefficients (e.g., which group is the reference group?). Please also drop missing cases using listwise deletion (any case that has missing information for any of the six variables will be dropped from the sample data).
- Check the descriptive statistics of these variables using the table and the summary function when appropriate. Note that when there is too much output (e.g., tabulation of income), you can present representative information.

```
> useddta <- subset(mygss,
+                   select=c(mntlhlth, age, sex, race, educ, inclk))

> #Create dummy variables female (male = 0)
> useddta$female <- as.numeric(useddta$sex==2)

> useddta$male <- as.numeric(useddta$sex == 1)

> #Create Binary Indicator Variables for Multi-Category Nomial Variables
>
> useddta$white <- ifelse(useddta$race == 1, 1, 0)

> useddta$black <- ifelse(useddta$race == 2, 1, 0)

> useddta$other <- ifelse(useddta$race == 3, 1, 0)

> nmdta <- useddta[complete.cases(useddta),] #no missing data

> #summarize data
> summary(nmdta$mntlhlth)
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
0.000  0.000   0.000   3.991  5.000   30.000

> summary(nmdta$inclk)
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
0.245  15.921  28.157   39.404  48.475  155.140

> summary(nmdta$age)
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
18.00	31.00	42.00	41.81	51.00	84.00

```
> table(nmdta$female)
```

0	1
381	369

```
> table(nmdta$white)
```

0	1
160	590

## Task 1 Create Ordinal Response Variables

- Generate an ordered response variable based on the mental health variable such that the new ordered response variable is coded as one if the number of days for poor mental health is zero, two if the number of days for having poor mental health is one-seven, three if the number of days for having poor mental health is eight-fourteen, and four if the number of days for having poor mental health is fifteen and above. Label this new variable as `mntlhlthOrd`. Please crosstab `mntlhlthOrd` with `mntlhlth`.

```
> #Task 1 Create Ordinal Response Variable
```

```
> # 1 number of days is 0
```

```
> # 2 days 1 to 7
```

```
> # 3 days 8-14
```

```
> # 4 days 15+
```

```
>
```

```
> nmdta$mntlhlthOrd <- cut(nmdta$mntlhlth, breaks = c(0,1,8,15,30), labels = c(1,2,3,4), right = FALSE)
```

```
> #Used the table functions to verify the correctness of the ordinal variable
```

```
> table(nmdta$mntlhlthOrd)
```

1	2	3	4
399	226	44	58

```
> table(nmdta$mntlhlth)
```

0	1	2	3	4	5	6	7	8	10	12	14	15	16	18	20	21	25	27	30
399	34	62	37	29	39	6	19	2	35	3	4	22	1	2	21	2	9	1	23

## Task 2 Run Proportional Odds Model (Cumulative Logit)

- Run a proportional odds (cumulative ordered logit) regression of `mntlhlthOrd` on age (age), sex (sex; male is used as the reference category), race (race; white is used as the reference category), education (educ), and income (inc1k). Let us call this model Model 1 in the following tasks (hint: when white is used as the reference category, the white dummy needs to be dropped from the equation and the other two dummy variables for race have to be retained).
- Then run a second proportional odds (cumulative ordered logit) regression of mental health ( `mntlhlthOrd`) on sex (sex; male is used as the reference category), race (race; white is used as the reference category), and income (inc1k), with age and educ excluded from the explanatory variables. Let's call this model Model 2.

```
> #Task 2 Run Proportional Odds Model
```

```

> library (MASS)

> ordlogit.model1 <- polr(mnt1hlthOrd ~ age + male + other + black + educ +
inclk, data = nmdata, method = c("logistic"))

> summary(ordlogit.model1)
Call:
polr(formula = mnt1hlthOrd ~ age + male + other + black + educ +
      inclk, data = nmdata, method = c("logistic"))

Coefficients:
              Value          Std. Error      t value
age          -0.009002         0.005631      -1.5987
male         -0.499754         0.147603      -3.3858
other        -0.472610         0.284010      -1.6641
black        -0.575629         0.224130      -2.5683
educ          0.014756         0.028746       0.5133
inclk        -0.002473         0.002246      -1.1013

Intercepts:
      Value  Std. Error t value
1|2 -0.4317  0.4696     -0.9194
2|3  1.2236  0.4729      2.5872
3|4  1.8633  0.4793      3.8873

Residual Deviance: 1523.136
AIC: 1541.136
(23 observations deleted due to missingness)

> ordlogit.model2 <- polr(mnt1hlthOrd ~ male + other + black + inclk, data =
nmdata, method = c("logistic"))

> summary(ordlogit.model2)
Call:
polr(formula = mnt1hlthOrd ~ male + other + black + inclk, data = nmdata,
      method = c("logistic"))

Coefficients:
              Value          Std. Error      t value
male        -0.503079         0.147368      -3.414
other       -0.435115         0.281733      -1.544
black       -0.586781         0.223041      -2.631
inclk       -0.002645         0.002053      -1.288

Intercepts:
      Value  Std. Error t value
1|2 -0.2711  0.1360     -1.9939
2|3  1.3781  0.1498      9.1978
3|4  2.0173  0.1715     11.7595

Residual Deviance: 1526.105
AIC: 1540.105
(23 observations deleted due to missingness)

```

### Task 3 Test Hypothesis Using the Likelihood Ratio Test

- Test whether the effects of education and income are simultaneous equal to zero in Model 1 using the Wald test. Please also provide a brief interpretation of the results.
- Test whether the effects of race are zero in Model 2 using the LR test. Please also provide a brief interpretation of the results.

```
> #Task 3 Test Hypothesis Using the Likelihood Ratio Test
> #Wald Test
> library(car)

> waldtest = linearHypothesis(ordlogit.model1, c("age = 0", "educ = 0" ))

> waldtest
Linear hypothesis test

Hypothesis:
age = 0
educ = 0
```

```
Model 1: restricted model
Model 2: mnt1hlthOrd ~ age + male + other + black + educ + inclk
```

	Res.Df	Df	Chisq	Pr(>Chisq)
1	720			
2	718	2	2.9446	0.2294

**Wald Test Interpretation:** After completing the test, with a degree of freedom of 2 and a test statistic  $X^2 = 2.9446$ , we have a p-value of 0.2294, so we would fail to reject the null hypothesis at the 0.05 significance level. This would lead us to conclude there is no significant advantage of using the restricted model over the original model.

```
> #Test the same hypothesis using LR test
>
> lrTest = anova(ordlogit.model2, ordlogit.model1, test = "Chisq")

> print(lrTest)
Likelihood ratio tests of ordinal regression models
```

```
Response: mnt1hlthOrd
      Model               Resid.df  Resid.Dev   Test   Df LR stat  Pr(Chi)
1 male + other + black + inclk         720  1526.105
2 age + male + other + black + educ + inclk  718  1523.136  1 vs 2  2   2.968299 0.226695
```

**LR Test Interpretation:** After completing the test, with a degree of freedom of 2 and a test statistic  $X^2 = 2.9683$ , we have a p-value of 0.2267, so we would fail to reject the null hypothesis at the 0.05 significance level. This would lead us to conclude there is no significant advantage of using the second model (ordlogit.model2) over the original model (ordlogit.model1)

### Task 4 Test Hypothesis Using AIC and BIC

- Compare models 1-2 using AIC, and interpret the results.
- Compare models 1-2 using BIC, and interpret the results.

```
> #TASK 4 Test using AIC and BIC
> library(stats4)

> AIC(ordlogit.model1, ordlogit.model2)
```

	df	AIC
ordlogit.model1	9	1541.136
ordlogit.model2	7	1540.105

**AIC Interpretation:** From the table below, the second model will fit better because it has a smaller AIC. However, the difference in AICs between model1(1541.136) and model2(1540.105) is less than 2, so it is weak evidence that model2 has a significant advantage over the original model.

```
> BIC(ordlogit.model1, ordlogit.model2)
      df      BIC
ordlogit.model1  9 1582.437
ordlogit.model2  7 1572.227
```

**BIC Interpretation:** By looking strictly at the BIC numbers, the second model would fit the data better because it has the smaller BIC. When we check the difference between the two BIC values,  $1582.437 - 1572.227 = 10.21$ . Since this difference greater than 10, we would conclude there is strong evidence that the second model is the best-fitting model between the two.

### Task 5 Compare Models Using R-squared

- Compare models 1-2 using the McFadden's pseudo R-squared, and interpret the results.
- Compare models 1-2 using the Nagel's (Nagelkerke / Cragg & Uhler's) pseudo R-squared, and interpret the results.

```
> #TASK 5 Compare Using Pseudo R-Squared
> library(DescTools)

> PseudoR2(ordlogit.model1, c("McFadden", "Nagel"))
      McFadden      Nagelkerke
0.01542965    0.03666645

> PseudoR2(ordlogit.model2, c("McFadden", "Nagel"))
      McFadden      Nagelkerke
0.01351092    0.03217211
```

**Interpretation:** With both McFadden's pseudo- $R^2$  and Nagel's pseudo- $R^2$ , we would arrive at the same conclusion that the original model (ordlogit.model1) is the better fitting model because the pseudo- $R^2$  values are greater for the original model than the second model (ordlogit.model2).

### R-File

```
#source("/Users/burrisfaculty/Desktop/DSCode/SOC686/Shepherd_Lab07_SOC686.r",
echo=T, max.deparse.length=10000)
library(foreign)
```

```
#Open Log and read in data
setwd("/Users/burrisfaculty/Desktop/DSCode/SOC686")
sink("Shepherd_asgn07.log", split=T)
rm(list=ls(all=TRUE))
mygss <- read.dta("gsscsm7212teach.dta", convert.factor=F)
```

```
#MANAGE DATA AND RUN LOGIT
#SELECT DATA
useddta <- subset(mygss,
                  select=c(mntlh1th, age, sex, race, educ, inclk))
```

```

#Create dummy variables female (male = 0)
useddta$female <- as.numeric(useddta$sex==2)
useddta$male <- as.numeric(useddta$sex == 1)

#Create Binary Indicator Variables for Multi-Category Nomial Variables

useddta$white <- ifelse(useddta$race == 1, 1, 0)
useddta$black <- ifelse(useddta$race == 2, 1, 0)
useddta$other <- ifelse(useddta$race == 3, 1, 0)

nmdta <- useddta[complete.cases(useddta),] #no missing data

#summarize data
summary(nmdta$mntlhlth)
summary(nmdta$inclk)
summary(nmdta$age)
table(nmdta$female)
table(nmdta$white)

#Task 1 Create Ordinal Response Variable
# 1 number of days is 0
# 2 days 1 to 7
# 3 days 8-14
# 4 days 15+

nmdta$mntlhlthOrd <- cut(nmdta$mntlhlth, breaks = c(0,1,8,15,30), labels =
c(1,2,3,4), right = FALSE)
#Used the table functions to verify the correctness of the ordinal variable
table(nmdta$mntlhlthOrd)
table(nmdta$mntlhlth)

#Task 2 Run Proportional Odds Model
library(MASS)
ordlogit.model1 <- polr(mntlhlthOrd ~ age + male + other + black + educ +
inclk, data = nmdta, method = c("logistic"))
summary(ordlogit.model1)
ordlogit.model2 <- polr(mntlhlthOrd ~ male + other + black + inclk, data =
nmdta, method = c("logistic"))
summary(ordlogit.model2)

#Task 3 Test Hypothesis Using the Likelihood Ratio Test
#Wald Test
library(car)
waldtest = linearHypothesis(ordlogit.model1, c("age = 0", "educ = 0" ))
waldtest

#Test the same hypothesis using LR test

lrTest = anova(ordlogit.model2, ordlogit.model1, test = "Chisq")
print(lrTest)

#TASK 4 Test using AIC and BIC
library(stats4)
AIC(ordlogit.model1,ordlogit.model2)

BIC(ordlogit.model1,ordlogit.model2)

```

```
#TASK 5 Compare Using Pseudo R-Squared
library(DescTools)

PseudoR2(ordlogit.model1, c("McFadden", "Nagel"))
PseudoR2(ordlogit.model2, c("McFadden", "Nagel"))

#Close log
save(nmdta, file = "Assignment_07.rdata")
sink()
```

## Log

```
> rm(list=ls(all=TRUE))

> mygss <- read.dta("gsscum7212teach.dta", convert.factor=F)

> #MANAGE DATA AND RUN LOGIT
> #SELECT DATA
> useddta <- subset(mygss,
+                   select=c(mntlhlth, age, sex, race, educ, inclk))

> #Create dummy variables female (male = 0)
> useddta$female <- as.numeric(useddta$sex==2)

> useddta$male <- as.numeric(useddta$sex == 1)

> #Create Binary Indicator Variables for Multi-Category Nomial Variables
>
> useddta$white <- ifelse(useddta$race == 1, 1, 0)

> useddta$black <- ifelse(useddta$race == 2, 1, 0)

> useddta$other <- ifelse(useddta$race == 3, 1, 0)

> nmdta <- useddta[complete.cases(useddta),] #no missing data

> #summarize data
> summary(nmdta$mntlhlth)
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
0.000  0.000   0.000   3.991  5.000  30.000

> summary(nmdta$inclk)
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
0.245  15.921  28.157  39.404  48.475 155.140

> summary(nmdta$age)
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
18.00  31.00   42.00   41.81  51.00   84.00

> table(nmdta$female)

 0    1
381 369

> table(nmdta$white)
```

```
0 1
160 590
```

```
> #Task 1 Create Ordinal Response Variable
> # 1 number of days is 0
> # 2 days 1 to 7
> # 3 days 8-14
> # 4 days 15+
>
> nmdta$mntlhlthOrd <- cut(nmdta$mntlhlth, breaks = c(0,1,8,15,30), labels =
c(1,2,3,4), right = FALSE)

> #Used the table functions to verify the correctness of the ordinal variable
> table(nmdta$mntlhlthOrd)
```

```
1 2 3 4
399 226 44 58
```

```
> table(nmdta$mntlhlth)
```

```
0 1 2 3 4 5 6 7 8 10 12 14 15 16 18 20 21 25 27
30
399 34 62 37 29 39 6 19 2 35 3 4 22 1 2 21 2 9 1
23
```

```
> #Task 2 Run Proportional Odds Model
> library (MASS)

> ordlogit.model1 <- polr(mntlhlthOrd ~ age + male + other + black + educ +
inclk, data = nmdta, method = c("logistic"))

> summary(ordlogit.model1)
Call:
polr(formula = mntlhlthOrd ~ age + male + other + black + educ +
inclk, data = nmdta, method = c("logistic"))
```

Coefficients:

	Value	Std. Error	t value
age	-0.009002	0.005631	-1.5987
male	-0.499754	0.147603	-3.3858
other	-0.472610	0.284010	-1.6641
black	-0.575629	0.224130	-2.5683
educ	0.014756	0.028746	0.5133
inclk	-0.002473	0.002246	-1.1013

Intercepts:

	Value	Std. Error	t value
1 2	-0.4317	0.4696	-0.9194
2 3	1.2236	0.4729	2.5872
3 4	1.8633	0.4793	3.8873

Residual Deviance: 1523.136

AIC: 1541.136

(23 observations deleted due to missingness)



```
> ordlogit.model2 <- polr(mntllhthOrd ~ male + other + black + inclk, data =
nmdta, method = c("logistic"))
```

```
> summary(ordlogit.model2)
```

Call:

```
polr(formula = mntllhthOrd ~ male + other + black + inclk, data = nmdta,
method = c("logistic"))
```

Coefficients:

	Value	Std. Error	t value
male	-0.503079	0.147368	-3.414
other	-0.435115	0.281733	-1.544
black	-0.586781	0.223041	-2.631
inclk	-0.002645	0.002053	-1.288

Intercepts:

	Value	Std. Error	t value
1 2	-0.2711	0.1360	-1.9939
2 3	1.3781	0.1498	9.1978
3 4	2.0173	0.1715	11.7595

Residual Deviance: 1526.105

AIC: 1540.105

(23 observations deleted due to missingness)

```
> #Task 3 Test Hypothesis Using the Likelihood Ratio Test
```

```
> #Wald Test
```

```
> library(car)
```

```
> waldtest = linearHypothesis(ordlogit.model1, c("age = 0", "educ = 0" ))
```

```
> waldtest
```

Linear hypothesis test

Hypothesis:

age = 0

educ = 0

Model 1: restricted model

Model 2: mntllhthOrd ~ age + male + other + black + educ + inclk

	Res.Df	Df	Chisq	Pr(>Chisq)
1	720			
2	718	2	2.9446	0.2294

```
> #Test the same hypothesis using LR test
```

```
>
```

```
> lrTest = anova(ordlogit.model2, ordlogit.model1, test = "Chisq")
```

```
> print(lrTest)
```

Likelihood ratio tests of ordinal regression models

Response: mntllhthOrd

	LR stat.	Pr(Chi)	Model	Resid. df	Resid. Dev	Test	Df
1			male + other + black + inclk	720	1526.105		

```
2 age + male + other + black + educ + inclk      718    1523.136 1 vs 2      2
2.968299 0.226695
```

```
> #TASK 4 Test using AIC and BIC
> library(stats4)
```

```
> AIC(ordlogit.model1,ordlogit.model2)
      df      AIC
ordlogit.model1  9    1541.136
ordlogit.model2  7    1540.105
```

```
> BIC(ordlogit.model1,ordlogit.model2)
      df      BIC
ordlogit.model1  9    1582.437
ordlogit.model2  7    1572.227
```

```
> #TASK 5 Compare Using Pseudo R-Squared
> library(DescTools)
```

```
> PseudoR2(ordlogit.model1, c("McFadden", "Nagel"))
      McFadden  Nagelkerke
0.01542965    0.03666645
```

```
> PseudoR2(ordlogit.model2, c("McFadden", "Nagel"))
      McFadden  Nagelkerke
0.01351092    0.03217211
```

```
> #Close log
> save(nmdta, file = "Assignment_07.rdata")

> sink()
```