

Cálculo de Programas

Resolução - Ficha 08

Eduardo Freitas Fernandes

2026

Exercício 1

$$\begin{array}{ll} T \textit{id} = \textit{id} & T(f \cdot g) = (Tf) \cdot (Tg) \\ \equiv \{ \text{(F1)} \} & \equiv \{ \text{(F1)} \} \\ \textit{id} \times \textit{id} = \textit{id} & (f \cdot g) \times (f \cdot g) = (f \times f) \cdot (g \times g) \\ \equiv \{ \text{def-}\times \} & \equiv \{ \text{def-}\times \text{(twice)} \} \\ \langle \textit{id} \cdot \pi_1, \textit{id} \cdot \pi_2 \rangle = \textit{id} & \langle f \cdot g \cdot \pi_1, f \cdot g \cdot \pi_2 \rangle = \langle f \cdot \pi_1, f \cdot \pi_2 \rangle \cdot (g \times g) \\ \equiv \{ \text{natural-id (twice)} \} & \equiv \{ \text{fusão-}\times \} \\ \langle \pi_1, \pi_2 \rangle = \textit{id} & \langle f \cdot g \cdot \pi_1, f \cdot g \cdot \pi_2 \rangle = \langle f \cdot \pi_1 \cdot (g \times g), f \cdot \pi_2 \cdot (g \times g) \rangle \\ \equiv \{ \text{reflexão-}\times \} & \equiv \{ \text{natural-}\pi_1, \text{natural-}\pi_2 \} \\ \textit{id} = \textit{id} & \langle f \cdot g \cdot \pi_1, f \cdot g \cdot \pi_2 \rangle = \langle f \cdot g \cdot \pi_1, f \cdot g \cdot \pi_2 \rangle \\ \square & \square \end{array}$$

$$\begin{array}{l} \mu \cdot (T u) = \mu \cdot u \\ \equiv \{ \text{def. } \mu, \text{def. } T u \} \\ (\pi_1 \times \pi_2) \cdot (\langle \textit{id}, \textit{id} \rangle \times \langle \textit{id}, \textit{id} \rangle) = (\pi_1 \times \pi_2) \cdot \langle \textit{id}, \textit{id} \rangle \\ \equiv \{ \text{def-}\times, \text{absorção-}\times, \text{natural-id} \} \\ (\pi_1 \times \pi_2) \cdot \langle \langle \textit{id}, \textit{id} \rangle \cdot \pi_1 \cdot (\langle \textit{id}, \textit{id} \rangle \cdot \pi_2) = \langle \pi_1, \pi_2 \rangle, \cdot \rangle \\ \equiv \{ \text{absorção-}\times, \text{reflexão-}\times \} \\ \langle \pi_1 \cdot \langle \textit{id}, \textit{id} \rangle \cdot \pi_1, \pi_2 \cdot \langle \textit{id}, \textit{id} \rangle \cdot \pi_2 \rangle = \langle \pi_1, \pi_2 \rangle \\ \equiv \{ \text{cancelamento-}\times \} \\ \langle \pi_1, \pi_2 \rangle = \langle \pi_1, \pi_2 \rangle \\ \square \end{array}$$

Exercício 2

$$\begin{array}{l} \textit{loop}(a, b) = (2 + b, 2 - a) \\ \equiv \{ \text{pointfree} \} \\ \textit{loop} = ((2+) \times (2-)) \cdot \textit{swap} \\ \langle f, g \rangle = \text{for } \textit{loop} \ 4, -2 \end{array}$$

$$\begin{aligned}
&\equiv \{ \text{def. for} \} \\
&\quad \langle f, g \rangle = \langle \llbracket \underline{4}, \underline{-2} \rrbracket, \text{loop} \rangle \\
&\equiv \{ \text{def. loop} \} \\
&\quad \langle f, g \rangle = \langle \llbracket \underline{4}, \underline{-2} \rrbracket, (2+) \times (2-) \cdot \text{swap} \rangle \\
&\equiv \{ \text{def-}\times, \text{fusão-}\times \} \\
&\quad \langle f, g \rangle = \langle \llbracket \underline{4}, \underline{-2} \rrbracket, \langle (2+) \cdot \pi_1 \cdot \text{swap}, (2-) \cdot \pi_2 \cdot \text{swap} \rangle \rangle \\
&\equiv \{ \text{lei da troca} \} \\
&\quad \langle f, g \rangle = \langle \llbracket \underline{4}, (2+) \cdot \pi_1 \cdot \text{swap} \rrbracket, \llbracket \underline{-2}, (2-) \cdot \pi_2 \cdot \text{swap} \rrbracket \rangle \\
&\equiv \{ \text{fokkinga} \} \\
&\quad \begin{cases} f \cdot \llbracket \underline{0}, \text{succ} \rrbracket = \llbracket \underline{4}, (2+) \cdot \pi_1 \cdot \text{swap} \rrbracket \cdot (\text{id} + \langle f, g \rangle) \\ g \cdot \llbracket \underline{0}, \text{succ} \rrbracket = \llbracket \underline{-2}, (2-) \cdot \pi_2 \cdot \text{swap} \rrbracket \cdot (\text{id} + \langle f, g \rangle) \end{cases} \\
&\equiv \{ \text{fusão-}+, \text{absorção-}+, \text{eq-}+ \} \\
&\quad \begin{cases} f \cdot \underline{0} = \underline{4} \\ f \cdot \text{succ} = (2+) \cdot \pi_1 \cdot \text{swap} \\ g \cdot \underline{0} = \underline{-2} \\ g \cdot \text{succ} = (2-) \cdot \pi_2 \cdot \text{swap} \end{cases} \\
&\equiv \{ \text{pointwise} \} \\
&\quad \begin{cases} f \ 0 = 4 \\ f \ (n+1) = 2 + g \ n \\ g \ 0 = -2 \\ g \ (n+1) = 2 - f \ n \end{cases}
\end{aligned}$$

Exercício 3

$$\begin{aligned}
&\langle \langle f, g \rangle, j \rangle = \langle \langle \langle h, k \rangle, l \rangle \rangle \\
&\equiv \{ \text{fokkinga} \} \\
&\quad \begin{cases} \langle f, g \rangle \cdot \text{in} = \langle h, k \rangle \cdot F \langle \langle f, g \rangle, h \rangle \\ j \cdot \text{in} = l \cdot F \langle \langle f, g \rangle, h \rangle \end{cases} \\
&\equiv \{ \text{fusão-}\times (\text{twice}) \} \\
&\quad \begin{cases} \langle f \cdot \text{in}, g \cdot \text{in} \rangle = \langle h \cdot F \langle \langle f, g \rangle, j \rangle, k \cdot F \langle \langle f, g \rangle, j \rangle \rangle \\ j \cdot \text{in} = l \cdot F \langle \langle f, g \rangle, j \rangle \end{cases} \\
&\equiv \{ \text{eq-}\times \} \\
&\quad \begin{cases} f \cdot \text{in} = h \cdot F \langle \langle f, g \rangle, j \rangle \\ g \cdot \text{in} = k \cdot F \langle \langle f, g \rangle, j \rangle \\ j \cdot \text{in} = l \cdot F \langle \langle f, g \rangle, j \rangle \end{cases}
\end{aligned}$$

Exercício 4

$$\begin{aligned}
&\begin{cases} \text{impar } 0 = \text{false} \\ \text{impar } (n+1) = \text{par } n \\ \text{par } 0 = \text{true} \\ \text{par } (n+1) = \text{impar } n \end{cases} \\
&\equiv \{ \text{pointfree} \}
\end{aligned}$$

$$\begin{aligned}
& \left\{ \begin{array}{l} \text{impar} \cdot \underline{0} = \underline{\text{false}} \\ \text{impar} \cdot \text{succ} = \text{par} \\ \text{par} \cdot \underline{0} = \underline{\text{true}} \\ \text{par} \cdot \text{succ} = \text{impar} \end{array} \right. \\
\equiv & \quad \{ \text{eq-+} \} \\
& \left\{ \begin{array}{l} [\text{impar} \cdot \underline{0}, \text{impar} \cdot \text{succ}] = [\underline{\text{false}}, \text{par}] \\ [\text{par} \cdot \underline{0}, \text{par} \cdot \text{succ}] = [\underline{\text{true}}, \text{impar}] \end{array} \right. \\
\equiv & \quad \{ \text{fusão-+}, \text{cancelamento-+ (twice)} \} \\
& \left\{ \begin{array}{l} \text{impar} \cdot \text{in} = [\underline{\text{false}}, \pi_2 \cdot \langle \text{impar}, \text{par} \rangle] \\ \text{par} \cdot \text{in} = [\underline{\text{true}}, \pi_1 \cdot \langle \text{impar}, \text{par} \rangle] \end{array} \right. \\
\equiv & \quad \{ \text{natural-id (twice)}, \text{absorção-+} \} \\
& \left\{ \begin{array}{l} \text{impar} \cdot \text{in} = [\underline{\text{false}}, \pi_2] \cdot (\text{id} + \langle \text{impar}, \text{par} \rangle) \\ \text{par} \cdot \text{in} = [\underline{\text{true}}, \pi_1] \cdot (\text{id} + \langle \text{impar}, \text{par} \rangle) \end{array} \right.
\end{aligned}$$

Podemos então concluir que $h = [\underline{\text{false}}, \pi_2]$ e $k = [\underline{\text{true}}, \pi_1]$.

$$\begin{aligned}
& \langle \text{impar}, \text{par} \rangle = \text{for swap false, true} \\
\equiv & \quad \{ \text{def. for} \} \\
& \langle \text{impar}, \text{par} \rangle = \langle \langle [\underline{\text{false}}, \underline{\text{true}}], \text{swap} \rangle \rangle \\
\equiv & \quad \{ \text{def. swap} \} \\
& \langle \text{impar}, \text{par} \rangle = \langle \langle [\underline{\text{false}}, \underline{\text{true}}], \langle \pi_2, \pi_1 \rangle \rangle \rangle \\
\equiv & \quad \{ \text{lei da troca} \} \\
& \langle \text{impar}, \text{par} \rangle = \langle \langle [\underline{\text{false}}, \pi_2], [\underline{\text{true}}, \pi_1] \rangle \rangle \\
\equiv & \quad \{ \text{fokkinga} \} \\
& \left\{ \begin{array}{l} \text{impar} \cdot [\underline{0}, \text{succ}] = [\underline{\text{false}}, \pi_2] \cdot (\text{id} + \langle \text{impar}, \text{par} \rangle) \\ \text{par} \cdot [\underline{0}, \text{succ}] = [\underline{\text{true}}, \pi_1] \cdot (\text{id} + \langle \text{impar}, \text{par} \rangle) \end{array} \right. \\
\equiv & \quad \{ \text{fusão-+}, \text{absorção-+} \} \\
& \left\{ \begin{array}{l} [\text{impar} \cdot \underline{0}, \text{impar} \cdot \text{succ}] = [\underline{\text{false}}, \pi_2 \cdot \langle \text{impar}, \text{par} \rangle] \\ [\text{par} \cdot \underline{0}, \text{par} \cdot \text{succ}] = [\underline{\text{true}}, \pi_1 \cdot \langle \text{impar}, \text{par} \rangle] \end{array} \right. \\
\equiv & \quad \{ \text{cancelamento-}\times \text{ (twice)}, \text{eq-+}, \text{pointwise} \} \\
& \left\{ \begin{array}{l} \text{impar } 0 = \text{false} \\ \text{impar } (n + 1) = \text{par } n \\ \text{par } 0 = \text{true} \\ \text{par } (n + 1) = \text{impar } n \end{array} \right.
\end{aligned}$$

Exercício 5

$$\begin{aligned}
& \left\{ \begin{array}{l} \text{insg } 0 = [] \\ \text{insg } (n + 1) = \text{fsuc } n : \text{insg } n \\ \text{fsuc } 0 = 1 \\ \text{fsuc } (n + 1) = \text{fsuc } n + 1 \end{array} \right. \\
\equiv & \quad \{ \text{pointfree} \}
\end{aligned}$$

$$\begin{aligned}
& \left\{ \begin{array}{l} \text{insg} \cdot \underline{0} = \text{nil} \\ \text{insg} \cdot \text{succ} = \text{cons} \cdot \langle \text{fsuc}, \text{insg} \rangle \\ \text{fsuc} \cdot \underline{0} = \underline{1} \\ \text{fsuc} \cdot \text{succ } 0 = (1+) \cdot \text{fsuc} \end{array} \right. \\
\equiv & \quad \{ \text{cancelamento-}\times \} \\
& \left\{ \begin{array}{l} \text{insg} \cdot \underline{0} = \text{nil} \\ \text{insg} \cdot \text{succ} = \text{cons} \cdot \langle \text{fsuc}, \text{insg} \rangle \\ \text{fsuc} \cdot \underline{0} = \underline{1} \\ \text{fsuc} \cdot \text{succ } 0 = (1+) \cdot \pi_1 \cdot \langle \text{fsuc}, \text{insg} \rangle \end{array} \right. \\
\equiv & \quad \{ \text{eq-+}, \text{fusão-+}, \text{absorção-+} \} \\
& \left\{ \begin{array}{l} \text{insg} \cdot \text{in} = [\text{nil}, \text{cons}] \cdot (\text{id} + \langle \text{fsuc}, \text{insg} \rangle) \\ \text{fsuc} \cdot \text{in} = [\underline{1}, (1+) \cdot \pi_1] \cdot (\text{id} + \langle \text{fsuc}, \text{insg} \rangle) \end{array} \right. \\
\equiv & \quad \{ \text{fokkinga} \} \\
& \langle \text{fsuc}, \text{insg} \rangle = \langle \langle \text{in}, [\underline{1}, (1+) \cdot \pi_1] \rangle \rangle
\end{aligned}$$

$$\begin{aligned}
& \text{insgfor} = \text{for } \langle (1+) \cdot \pi_1, \text{cons} \rangle \langle \underline{1}, \text{nil} \rangle \\
\equiv & \quad \{ \text{def. for} \} \\
& \text{insgfor} = \langle \langle \underline{1}, \text{nil} \rangle, \langle (1+) \cdot \pi_1, \text{cons} \rangle \rangle \\
\equiv & \quad \{ \text{lei da troca} \} \\
& \text{insgfor} = \langle \langle [\underline{1}, (1+) \cdot \pi_1], [\text{nil}, \text{cons}] \rangle \rangle \\
\equiv & \quad \{ \} \\
& \langle f, \text{insg} \rangle = \langle \langle [\underline{1}, (1+) \cdot \pi_1], [\text{nil}, \text{cons}] \rangle \rangle \\
\equiv & \quad \{ \text{fokkinga} \} \\
& \left\{ \begin{array}{l} f \cdot [\underline{0}, \text{succ}] = [\underline{1}, (1+) \cdot \pi_1] \cdot F \langle f, \text{insg} \rangle \\ \text{insg} \cdot [\underline{0}, \text{succ}] = [\text{nil}, \text{cons}] \cdot F \langle f, \text{insg} \rangle \end{array} \right. \\
\equiv & \quad \{ \text{def. functor } \mathbb{N}_0, \text{fusão-+}, \text{absorção-+}, \text{eq-+} \} \\
& \left\{ \begin{array}{l} f \cdot \underline{0} = \underline{1} \\ f \cdot \text{succ} = (1+) \cdot \pi_1 \cdot \langle f, \text{insg} \rangle \\ \text{insg} \cdot \underline{0} = \text{nil} \\ \text{insg} \cdot \text{succ} = \text{cons} \cdot \langle f, \text{insg} \rangle \end{array} \right. \\
\equiv & \quad \{ \text{pointwise} \} \\
& \left\{ \begin{array}{l} f \ 0 = 1 \\ f \ (n+1) = 1 + f \ n \\ \text{insg} \ 0 = [] \\ \text{insg} \ (n+1) = f \ n : \text{insg} \ n \end{array} \right.
\end{aligned}$$

Exercício 6

$$\left\{ \begin{array}{l} f_1 [] = [] \\ f_1 (h : t) = h : f_2 t \\ f_2 [] = [] \\ f_2 (h : t) = f_1 t \end{array} \right.$$

$$\begin{aligned}
&\equiv \{ \text{def. comp, pointfree} \} \\
&\quad \left\{ \begin{array}{l} f_1 \cdot \text{nil} = \text{nil} \\ f_1 \cdot \text{cons} = \text{cons} \cdot (\text{id} \times f_2) \\ f_2 \cdot \text{nil} = \text{nil} \\ f_2 \cdot \text{cons} = f_1 \cdot \pi_2 \end{array} \right. \\
&\equiv \{ \text{eq-+}, \text{fusão-+} \} \\
&\quad \left\{ \begin{array}{l} f_1 \cdot [\text{nil}, \text{cons}] = [\text{nil}, \text{cons} \cdot (\text{id} \times f_2)] \\ f_2 \cdot [\text{nil}, \text{cons}] = [\text{nil}, f_1 \cdot \pi_2] \end{array} \right. \\
&\equiv \{ \text{def. in, natural-id (twice), natural-}\pi_2, \text{cancelamento-}\times \} \\
&\quad \left\{ \begin{array}{l} f_1 \cdot \text{in} = [\text{nil} \cdot \text{id}, \text{cons} \cdot (\text{id} \times \pi_2 \cdot \langle f_1, f_2 \rangle)] \\ f_2 \cdot \text{in} = [\text{nil} \cdot \text{id}, \pi_2 \cdot (\text{id} \times \pi_1 \cdot \langle f_1, f_2 \rangle)] \end{array} \right. \\
&\equiv \{ \text{functor-}\times \} \\
&\quad \left\{ \begin{array}{l} f_1 \cdot \text{in} = [\text{nil} \cdot \text{id}, \text{cons} \cdot (\text{id} \times \pi_2) \cdot (\text{id} \times \langle f_1, f_2 \rangle)] \\ f_2 \cdot \text{in} = [\text{nil} \cdot \text{id}, \pi_2 \cdot (\text{id} \times \pi_1) \cdot (\text{id} \times \langle f_1, f_2 \rangle)] \end{array} \right. \\
&\equiv \{ \text{absorção-+}, \text{def. functor de listas, natural-}\pi_2 \} \\
&\quad \left\{ \begin{array}{l} f_1 \cdot \text{in} = [\text{nil}, \text{cons} \cdot (\text{id} \times \pi_2)] \cdot F \langle f_1, f_2 \rangle \\ f_2 \cdot \text{in} = [\text{nil}, \pi_1 \cdot \pi_2] \cdot F \langle f_1, f_2 \rangle \end{array} \right. \\
&\equiv \{ \text{fokkinga} \} \\
&\quad \langle f_1, f_2 \rangle = \langle \langle \text{in} \cdot (\text{id} \times \pi_2), [\text{nil}, \pi_1 \cdot \pi_2] \rangle \rangle
\end{aligned}$$





Diagrama do gene do catamorfismo:



A função f_1 seleciona os elementos de uma lista nas posições pares, e a função f_2 seleciona os elementos de uma lista nas posições ímpares.

Exercício 7

$$\begin{aligned}
 H(g \cdot h) &= (H g) \cdot (H h) \\
 \equiv \quad &\{ \text{def. functor } H(3^*) \} \\
 F(g \cdot h) + G(g \cdot h) &= (F g + G g) \cdot (F h + G h) \\
 \equiv \quad &\{ \text{def. functor } F(3^*), \text{ def. functor } G(3^*) \} \\
 id + (g \cdot h) &= (id + g) \cdot (id + h) \\
 \equiv \quad &\{ \text{natural-id, functor-+} \} \\
 (id \cdot id) + (g \cdot h) &= (id \cdot id) + (g \cdot h) \\
 \square
 \end{aligned}$$

$$\begin{aligned}
 K(g \cdot h) &= (K g) \cdot (K h) \\
 \equiv \quad &\{ \text{def. functor } K \} \\
 G(g \cdot h) \times F(g \cdot h) &= (G g \times F g) \cdot (G h \times F h) \\
 \equiv \quad &\{ \text{def. functor } F(3^*), \text{ def. functor } G(3^*) \} \\
 (g \cdot h) \times id &= (g \times id) \cdot (h \times id) \\
 \equiv \quad &\{ \text{natural-id, functor-}\times \} \\
 (g \cdot h) \times (id \cdot id) &= (g \cdot h) \times (id \cdot id) \\
 \square
 \end{aligned}$$

$$\begin{aligned}
 H id &= id \\
 \equiv \quad &\{ \text{def. functor } H \} \\
 F id + G id &= id \\
 \equiv \quad &\{ \text{def. functor } F, \text{ def. functor } G \} \\
 id + id &= id \\
 \equiv \quad &\{ \text{functor-id-+} \} \\
 id &= id \\
 \square
 \end{aligned}$$

$$\begin{aligned}
 K id &= id \\
 \equiv \quad &\{ \text{def. functor } K \} \\
 G id \times F id &= id \\
 \equiv \quad &\{ \text{def. functor } F, \text{ def. functor } G \} \\
 id \times id &= id \\
 \equiv \quad &\{ \text{functor-id-}\times \} \\
 id &= id \\
 \square
 \end{aligned}$$

Exercício 8

$$\begin{aligned}
 H id &= id \\
 \equiv \quad &\{ \text{def. functor } H \} \\
 (F \cdot G) id &= id \\
 \equiv \quad &\{ \text{composição de funtores} \} \\
 F(G id) &= id \\
 \equiv \quad &\{ \text{functor-id-G} \} \\
 F id &= id \\
 \equiv \quad &\{ \text{functor-id-F} \} \\
 id &= id \\
 \square
 \end{aligned}$$

$$\begin{aligned}
 H(f \cdot g) &= (H f) \cdot (H g) \\
 \equiv \quad &\{ \text{def. functor } H \} \\
 (F \cdot G)(f \cdot g) &= ((F \cdot G) f) \cdot ((F \cdot G) g) \\
 \equiv \quad &\{ \text{composição de funtores} \} \\
 F(G(f \cdot g)) &= (F(G f)) \cdot (F(G g)) \\
 \equiv \quad &\{ \text{Functor-F, Functor-G} \} \\
 F(G f \cdot G g) &= F(G f \cdot G g) \\
 \square
 \end{aligned}$$

Exercício 9