



planetmath.org

Math for the people, by the people.

properties of X -integrable processes

Canonical name	PropertiesOfXintegrableProcesses
Date of creation	2013-03-22 18:40:59
Last modified on	2013-03-22 18:40:59
Owner	gel (22282)
Last modified by	gel (22282)
Numerical id	5
Author	gel (22282)
Entry type	Theorem
Classification	msc 60H10
Classification	msc 60G07
Classification	msc 60H05
Related topic	StochasticIntegration

Let X be a semimartingale. Then a predictable process ξ is X -integrable if the stochastic integral $\int \xi dX$ is defined, which is equivalent to the set

$$\left\{ \int_0^t \alpha dX : |\alpha| \leq |\xi| \text{ is predictable} \right\}$$

being bounded in probability, for each $t > 0$. We list some properties of X -integrable processes.

1. Every locally bounded predictable process is X -integrable.
2. The X -integrable processes are closed under linear combinations. That is, if α, β are X -integrable and $\lambda, \mu \in \mathbb{R}$, then $\lambda\alpha + \mu\beta$ is X -integrable.
3. If $|\alpha| \leq |\beta|$ are predictable processes and β is X -integrable, then so is α .
4. A process is X -integrable if it is locally X -integrable. That is, if there are stopping times τ_n almost surely increasing to infinity and such that $1_{\{t \leq \tau_n\}} \xi_t$ is X -integrable, then ξ is X -integrable.